

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 November 2003 (13.11.2003)

PCT

(10) International Publication Number  
**WO 03/092612 A2**

- (51) International Patent Classification<sup>7</sup>: **A61K**
- (21) International Application Number: PCT/US03/13672
- (22) International Filing Date: 2 May 2003 (02.05.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/377,503 2 May 2002 (02.05.2002) US
- (71) Applicant (*for all designated States except US*): **UNIVERSITY OF ROCHESTER** [US/US]; 601 Elmwood Avenue, Box 706, Rochester, NY 14642 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): **KYRKANIDES, Stephanos** [GR/US]; 1963 Harris Road, Penfield, NY 14526 (US).
- (74) Agents: **HUIZENGA, David, E.** et al.; Needle & Rosenberg, P.C., The Candler Building, 127 Peachtree Street, Atlanta, GA 30303-1811 (US).

**Published:**

— *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **VECTORS HAVING BOTH ISOFORMS OF  $\beta$ -HEXOSAMINIDASE**

(57) Abstract: Disclosed are compositions and methods related to nucleic acid constructs containing a HexB encoding element and a HexA encoding element. These constructs can be used in the treatment of Tay-Sachs and Sandoff disease.

WO 03/092612 A2

**VECTORS HAVING BOTH ISOFORMS OF  $\beta$ -HEXOSAMINIDASE****I. ACKNOWLEDGMENTS**

[01] This application claims priority to United States Provisional Application No. 60/377,503 filed on May 2, 2003 for Vectors Having Both Isoforms of  $\beta$ -hexosamidase.

5 This application is herein incorporated by reference in its entirety.

**II. BACKGROUND OF THE INVENTION**

[02] Lysosomal storage disorders are disorders that typically arise from the aberrant or non-existent proteins involved in degradation function within the lysosomes. This causes a decrease in the lysosomal activity, which in turn causes an accumulation of  
10 unwanted materials in the cell. These unwanted materials can cause severe cellular toxicity and can impair, for example, neuronal function. These diseases severely impair the quality of life of those who have them, and can even result in death. Two diseases, Tay-Sachs and Sandoffs, are related to the functional impairment of the lysosomal protein  $\beta$ -hexosaminidase.  $\beta$ -hexosaminidase is a hetero or homo dimer made up of two subunits  
15 arising from two separate genes, HexA and HexB. Mutation of the HexA gene, causing functional problems with the HEX- $\alpha$  (HexA/HexB) polypeptide, results in Tay-Sachs disease, whereas mutation of the HexB gene, causing functional problems in the HEX- $\alpha$  (HexA/HexB) and HEX- $\beta$  (HexB/HexB) polypeptides, results in Sandhoff's disease. Clinically, it is not uncommon for patients to display only mild features at infancy, but due  
20 to increasing lysosomal storage over time, progress to severe forms of the disease by adolescence.

[03] Current treatments include bone marrow transplantation, which has been employed in some cases of individuals during childhood but with modest outcomes. A significant problem with the bone marrow transplantation approach is that it may address  
25 the lack of specific metabolic activity in peripheral tissues, but due to the presence of the blood-brain-barrier it fails to avert disease progression in the central nervous system. Hence patients often continue to clinically deteriorate due to central nervous system involvement with subsequent development of neurodegeneration, blindness, mental retardation, paralysis and dementia.

30 [04] Enzyme replacement strategies targeting peripheral and central nervous system tissues utilizing gene therapy is a logical approach for treating inherited metabolic disorders. In a study by Akli *et al.* (1996), the authors report successful restoration of  $\beta$ -

hexosaminidase in fibroblasts derived from patients with *HexA* deficiency via adenoviral-mediated gene transfer *in vitro*. Likewise a *HexA* transgene and a *HexB* transgene was successfully introduced into neural progenitor cells utilizing retroviral vectors (Lacorazza *et al.*, "Expression of human beta-hexosaminidase alpha-subunit gene (the gene defect of Tay-Sachs disease) in mouse brains upon engraftment of transduced progenitor cells". Nat Med 2(4):424-9 (1996 Apr).

[05] Disclosed herein are vectors and methods which solve the problems associated with enzyme replacement therapies directed to  $\beta$ -hexosaminidase deficiencies.

### III. SUMMARY OF THE INVENTION

10 [06] In accordance with the purposes of this invention, as embodied and broadly described herein, this invention, in one aspect, relates to vector constructs that comprise sequence encoding the HEX- $\beta$  polypeptide. Also disclosed are vector constructs comprising sequence encoding the HEX- $\beta$  and the HEX- $\alpha$  polypeptides. Also disclosed are vectors for perinatal gene delivery, including delivery of HEX- $\alpha$  and HEX- $\beta$ , which can be  
15 used for inherited lysosomal disorders such as Tay-Sachs and Sandoffs disease.

[07] Additional advantages of the invention will be set forth in part in the description which follows, and in part will be obvious from the description, or may be learned by practice of the invention. The advantages of the invention will be realized and attained by means of the elements and combinations particularly pointed out in the  
20 appended claims. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

### IV. BRIEF DESCRIPTION OF THE DRAWINGS

[08] The accompanying drawings, which are incorporated in and constitute a part  
25 of this specification, illustrate several embodiments of the invention and together with the description, serve to explain the principles of the invention.

[09] Figure 1 shows that HEX*lacZ* encodes for both isoforms of human  $\beta$ -hexosaminidase, HexA & HexB. Figure 1(A) shows pHEX*lacZ* vector. BHK<sup>Hex*lacZ*</sup> are developed by stable Hex*lacZ* transduction. Figure 3(B) shows cells stain positively by X-gal histochemistry. Figure 3(C) shows HexA & HexB mRNA is detected by RT-PCR in  
30 total RNA extracts. Figure 3(D<sub>1</sub>) shows human HEXA & figure 3(E<sub>1</sub>) shows human HEXB proteins are detected in BHK<sup>Hex*lacZ*</sup> by imunocytochemistry. Figure 3(F<sub>1</sub>) shows HEXA &

HEXA+HEXB activity is measured by 4MUGS & 4MUG fluorometry, respectively. Figure (G)  $\beta$ -hexosaminidase detection by Fast Garnet histochemistry. (D<sub>2</sub>, E<sub>2</sub>, G<sub>2</sub> are controls for D<sub>1</sub>, E<sub>1</sub>, G<sub>1</sub>, respectively).

[10] Figure 2 shows that the  $\beta$ -Hex therapeutic gene cross-corrects. An important property of the  $\beta$ -Hex transgene is the products hHEXA & hHEXB have the ability to cross-correct, specifically, to be released extracellularly and then to be absorbed via paracrine pathways by other cells whereby they contribute to  $\beta$ -hexosaminidase activity. For this purpose, BHK<sup>HexlacZ</sup> cells were cultured and the supernatant was collected (conditioned medium), filtered (.45mm) and applied on normal mouse kidney fibroblasts in culture. Forty-eight hours later, the cells were washed thoroughly with phosphate buffered saline, and briefly treated with a trypsin solution to remove extracellular proteins from the cell surfaces. Following trypsin inactivation with Tris/EDTA buffer, the cells were fixed with 4% paraformaldehyde solution and processed by Fast Garnet histochemistry for  $\beta$ -hexosaminidase activity. Fast Garnet histochemistry of murine fibroblasts exposed to (A) conditioned medium collected from BHK<sup>HexlacZ</sup> cells compared to cells exposed to medium from normal parent BHK-21 cells (B). These results demonstrate that hHEXA & hHEXB, products of the  $\beta$ -Hex transgene, are released into the extracellular medium and can be absorbed by other cells via paracrine pathways resulting in induction of the cellular  $\beta$ -hexosaminidase.

[11] Figure 3 shows a representation of a lentiviral system containing the HexA and HexB genes. The 3-vector FIV(Hex) system The FIV(Hex) lentiviral system is comprised of 3 vectors: Packaging vector providing the packaging instructions in trans, - VSV-G envelop vector providing the envelop instructions in trans, - FIV(Hex) vector containing the therapeutic bicistronic gene.

[12] Figure 4 shows a representation of a Fiv(Hex) vector. Backbone FIV vector constructed by Proeschla et al. (1998)

[13] Figure 5 shows restriction fragment pattern of Feline immunodeficiency viral vector comprising a  $\beta$ -Hex construct. A maxi prep of FIV(Hex) clone 6.2 in 500 TB with 3X solution run through 2 columns. Yield of DNA was 1.095 mg. Final concentration is 1microg/microl. Restriction enzyme digest with ScaI, notI, SalI, and XhoI. The bands are as expected.

[14] Figure 6 shows fibroblast infection by FIV(Hex) in vitro.

[15] Figure 7 shows an FIV(Hex) titration experiment.

[16] Figure 8 shows FIV(lacZ) administration to adult mice. FIV(lacZ) infection of murine fibroblasts (CrFK's) *in vitro*, as well as of liver cells following direct transdermal intra-hepatic injection. Liver, brain and spleen sections stained for  $\beta$ -galactosidase following intraperitoneal injection to 3 month old mice. lacZ expression was detected by X-gal staining (blue stain) and immunocytochemistry (ICC; black stain) on fixed tissue sections harvested 1 month post-treatment.

[17] Figure 9 shows FIV(lacZ) administration to P4 mice. Liver, brain, spleen and kidney sections stained for  $\beta$ -galactosidase following intraperitoneal injection to mice of perinatal age (4 days old). lacZ expression was detected by X-gal staining (blue stain) on fixed tissue sections harvested 3 months post-treatment.

[18] Figure 10 shows dose response of IP injections. Young adult mice (6 weeks old) were injected intra-peritoneally with different doses of FIV(lacZ) {0.1 mL, 0.5 mL, 1.0 mL and 2.0mL of  $10^3$  infectious particles per mL} viral solution. One month following treatment the animals were sacrificed and lacZ reporter gene expression was measured. It was found that increasing doses of FIV result in increasing levels of gene therapy efficacy. In the clinical, human disease arena, this would optimally translate into intravenous administration of  $10^5$ - $10^6$  infectious FIV particles to ensure similar efficacy levels of gene therapy.

[19] Figure 11 shows diagrams of the vectors used to make the constructs discussed in Examples 1 and 2. FIV(Hex) is constructed by ligating the backbone part of FIV(LacZ), and the fragment of HexB-IRES-HexA from pHexLacZ. FIV(LacZ) is 12750 bp, after cut with SstII and NotI (generate 4500 bp and 8250 bp bands). Purify the 8250 bp band which contains the FIV backbone with CMV promoter. pHexlacZ is a construct of 10150 bp. Cut with NheI and NotI, there are 4700 bp and 5450 bp fragments. The 4700 bp band contains the structure of HexB-IRES-HexA, which doesn't have CMV.

[20] Figure 12 shows how the structure of FIV(Hex) was confirmed. The constructs were digested with different restriction enzymes: (Result see Figure 5). ScaI: cut once in the FIV backbone (generated one band 13 Kb). NotI: the site of ligation, and it is the only site (generated one band 13 Kb). Sal I: one site in HexB-IRFS-HexA and 3 sites in FIV backbone (generated one band t 8.5 Kb, one wide band with 2184 bp and 2400 bp, one band 34 bp which is invisible). Xho I: there is one site in HexB-IRFS-HexA and six

sites in the FIV backbone (FIV(LacZ) : at 502, 1410, 1453, 7559, 7883 and 9949 bp). These generated 6 bands (908 bp, 43 bp(invisible), 1.7Kb, 324 bp, 2066 bp, 3.3 Kb, and 2.8 Kb).

[21] Figure 13 shows a transcription termination cassette (STOP) flanked by 2 loxP sites was inserted between the promoter CMV and the therapeutic gene HexB-IRES-HexA. This results in inhibition of gene expression, until the STOP cassette is excisionally removed via the action of cre recombinase. The termination stop can consist of for example, a neomycin gene, whose termination signal acts as a termination signal for the rest of the transcript. Any reporter gene could be inserted and used in this way.

[22] Figure 14 shows a dually regulated inducible cre-recombinase system which was constructed. The activity of this construct is regulated exogenously by RU486. Furthermore, a stable cell line for this system was developed, whereby addition of RU486 in the culture media results in activation of cre-recombinase and subsequently excisional recombination of DNA, such as a transcription termination cassette flanked by 2 loxP sites.

[23] Figure 15 shows an example of the function of stable cell line, named GLVP/CrePr cell line, described in figure 14. In this case, the dual reporter vector CMV-lox-Luc-lox-AP was transiently transfected into the cell line. Alkaline phosphatase (AP) activity was evaluated in vitro after the addition of RU486 to the culture media by an AP histochemical staining method.

[24] Figure 16A shows the excisionally activated  $\beta$ -hexosaminidase gene Hex<sup>XAT</sup> was constructed by placing a floxed transcription termination cassette (STOP) upstream to the first open reading frame: CMV-loxP-STOP-loxP-HexB-IRES-HexA. Figure 16B shows Hex<sup>XAT</sup> was transiently transfected into our inducible cre cell line. Activation of cre-recombinase resulted in loxP directed DNA recombination and excision of the STOP cassette. Figure 16C Cre-mediated activation of Hex<sup>XAT</sup> resulted in HexA and HexB upregulation (column 1). RU486 stimulation of GLVP/CrePr results in site-directed recombination and subsequent activation of a dormant transcriptional unit. A. shows the p Hex<sup>XAT</sup>, a bicistronic transgene comprised of a "floxed" transcription-termination cassette (STOP), and both isoforms of the human  $\beta$ -hexosaminidase, was transiently transacted into the GLVP/CrePr cell line. B. RU-486 administration resulted in loxP-directed excisional recombination, C. resulting in transcriptional activation and synthesis of HexA and HexB mRNA.

[25] Figure 17 shows the semi-quantitative analysis for HexA and HexB showed induction of gene transcription following Hex<sup>XAT</sup> activation at (A) the mRNA level, (B) enzyme activity level in vitro, as well as (C) histochemical level in situ. RU486 significantly induces  $\beta$ -hexosaminidase expression in the GLVP/CrePr cell line.  $\beta$ -hexosaminidase activity was found significantly upregulated in p Hex<sup>XAT</sup>-transfected GLVP/CrePr cells 4 days after RU486 administration at the (A) HexA & HexB mRNA, (B) enzyme activity in vitro, as well as (C) in fixed monolayers in situ, as assessed by RTPCR, 4-MUG fluorescence and X-Hex histochemistry, respectively.

[26] Figure 18 shows Hex<sup>XAT</sup> was stably expressed in fibroblasts derived from a patient with Tay-Sachs disease (TSD). Gene activation was mediated by infection of the cells with a HSV aplicon viral vector capable of transducing cells with the cre recombinase. This figure demonstrates that activation of the Hex gene results in protection of the TSD cells from death following GM<sub>2</sub> substrate challenge.

[27] Figure 19 shows that the virus produced in Figure 3 above can resolve GM<sub>2</sub> storage in TSD cells cultured in vitro.

[28] Figure 20 shows the Hex gene was cloned in the FIV backbone as shown in Fig.3 producing the virus FIV(Hex), which was then used to infect TSD cells challenged with GM<sub>2</sub> substrate. This figures shows that delivery of our Hex gene with FIV(Hex) in TSD cells in vitro confers protection to cell death following GM<sub>2</sub> administration.

[29] Figure 21 shows HexB<sup>-/-</sup> knock out pups (2 days) were injected 100uL of FIV(Hex) virus intraperitoneally. The animals were monitored weekly while they assumed growth until sacrificed (16-18 weeks of age).

[30] Figure 22 shows expression of HEXB protein in adult mice that were injected with the FIV(Hex) virus as infants 2 days after birth. HEXB protein expression was detected by immunocytochemistry in the liver and brain of these mice.

[31] Figure 23 shows locomotive performance in relation to age (in weeks) of 6 mice that were treated 2 days after birth: 3 mice were injected with FIV(Hex) and 3 with FIV(lacZ) and served as controls. At 16 weeks of age, the "classic" stage that the hexB knockout mice display the disease, there was significant disease difference between the two groups.

## V. DETAILED DESCRIPTION

[32] The present invention may be understood more readily by reference to the following detailed description of preferred embodiments of the invention and the Examples included therein and to the Figures and their previous and following description.

5 [33] Before the present compounds, compositions, articles, devices, and/or methods are disclosed and described, it is to be understood that this invention is not limited to specific synthetic methods, specific recombinant biotechnology methods unless otherwise specified, or to particular reagents unless otherwise specified, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose  
10 of describing particular embodiments only and is not intended to be limiting.

[34] Disclosed are the components to be used to prepare the disclosed compositions as well as the compositions themselves to be used within the methods disclosed herein. These and other materials are disclosed herein, and it is understood that when combinations, subsets, interactions, groups, etc. of these materials are disclosed that  
15 while specific reference of each various individual and collective combinations and permutation of these compounds may not be explicitly disclosed, each is specifically contemplated and described herein. For example, if a particular  $\beta$ -Hex vector is disclosed and discussed and a number of modifications that can be made to a number of molecules including the  $\beta$ -Hex vector are discussed, specifically contemplated is each and every  
20 combination and permutation of the  $\beta$ -Hex vector and the modifications that are possible unless specifically indicated to the contrary. Thus, if a class of molecules A, B, and C are disclosed as well as a class of molecules D, E, and F and an example of a combination molecule, A-D is disclosed, then even if each is not individually recited each is individually and collectively contemplated meaning combinations, A-E, A-F, B-D, B-E, B-F, C-D, C-E,  
25 and C-F are considered disclosed. Likewise, any subset or combination of these is also disclosed. Thus, for example, the sub-group of A-E, B-F, and C-E would be considered disclosed. This concept applies to all aspects of this application including, but not limited to, steps in methods of making and using the disclosed compositions. Thus, if there are a variety of additional steps that can be performed it is understood that each of these  
30 additional steps can be performed with any specific embodiment or combination of embodiments of the disclosed methods.



#### A. Definitions

[35] As used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a pharmaceutical carrier" includes mixtures of two or more such carriers, and the like.

[36] Ranges may be expressed herein as from "about" one particular value, and/or to "about" another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent "about," it will be understood that the particular value forms another embodiment. It will be further understood that the endpoints of each of the ranges are significant both in relation to the other endpoint, and independently of the other endpoint. It also understood that for every value disclosed, "about" that value is also disclosed. For example, if the value "10" is disclosed, then "about 10" is also disclosed, even if not specifically recited out as "about 10."

[37] In this specification and in the claims which follow, reference will be made to a number of terms which shall be defined to have the following meanings:

[38] "Optional" or "optionally" means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances where it does not.

[39] "Primers" are a subset of probes which are capable of supporting some type of enzymatic manipulation and which can hybridize with a target nucleic acid such that the enzymatic manipulation can occur. A primer can be made from any combination of nucleotides or nucleotide derivatives or analogs available in the art which do not interfere with the enzymatic manipulation.

[40] "Probes" are molecules capable of interacting with a target nucleic acid, typically in a sequence specific manner, for example through hybridization. The hybridization of nucleic acids is well understood in the art and discussed herein. Typically a probe can be made from any combination of nucleotides or nucleotide derivatives or analogs available in the art.

[41] Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into

this application in order to more fully describe the state of the art to which this invention pertains. The references disclosed are also individually and specifically incorporated by reference herein for the material contained in them that is discussed in the sentence in which the reference is relied upon.

## 5 B. Compositions and methods

### 1. Lysosomal disorders

[42] Lysosomal storage disorders are a group of closely related metabolic diseases resulting from deficiency in enzymes essential for the degradation of gangliosides, mucopolysaccharides, as well as other complex macromolecules. With the dysfunction of a lysosomal enzyme, catabolism of correlate substrates remains incomplete, leading to accumulation of insoluble complex macromolecules within the lysosomes. For example,  $\beta$ -hexosaminidase defects result in lysosomal storage of GM<sub>2</sub> gangliosides leading to the development of Tay-Sachs or Sandhoff's disease. Similarly, mucopolysaccharidoses (MPS) are a group of closely related metabolic disorders that result from deficiencies in lysosomal enzymes involved in glycosaminoglycan metabolism, leading to lysosomal mucopolysaccharide storage. Affected patients, depending on the specific disorder and clinical severity, may present with neurodegeneration, mental retardation, paralysis, dementia and blindness, dysostosis multiplex, craniofacial malformations and facial dysfiguration. Below, some of the most common conditions of this family of diseases are summarized.

Representative examples of common lysosomal storage disorders

Disease	Enzyme Deficiency	Storage Metabolite
Glycogenosis-Type 2	$\alpha$ -1,4-Glucosidase	Glycogen
Gangliosidoses		
GM <sub>1</sub> Gangliosidosis	GM <sub>1</sub> ganglioside $\beta$ -galactosidase	GM <sub>1</sub> ganglioside
Tay-Sachs disease	Hexosaminidase – $\alpha$ subunit	GM <sub>2</sub> ganglioside
Sandhoff disease	Hexosaminidase – $\beta$ subunit	GM <sub>2</sub> ganglioside
Sulfatidoses		
Krabbe disease	Galactosylceramidase	galactocerebroside
Fabry disease	$\alpha$ -Galactosidase A	ceramide trihexoside
Gaucher disease	Glucocerebrosidase	glucocerebroside
Niemann-Pick – types A & B	Sphingomyelinase	sphingomyelin
Mucopolysaccharidoses		
Hurler's syndrome	$\alpha$ -L-Iduronidase	dermatan/heparan sulfate
Hunter's syndrome	L-Iduronosulfate sulfatase	
Mucolipidoses		
Mucopolipidosis – II	Mannose-6-phosphate kinases	mucopolysaccharide/ glycolipid
Pseudo-Hurler's		

Fucosidosis	$\alpha$ -Fucosidase	Glycoproteins
Mannosidosis	$\alpha$ -Mannosidase	oligosaccharides
Wolman Disease	Acid Lipase	triglycerides

## 2. Histopathology & Pathophysiology - A progressive disorder

[43] In storage diseases, the affected cells become distended and display vacuolated cytoplasm, which appear as swollen lysosomes under the electronic microscope. For example, in the central nervous system, the neurons of the brain, trigeminal and spinal root ganglia in patients suffering from GM<sub>2</sub> gangliosidosis display swollen vacuolated perikarya stored with excessive amounts of lysosomal storage. As a result, these organelles become large in size and numbers, interfering with normal cell functions. The formation of meganeurites, axon hillock enlargements accompanied by secondary neuritic sprouting, present as cardinal histopathological feature of gangliosidosis and mucopolysaccharidoses (Purpura and Suzuki, 1976; Walkley *et al.*, 1988). Purpura and Suzuki proposed that meganeurites, and the synapses they develop, contribute to the onset and progression of neuronal dysfunction in storage diseases, by altering electrical properties of neurons and modifying integrative operations of somatodendritic synaptic inputs. In addition, Walkley *et al.* (1991) suggested that this neuroaxonal dystrophy commonly involved GABAergic neurons, and proposed that the resulting defect in neurotransmission in inhibitory circuits may be an important factor underlying brain dysfunction in lysosomal storage diseases. Consequently, the clinical phenotype often includes neurodegeneration, mental retardation, paralysis, dementia and blindness. In addition, some storage disorders also affect peripheral tissues, such as cartilage and bone, resulting in abnormal growth & development of long bones, vertebrae, ribs and jaws, ultimately leading to anomalies of the skeleton, the cranium and dysfiguration of the face (Mucopolysaccharidoses, and Sandhoff's disease to some degree).

[44] One cardinal characteristic of storage disorders is their progressively worsening (progressive) nature. The deficiency of metabolic enzymes results in accumulation of insoluble metabolites in the lysosomes, which becomes excessive and deleterious over time due to the additive effects of accumulating insoluble metabolite storage. For example, patients suffering from mucopolysaccharidoses (Hurler's or Hunter's) display only a mild degree of the disease's phenotype at infancy, but, due to increasing storage over time, progress to severe forms by adolescence, often leading to death (Gorlin *et al.*, 1990). This provides a window of opportunity in mammalian development during

which the pathophysiological process of the disease can be attenuated by restoring lysosomal enzymatic activity early enough in life to prevent the development of a "full-blown" disease and, perhaps, to reverse its progression.

### 3. Tay-Sachs & Sandhoffs disorders

5           [45]   The lysosomal enzyme  $\beta$ -hexosaminidase is comprised of 2 subunits (peptides), HEX- $\alpha$  and HEX- $\beta$ , encoded by two distinct genes, HexA and HexB, respectively.  $\beta$ -hexosaminidase exists in 3 isoforms (proteins), HEXA ( $\alpha/\beta$  heterodimer), HEXB ( $\beta/\beta$  homodimer) and HEXS ( $\alpha/\alpha$  homodimer). Mutation of the HexA gene, causing functional problems with the HEX- $\alpha$  polypeptide in humans results in Tay Sachs disease, 10   whereas mutation of the HexB gene, causing functional problems in the  $\beta$ -Hex polypeptide, in Sandhoff's disease. In Tay Sachs disease, HexA mutation results in loss of HEXA isoform ( $\alpha/\beta$  heterodimer), whereas in Sandhoff's disease, HexB mutation results in loss of both HEXA ( $\alpha/\beta$  heterodimer) and HexB ( $\beta/\beta$  homodimer) isoforms, leading to a more severe clinical phenotype. Affected patients, depending on the clinical severity, may 15   present with neurodegeneration, mental and motor deterioration, dysarthria, impaired thermal sensitivity, blindness, as well as facial dysfiguration (doll-like and coarse facies). Histopathologically, the cells of the brain (neurons and glia), spleen and cartilage appear swollen with vacuolated/clear perikarya suggestive of lysosomal storage. Biochemical analysis reveals a complete lack of  $\beta$ -hexosaminidase activity accompanied by lysosomal 20   accumulation of GM<sub>2</sub> gangliosides. As a result, the lysosomes become large in size and numbers, significantly crippling normal cellular function. Clinically, it is not uncommon for patients to display only mild features at infancy, but due to increasing storage over time, progress to severe forms of the disease by adolescence (Gorlin *et al.*, 1990). Similarly, other affected mammals, such as affected mice pups, display only mild anomalies at birth, but 25   quickly develop their distinct abnormal features (1 month of age).

### 4. Blood brain barrier formation

          [46]   The blood-brain barrier (BBB) is a structure unique to the central nervous system and is the result of tight junctions between the brain endothelial cells (Goldstein *et al.*, 1986). Previous work (Risau *et al.*, 1986) on the development of mouse BBB using large 30   protein molecules (horse radish peroxidase) suggested BBB formation during the late days of embryonic life (E17 in mouse). Furthermore, BBB in the adult is not absolute; whereby certain areas of the brain do not develop BBB and thus allow for free exchange of

molecules through them. These areas include the median eminence (hypothalamus), pituitary, choroids plexus, pineal gland, subfornical organ, organum vasculosum lamina terminalis and area postrema (Risau & Wolburg, 1990). This allows for the intrusion of FIV(Hex) virions into the brain matter through an incomplete BBB as well as through areas lacking BBB during the first few days after birth as discussed in the examples herein. Disclosed herein a diffuse expression of lacZ throughout the brain of P4 mice injected with FIV(lacZ) versus periventricular only localization following "adult administration" was shown.

#### 5. Immune system development

[47] Specific immunity in vertebrates is dependent on the host's ability to generate a heterogeneous repertoire of antigen-binding structures that are displayed on the surface of lymphocytes. Immunologic competence arises early in mammalian development. Since the expression of  $\beta$ -Hex therapeutic gene in *hexA<sup>-/-</sup>/hexB<sup>-/-</sup>* mice may be perceived as presentation of "non-self" antigens, one needs to consider the possibility of an immune response against human HEXA and HEXB following gene therapy. In these terms, perinatal administration can offer a unique opportunity in gene therapy application. Specifically, numerous studies have documented that the human and mouse neonate is unable to mount satisfactory responses to various antigenic challenges, which in many instances is delayed well beyond infancy (Schroeder et al., 1995). Therefore, due to this "immature" immunologic state of mice and humans early in their postnatal life, perinatal gene therapy is consistent with adequate "training" of the immune system to recognize HEXA and HEXB as "self" antigens circumventing any potential immunologic rejection. It is understood that the transtherapy can take place in an infant as well.

[48] Disclosed are nucleic acids comprising sequence encoding HEX- $\alpha$  and sequence encoding HEX- $\beta$ . Also disclosed are nucleic acids, wherein the nucleic acid further comprises an IRES sequence, wherein the nucleic acids express more than one IRES sequence, wherein the vectors express an IRES sequence after each Hex nucleic acid, wherein the nucleic acid further comprises a promoter sequence, wherein the nucleic acid further comprises a promoter sequence, wherein the HEX- $\beta$  has at least 80% identity to the sequence set forth in SEQ ID NO:3 and the HEX- $\alpha$  has at least 80% identity to the sequence set forth in SEQ ID NO:1, wherein the HEX- $\beta$  has at least 85% identity to the sequence set forth in SEQ ID NO:3 and the HEX- $\alpha$  has at least 80% identity to the

sequence set forth in SEQ ID NO:1, wherein the HEX- $\beta$  has at least 90% identity to the sequence set forth in SEQ ID NO:3 and the HEX- $\alpha$  has at least 80% identity to the sequence set forth in SEQ ID NO:1, wherein the HEX- $\beta$  has at least 95% identity to the sequence set forth in SEQ ID NO:3 and the HEX- $\alpha$  has at least 80% identity to the sequence set forth in SEQ ID NO:1, wherein the HEX- $\beta$  has the sequence set forth in SEQ ID NO:3 and the HEX- $\alpha$  has the sequence set forth in SEQ ID NO:1, wherein the sequence encoding the HEX- $\beta$  is orientated 5' to the sequence encoding HEX- $\alpha$ , wherein the sequence encoding the HEX- $\beta$  is orientated 5' to the IRES sequence and the IRES sequence is located 5' to the sequence encoding HEX- $\alpha$ , wherein the promoter is located 5' to the sequence encoding the HEX- $\beta$  and the sequence encoding the HEX- $\beta$  is orientated 5' to the IRES sequence and the IRES sequence is located 5' to the sequence encoding HEX- $\alpha$ .

[49] Also disclosed are vectors comprising the disclosed nucleic acids. Also disclosed are cells comprising the disclosed nucleic acids and vectors.

[50] Also disclosed are non-human mammal comprising the disclosed nucleic acids, vectors, and cells disclosed herein.

[51] Also disclosed are methods of providing HEX- $\alpha$  in a cell comprising transfecting the cell with the nucleic acids, also disclosed are methods of providing HEX- $\beta$  in a cell comprising transfecting the cell with the nucleic acids, also disclosed are method of providing HEX- $\alpha$  and HEX- $\beta$  in a cell comprising transfecting the cell with the nucleic acid of claims 1-4.

[52] Also disclosed are method of delivering the disclosed compositions, wherein the transfection occurs in vitro or in vivo.

[53] Disclosed are methods of making a transgenic organism comprising administering the disclosed nucleic acids, vectors and/or cells.

[54] Disclosed are methods of making a transgenic organism comprising transfecting a lentiviral vector to the organism at during a perinatal stage of the organism's development.

[55] Also disclosed are methods of treating a subject having Tay Sachs disease and/or Sandoff disease comprising administering any of the disclosed compounds and compositions.

### C. Compositions

#### 1. $\beta$ -Hexosaminidase transgene ( $\beta$ -Hex)

[56] The  $\beta$ -Hexosaminidase protein is a protein comprised of two subunits, one subunit is encoded by the HexA gene and a second subunit encoded by the gene HexB. The human HexA Exon 1 can be found 316 bp upstream of MstII site; chromosome 15q11-15qter. The human HexA gene can be found at human chromosomal region 15q23----q24. The human HexB gene can be found on chromosome 5, map 5q13..

[57] Disclosed are constructs capable of expressing both the HexA gene product and the HexB gene product, from a single construct. Any construct capable of expressing both the HexA and HexB gene products is referred to as a  $\beta$ -Hex construct herein. The  $\beta$ -Hex construct allows for synthesis of all  $\beta$ -hexosaminidase protein isoforms, HEXA ( $\alpha/\beta$  heterodimer), HEXB ( $\beta/\beta$  homodimer) and HEXS ( $\alpha/\alpha$  homodimer). Disclosed are nucleic acid constructs comprising a cytomegalovirus (CMV) promoter-driven bicistronic gene ( $\beta$ -Hex) that encodes for both human HexA and HexB genes, which can lead to the synthesis of functional  $\beta$ -hexosaminidase isoenzymes.

[58] The  $\beta$ -Hex construct typically comprises four parts: 1) a promoter, 2) the HexA coding sequence, 3) the HexB coding sequence, and 4) an IRES sequence (integrated ribosomal entry site). These four parts can be integrated into any vector delivery system. In preferred embodiments, the orientation of the four parts is 5'-promoter-HexB-IRES-HexA-3'.

[59] The promoter can be any promoter, such as those discussed herein. It is understood as discussed herein that there are functional variants of the HexA and HexB which can be made. Furthermore, it is understood that there are functional variants of the IRES element, for example as discussed herein. Typically the genes to be expressed are placed on either side of the IRES sequence.

[60] The IRES element is an internal ribosomal entry sequence which can be isolated from the encephalomyocarditis virus (ECMV). This element allows multiple genes to be expressed and correctly translated when the genes are on the same construct. IRES sequences are discussed in for example, United States Patent No: 4,937,190 which is herein incorporated by reference at least for material related to IRES sequences and their use.

[61] HexA and HexB cDNA can be obtained from the American Tissue Culture Collection. (American Tissue Culture Collection, Manassas, VA 20110-2209; Hex- $\alpha$ : ATCC# 57206; Hex- $\beta$  ATCC# 57350) The IRES sequence can be obtained from a number of sources including commercial sources, such as the pIRES expressing vector from  
5 Clontech (Clontech, Palo Alto CA 94303-4230).

[62] Also disclosed are tricistronic constructs encoding for both isoforms of human  $\beta$ -hexosaminidase, hHexA & hHexB, as well as the  $\beta$ -galactosidase reporter gene (*lacZ*).

[63] Global delivery of the disclosed constructs is also disclosed. Disclosed is a  
10 pseudotyped feline immunodeficiency virus (FIV) for global  $\beta$ -Hex delivery. Stable expression of the therapeutic gene aids prolonged restoration of the genetic anomaly enhancing treatment efficacy and contributing to long-term therapeutic outcomes. The backbone FIV system has been shown to effectively incorporate, due to its lentiviral properties, the transgene of interest into the host's genome, allowing for stable gene  
15 expression (Poeschla et al., 1998). Disclosed herein is stable expression of the reporter gene *lacZ* for over 3 months in mice following perinatal systemic FIV(*lacZ*) administration.

[64] A model system for the study of these vectors is a mouse that is knockout mouse deficient in both HexA and HexB, since the *hexA*<sup>-/-</sup>/*hexB*<sup>-/-</sup> mouse is characterized by global disruption of the *hexA* and *hexB* genes. Gene disruption in this mouse is global, and  
20 therefore, can be used as a model for global replacement. The timing of gene therapy is important as it is closely related to the temporal development of the disorder. *HexA*<sup>-/-</sup>/*hexB*<sup>-/-</sup> mice display mild phenotype aberrations at birth and quickly develop craniofacial dysplasia by 4-5 weeks of age. Similarly, it is not uncommon for patients suffering from this class of genetic disorders to display only mild degree of the disease at infancy, and to  
25 progress to severe forms by adolescence.

## 2. Delivery of the compositions to cells

[65] Delivery can be applied, in general, via local or systemic routes of administration. Local administration includes virus injection directly into the region or organ of interest, versus intravenous (*IV*) or intraperitoneal (*IP*) injections (systemic)  
30 aiming at viral delivery to multiple sites and organs via the blood circulation. Previous research on the effects of local administration demonstrated gene expression limited to the site/organ of the injection, which did not extend to the rest of the body (Daly et al., 1999a;



Kordower et al., 1999). Furthermore, previous studies have demonstrated successful global gene transfer to multiple tissues and organs in rodents and primates following viral *IV* and *IP* injections (Daly et al., 1999b; Tarntal et al., 2001; McCormack et al., 2001; Lipschutz et al., 2001). Disclosed herein *IP* injection of FIV(lacZ) in mice of adult (3 months old) as well as of perinatal age (P4) resulted in global transfer and expression of the reporter gene lacZ in brain, liver, spleen and kidney. Also disclosed, the levels of expression achieved via *IP* injections were superior to those acquired following local administration directly into the liver.

[66] There are a number of compositions and methods which can be used to deliver nucleic acids to cells, either in vitro or in vivo. These methods and compositions can largely be broken down into two classes: viral based delivery systems and non-viral based delivery systems. For example, the nucleic acids can be delivered through a number of direct delivery systems such as, electroporation, lipofection, calcium phosphate precipitation, plasmids, viral vectors, viral nucleic acids, phage nucleic acids, phages, cosmids, or via transfer of genetic material in cells or carriers such as cationic liposomes. Appropriate means for transfection, including viral vectors, chemical transfectants, or physico-mechanical methods such as electroporation and direct diffusion of DNA, are described by, for example, Wolff, J. A., et al., Science, 247, 1465-1468, (1990); and Wolff, J. A. Nature, 352, 815-818, (1991). Such methods are well known in the art and readily adaptable for use with the compositions and methods described herein. In certain cases, the methods will be modified to specifically function with large DNA molecules. Further, these methods can be used to target certain diseases and cell populations by using the targeting characteristics of the carrier.

a) Nucleic acid based delivery systems

[67] Transfer vectors can be any nucleotide construction used to deliver genes into cells (e.g., a plasmid), or as part of a general strategy to deliver genes, e.g., as part of recombinant retrovirus or adenovirus (Ram et al. Cancer Res. 53:83-88, (1993)).

[68] As used herein, plasmid or viral vectors are agents that transport the disclosed nucleic acids, such as the  $\beta$ -Hex construct into the cell without degradation and include a promoter yielding expression of the HexA and HexB encoding sequences in the cells into which it is delivered. In some embodiments the vectors for the  $\beta$ -Hex constructs are derived from either a virus, retrovirus, or lentivirus. Viral vectors can be, for example,

Adenovirus, Adeno-associated virus, Herpes virus, Vaccinia virus, Polio virus, AIDS virus, neuronal trophic virus, Sindbis and other RNA viruses, including these viruses with the HIV backbone, and lentiviruses. Also preferred are any viral families which share the properties of these viruses which make them suitable for use as vectors. Retroviruses include Murine  
5 Maloney Leukemia virus, MMLV, and retroviruses that express the desirable properties of MMLV as a vector. Retroviral vectors are able to carry a larger genetic payload, i.e., a transgene, such as, the disclosed  $\beta$ -Hex constructs or marker gene, than other viral vectors, and for this reason are a commonly used vector. However, they are not as useful in non-proliferating cells. Adenovirus vectors are relatively stable and easy to work with, have  
10 high titers, and can be delivered in aerosol formulation, and can transfect non-dividing cells. Pox viral vectors are large and have several sites for inserting genes, they are thermostable and can be stored at room temperature. A preferred embodiment is a viral vector, which has been engineered so as to suppress the immune response of the host organism, elicited by the viral antigens. Preferred vectors of this type will carry coding regions for Interleukin 8 or  
15 10.

[69] Viral vectors can have higher transaction (ability to introduce genes) abilities than chemical or physical methods to introduce genes into cells. Typically, viral vectors contain, nonstructural early genes, structural late genes, an RNA polymerase III transcript, inverted terminal repeats necessary for replication and encapsidation, and promoters to  
20 control the transcription and replication of the viral genome. When engineered as vectors, viruses typically have one or more of the early genes removed and a gene or gene/promotor cassette is inserted into the viral genome in place of the removed viral DNA. Constructs of this type can carry up to about 8 kb of foreign genetic material. The necessary functions of the removed early genes are typically supplied by cell lines which have been engineered to  
25 express the gene products of the early genes in trans.

#### (1) Retroviral Vectors

[70] A retrovirus is an animal virus belonging to the virus family of Retroviridae, including any types, subfamilies, genus, or tropisms. Retroviral vectors, in general, are described by Verma, I.M., Retroviral vectors for gene transfer. In Microbiology-1985,  
30 American Society for Microbiology, pp. 229-232, Washington, (1985), which is incorporated by reference herein. Examples of methods for using retroviral vectors for gene therapy are described in U.S. Patent Nos. 4,868,116 and 4,980,286; PCT applications WO

90/02806 and WO 89/07136; and Mulligan, (Science 260:926-932 (1993)); the teachings of which are incorporated herein by reference.

[71] A retrovirus is essentially a package which has packed into it nucleic acid cargo. The nucleic acid cargo carries with it a packaging signal, which ensures that the replicated daughter molecules will be efficiently packaged within the package coat. In addition to the package signal, there are a number of molecules which are needed in cis, for the replication, and packaging of the replicated virus. Typically a retroviral genome, contains the gag, pol, and env genes which are involved in the making of the protein coat. It is the gag, pol, and env genes which are typically replaced by the foreign DNA that it is to be transferred to the target cell. Retrovirus vectors typically contain a packaging signal for incorporation into the package coat, a sequence which signals the start of the gag transcription unit, elements necessary for reverse transcription, including a primer binding site to bind the tRNA primer of reverse transcription, terminal repeat sequences that guide the switch of RNA strands during DNA synthesis, a purine rich sequence 5' to the 3' LTR that serve as the priming site for the synthesis of the second strand of DNA synthesis, and specific sequences near the ends of the LTRs that enable the insertion of the DNA state of the retrovirus to insert into the host genome. The removal of the gag, pol, and env genes allows for about 8 kb of foreign sequence to be inserted into the viral genome, become reverse transcribed, and upon replication be packaged into a new retroviral particle. This amount of nucleic acid is sufficient for the delivery of a one to many genes depending on the size of each transcript. It is preferable to include either positive or negative selectable markers along with other genes in the insert.

[72] Since the replication machinery and packaging proteins in most retroviral vectors have been removed (gag, pol, and env), the vectors are typically generated by placing them into a packaging cell line. A packaging cell line is a cell line which has been transfected or transformed with a retrovirus that contains the replication and packaging machinery, but lacks any packaging signal. When the vector carrying the DNA of choice is transfected into these cell lines, the vector containing the gene of interest is replicated and packaged into new retroviral particles, by the machinery provided in cis by the helper cell. The genomes for the machinery are not packaged because they lack the necessary signals.

## (2) Adenoviral Vectors

- [73] The construction of replication-defective adenoviruses has been described (Berkner et al., *J. Virology* 61:1213-1220 (1987); Massie et al., *Mol. Cell. Biol.* 6:2872-2883 (1986); Haj-Ahmad et al., *J. Virology* 57:267-274 (1986); Davidson et al., *J. Virology* 61:1226-1239 (1987); Zhang "Generation and identification of recombinant adenovirus by liposome-mediated transfection and PCR analysis" *BioTechniques* 15:868-872 (1993)). The benefit of the use of these viruses as vectors is that they are limited in the extent to which they can spread to other cell types, since they can replicate within an initial infected cell, but are unable to form new infectious viral particles. Recombinant adenoviruses have been shown to achieve high efficiency gene transfer after direct, in vivo delivery to airway epithelium, hepatocytes, vascular endothelium, CNS parenchyma and a number of other tissue sites (Morsy, *J. Clin. Invest.* 92:1580-1586 (1993); Kirshenbaum, *J. Clin. Invest.* 92:381-387 (1993); Roessler, *J. Clin. Invest.* 92:1085-1092 (1993); Moullier, *Nature Genetics* 4:154-159 (1993); La Salle, *Science* 259:988-990 (1993); Gomez-Foix, *J. Biol. Chem.* 267:25129-25134 (1992); Rich, *Human Gene Therapy* 4:461-476 (1993); Zabner, *Nature Genetics* 6:75-83 (1994); Guzman, *Circulation Research* 73:1201-1207 (1993); Bout, *Human Gene Therapy* 5:3-10 (1994); Zabner, *Cell* 75:207-216 (1993); Caillaud, *Eur. J. Neuroscience* 5:1287-1291 (1993); and Ragot, *J. Gen. Virology* 74:501-507 (1993)). Recombinant adenoviruses achieve gene transduction by binding to specific cell surface receptors, after which the virus is internalized by receptor-mediated endocytosis, in the same manner as wild type or replication-defective adenovirus (Chardonnet and Dales, *Virology* 40:462-477 (1970); Brown and Burlingham, *J. Virology* 12:386-396 (1973); Svensson and Persson, *J. Virology* 55:442-449 (1985); Seth, et al., *J. Virol.* 51:650-655 (1984); Seth, et al., *Mol. Cell. Biol.* 4:1528-1533 (1984); Varga et al., *J. Virology* 65:6061-6070 (1991); Wickham et al., *Cell* 73:309-319 (1993)).

[74] A viral vector can be one based on an adenovirus which has had the E1 gene removed and these virions are generated in a cell line such as the human 293 cell line. In another preferred embodiment both the E1 and E3 genes are removed from the adenovirus genome.

30

## (3) Adeno-associated viral vectors

[75] Another type of viral vector is based on an adeno-associated virus (AAV). This defective parvovirus is a preferred vector because it can infect many cell types and is nonpathogenic to humans. AAV type vectors can transport about 4 to 5 kb and wild type

AAV is known to stably insert into chromosome 19. Vectors which contain this site specific integration property are preferred. An especially preferred embodiment of this type of vector is the P4.1 C vector produced by Avigen, San Francisco, CA, which can contain the herpes simplex virus thymidine kinase gene, HSV-tk, and/or a marker gene, such as the  
5 gene encoding the green fluorescent protein, GFP.

[76] In another type of AAV virus, the AAV contains a pair of inverted terminal repeats (ITRs) which flank at least one cassette containing a promoter which directs cell-specific expression operably linked to a heterologous gene. Heterologous in this context refers to any nucleotide sequence or gene which is not native to the AAV or B19  
10 parvovirus.

[77] Typically the AAV and B19 coding regions have been deleted, resulting in a safe, noncytotoxic vector. The AAV ITRs, or modifications thereof, confer infectivity and site-specific integration, but not cytotoxicity, and the promoter directs cell-specific expression. United states Patent No. 6,261,834 is herein incorporated by reference for  
15 material related to the AAV vector.

[78] The vectors of the present invention thus provide DNA molecules which are capable of integration into a mammalian chromosome without substantial toxicity.

[79] The inserted genes in viral and retroviral usually contain promoters, and/or enhancers to help control the expression of the desired gene product. A promoter is  
20 generally a sequence or sequences of DNA that function when in a relatively fixed location in regard to the transcription start site. A promoter contains core elements required for basic interaction of RNA polymerase and transcription factors, and may contain upstream elements and response elements.

#### (4) Lentiviral vectors

[01] The vectors can be lentiviral vectors, including but not limited to, SIV  
25 vectors, HIV vectors or a hybrid construct of these vectors, including viruses with the HIV backbone. These vectors also include first, second and third generation lentiviruses. Third generation lentiviruses have lentiviral packaging genes split into at least 3 independent plasmids or constructs. Also vectors can be any viral family that share the properties of  
30 these viruses which make them suitable for use as vectors. Lentiviral vectors are a special type of retroviral vector which are typically characterized by having a long incubation period for infection. Furthermore, lentiviral vectors can infect non-dividing cells.

Lentiviral vectors are based on the nucleic acid backbone of a virus from the lentiviral family of viruses. Typically, a lentiviral vector contains the 5' and 3' LTR regions of a lentivirus, such as SIV and HIV. Lentiviral vectors also typically contain the Rev Responsive Element (RRE) of a lentivirus, such as SIV and HIV.

5

(a) *Feline immunodeficiency viral vectors*

[80] One type of vector that the disclosed constructs can be delivered in is the VSV-G pseudotyped Feline Immunodeficiency Virus system developed by Poeschla *et al.* (1998). This lentivirus has been shown to efficiently infect dividing, growth arrested as well as post-mitotic cells. Furthermore, due to its lentiviral properties, it allows for incorporation of the transgene into the host's genome, leading to stable gene expression. This is a 3-vector system, whereby each confers distinct instructions: the FIV vector carries the transgene of interest and lentiviral apparatus with mutated packaging and envelope genes. A vesicular stomatitis virus G-glycoprotein vector (VSV-G; Burns *et al.*, 1993) contributes to the formation of the viral envelope *in trans*. The third vector confers packaging instructions *in trans* (Poeschla *et al.*, 1998). FIV production is accomplished *in vitro* following co-transfection of the aforementioned vectors into 293-T cells. The FIV-rich supernatant is then collected, filtered and can be used directly or following concentration by centrifugation. Titers routinely range between  $10^4 - 10^7$  bfu/ml..

10

15

(5) Packaging vectors

20

25

30

[81] As discussed above, retroviral vectors are based on retroviruses which contain a number of different sequence elements that control things as diverse as integration of the virus, replication of the integrated virus, replication of un-integrated virus, cellular invasion, and packaging of the virus into infectious particles. While the vectors in theory could contain all of their necessary elements, as well as an exogenous gene element (if the exogenous gene element is small enough) typically many of the necessary elements are removed. Since all of the packaging and replication components have been removed from the typical retroviral, including lentiviral, vectors which will be used within a subject, the vectors need to be packaged into the initial infectious particle through the use of packaging vectors and packaging cell lines. Typically retroviral vectors have been engineered so that the myriad functions of the retrovirus are separated onto at least two vectors, a packaging vector and a delivery vector. This type of system then requires the presence of all of the vectors providing all of the elements in the same cell before an infectious particle can be produced. The packaging vector typically carries the structural and replication genes

derived from the retrovirus, and the delivery vector is the vector that carries the exogenous gene element that is preferably expressed in the target cell. These types of systems can split the packaging functions of the packaging vector into multiple vectors, e.g., third-generation lentivirus systems. Dull, T. et al., "A Third-generation lentivirus vector with a conditional packaging system" J. Virol 72(11):8463-71 (1998)

[82] Retroviruses typically contain an envelope protein (env). The Env protein is in essence the protein which surrounds the nucleic acid cargo. Furthermore cellular infection specificity is based on the particular Env protein associated with a typical retrovirus. In typical packaging vector/delivery vector systems, the Env protein is expressed from a separate vector than for example the protease (pro) or integrase (in) proteins.

#### (6) Packaging cell lines

[83] The vectors are typically generated by placing them into a packaging cell line. A packaging cell line is a cell line which has been transfected or transformed with a retrovirus that contains the replication and packaging machinery, but lacks any packaging signal. When the vector carrying the DNA of choice is transfected into these cell lines, the vector containing the gene of interest is replicated and packaged into new retroviral particles, by the machinery provided in cis by the helper cell. The genomes for the machinery are not packaged because they lack the necessary signals. One type of packaging cell line is a 293 cell line.

#### (7) Large payload viral vectors

[84] Molecular genetic experiments with large human herpesviruses have provided a means whereby large heterologous DNA fragments can be cloned, propagated and established in cells permissive for infection with herpesviruses (Sun et al., Nature genetics 8: 33-41, 1994; Cotter and Robertson, Curr Opin Mol Ther 5: 633-644, 1999). These large DNA viruses (herpes simplex virus (HSV) and Epstein-Barr virus (EBV), have the potential to deliver fragments of human heterologous DNA > 150 kb to specific cells. EBV recombinants can maintain large pieces of DNA in the infected B-cells as episomal DNA. Individual clones carried human genomic inserts up to 330 kb appeared genetically stable. The maintenance of these episomes requires a specific EBV nuclear protein, EBNA1, constitutively expressed during infection with EBV. Additionally, these vectors can be used for transfection, where large amounts of protein can be generated transiently in vitro.

Herpesvirus amplicon systems are also being used to package pieces of DNA > 220 kb and to infect cells that can stably maintain DNA as episomes.

[85] Other useful systems include, for example, replicating and host-restricted non-replicating vaccinia virus vectors.

5                                   b) Non-nucleic acid based systems

[86] The disclosed compositions can be delivered to the target cells in a variety of ways. For example, the compositions can be delivered through electroporation, or through lipofection, or through calcium phosphate precipitation. The delivery mechanism chosen will depend in part on the type of cell targeted and whether the delivery is occurring for  
10   example in vivo or in vitro.

[87] Thus, the compositions can comprise, in addition to the disclosed constructs or vectors for example, lipids such as liposomes, such as cationic liposomes (e.g., DOTMA, DOPE, DC-cholesterol) or anionic liposomes. Liposomes can further comprise proteins to facilitate targeting a particular cell, if desired. Administration of a composition comprising  
15   a compound and a cationic liposome can be administered to the blood afferent to a target organ or inhaled into the respiratory tract to target cells of the respiratory tract. Regarding liposomes, see, e.g., Brigham et al. *Am. J. Resp. Cell. Mol. Biol.* 1:95-100 (1989); Felgner et al. *Proc. Natl. Acad. Sci USA* 84:7413-7417 (1987); U.S. Pat. No.4,897,355. Furthermore, the compound can be administered as a component of a microcapsule that can be targeted to  
20   specific cell types, such as macrophages, or where the diffusion of the compound or delivery of the compound from the microcapsule is designed for a specific rate or dosage.

[88] In the methods described above which include the administration and uptake of exogenous DNA into the cells of a subject (i.e., gene transduction or transfection), delivery of the compositions to cells can be via a variety of mechanisms. As one example,  
25   delivery can be via a liposome, using commercially available liposome preparations such as LIPOFECTIN, LIPOFECTAMINE (GIBCO-BRL, Inc., Gaithersburg, MD), SUPERFECT (Qiagen, Inc. Hilden, Germany) and TRANSFECTAM (Promega Biotec, Inc., Madison, WI), as well as other liposomes developed according to procedures standard in the art. In addition, the nucleic acid or vector of this invention can be delivered *in vivo* by  
30   electroporation, the technology for which is available from Genetronics, Inc. (San Diego, CA) as well as by means of a SONOPORATION machine (ImaRx Pharmaceutical Corp., Tucson, AZ).



[89] The materials may be in solution, suspension (for example, incorporated into microparticles, liposomes, or cells). These may be targeted to a particular cell type via antibodies, receptors, or receptor ligands. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Senter, et al., Bioconjugate Chem., 2:447-451, (1991); Bagshawe, K.D., Br. J. Cancer, 60:275-281, (1989); Bagshawe, et al., Br. J. Cancer, 58:700-703, (1988); Senter, et al., Bioconjugate Chem., 4:3-9, (1993); Battelli, et al., Cancer Immunol. Immunother., 35:421-425, (1992); Pietersz and McKenzie, Immunolog. Reviews, 129:57-80, (1992); and Roffler, et al., Biochem. Pharmacol., 42:2062-2065, (1991)). These techniques can be used for a variety of other specific cell types. Vehicles such as "stealth" and other antibody conjugated liposomes (including lipid mediated drug targeting to colonic carcinoma), receptor mediated targeting of DNA through cell specific ligands, lymphocyte directed tumor targeting, and highly specific therapeutic retroviral targeting of murine glioma cells *in vivo*. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Hughes et al., Cancer Research, 49:6214-6220, (1989); and Litzinger and Huang, Biochimica et Biophysica Acta, 1104:179-187, (1992)). In general, receptors are involved in pathways of endocytosis, either constitutive or ligand induced. These receptors cluster in clathrin-coated pits, enter the cell via clathrin-coated vesicles, pass through an acidified endosome in which the receptors are sorted, and then either recycle to the cell surface, become stored intracellularly, or are degraded in lysosomes. The internalization pathways serve a variety of functions, such as nutrient uptake, removal of activated proteins, clearance of macromolecules, opportunistic entry of viruses and toxins, dissociation and degradation of ligand, and receptor-level regulation. Many receptors follow more than one intracellular pathway, depending on the cell type, receptor concentration, type of ligand, ligand valency, and ligand concentration. Molecular and cellular mechanisms of receptor-mediated endocytosis has been reviewed (Brown and Greene, DNA and Cell Biology 10:6, 399-409 (1991)).

[90] Nucleic acids that are delivered to cells which are to be integrated into the host cell genome, typically contain integration sequences. These sequences are often viral related sequences, particularly when viral based systems are used. These viral integration systems can also be incorporated into nucleic acids which are to be delivered using a non-nucleic acid based system of delivery, such as a liposome, so that the nucleic acid contained in the delivery system can be come integrated into the host genome.

[91] Other general techniques for integration into the host genome include, for example, systems designed to promote homologous recombination with the host genome. These systems typically rely on sequence flanking the nucleic acid to be expressed that has enough homology with a target sequence within the host cell genome that recombination  
5 between the vector nucleic acid and the target nucleic acid takes place, causing the delivered nucleic acid to be integrated into the host genome. These systems and the methods necessary to promote homologous recombination are known to those of skill in the art.

c) *In vivo/ex vivo*

10 [92] As described herein, the compositions can be administered in a pharmaceutically acceptable carrier and can be delivered to the subjects cells *in vivo* and/or *ex vivo* by a variety of mechanisms well known in the art (e.g., uptake of naked DNA, liposome fusion, intramuscular injection of DNA via a gene gun, endocytosis and the like).

[93] If *ex vivo* methods are employed, cells or tissues can be removed and  
15 maintained outside the body according to standard protocols well known in the art. The compositions can be introduced into the cells via any gene transfer mechanism, such as, for example, calcium phosphate mediated gene delivery, electroporation, microinjection or proteoliposomes. The transduced cells can then be infused (e.g., in a pharmaceutically acceptable carrier) or homotopically transplanted back into the subject per standard  
20 methods for the cell or tissue type. Standard methods are known for transplantation or infusion of various cells into a subject.

[94] If *in vivo* delivery methods are performed the methods can be designed to deliver the nucleic acid constructs directly to a particular cell type, via any delivery mechanism, such as intra-peritoneal injection of a vector construct. In this type of delivery  
25 situation, the nucleic acid constructs can be delivered to any type of tissue, for example, brain or neural or muscle. The nucleic acid constructs can also be delivered such that they generally deliver the nucleic acid constructs to more than one type of cell. This type of delivery can be accomplished, by for example, injecting the constructs intraperitoneally into the flank of the organism. (See Example 2 and figures 8-10). In certain delivery  
30 methods, the timing of the delivery is monitored. For example, the nucleic acid constructs can be delivered at the perinatal stage of the recipients life or at the adult stage.

[95] The disclosed compositions, can be delivered to any type of cell. For example, they can be delivered to any type of mammalian cell. Exemplary types of cells neuron, glia, fibroblast, chondrocyte, osteocyte, endothelial, and hepatocyte.

### 3. Expression systems

5 [96] The nucleic acids that are delivered to cells typically contain expression controlling systems. For example, the inserted genes in viral and retroviral systems usually contain promoters, and/or enhancers to help control the expression of the desired gene product. A promoter is generally a sequence or sequences of DNA that function when in a relatively fixed location in regard to the transcription start site. A promoter contains core  
10 elements required for basic interaction of RNA polymerase and transcription factors, and may contain upstream elements and response elements.

#### a) Viral Promoters and Enhancers

[97] Preferred promoters controlling transcription from vectors in mammalian host cells may be obtained from various sources, for example, the genomes of viruses such  
15 as: polyoma, Simian Virus 40 (SV40), adenovirus, retroviruses, hepatitis-B virus and most preferably cytomegalovirus, or from heterologous mammalian promoters, e.g. beta actin promoter. The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment which also contains the SV40 viral origin of replication (Fiers et al., Nature, 273: 113 (1978)). The immediate early promoter of the human  
20 cytomegalovirus is conveniently obtained as a HindIII E restriction fragment (Greenway, P.J. et al., Gene 18: 355-360 (1982)). Of course, promoters from the host cell or related species also are useful herein.

[98] Enhancer generally refers to a sequence of DNA that functions at no fixed distance from the transcription start site and can be either 5' (Laimins, L. et al., Proc. Natl. Acad. Sci. 78: 993 (1981)) or 3' (Lusky, M.L., et al., Mol. Cell Bio. 3: 1108 (1983)) to the  
25 transcription unit. Furthermore, enhancers can be within an intron (Banerji, J.L. et al., Cell 33: 729 (1983)) as well as within the coding sequence itself (Osborne, T.F., et al., Mol. Cell Bio. 4: 1293 (1984)). They are usually between 10 and 300 bp in length, and they function in cis. Enhancers function to increase transcription from nearby promoters.  
30 Enhancers also often contain response elements that mediate the regulation of transcription. Promoters can also contain response elements that mediate the regulation of transcription. Enhancers often determine the regulation of expression of a gene. While many enhancer

sequences are now known from mammalian genes (globin, elastase, albumin, -fetoprotein and insulin), typically one will use an enhancer from a eukaryotic cell virus for general expression. Preferred examples are the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer  
5 on the late side of the replication origin, and adenovirus enhancers.

[99] The promoter and/or enhancer may be specifically activated either by light or specific chemical events which trigger their function. Systems can be regulated by reagents such as tetracycline and dexamethasone. There are also ways to enhance viral vector gene expression by exposure to irradiation, such as gamma irradiation, or alkylating  
10 chemotherapy drugs.

[100] In certain embodiments the promoter and/or enhancer region can act as a constitutive promoter and/or enhancer to maximize expression of the region of the transcription unit to be transcribed. In certain constructs the promoter and/or enhancer region be active in all eukaryotic cell types, even if it is only expressed in a particular type  
15 of cell at a particular time. A preferred promoter of this type is the CMV promoter (650 bases). Other preferred promoters are SV40 promoters, cytomegalovirus (full length promoter), and retroviral vector LTF.

[101] It has been shown that all specific regulatory elements can be cloned and used to construct expression vectors that are selectively expressed in specific cell types such  
20 as melanoma cells. The glial fibrillary acetic protein (GFAP) promoter has been used to selectively express genes in cells of glial origin.

[102] Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human or nucleated cells) may also contain sequences necessary for the termination of transcription which may affect mRNA expression. These regions are transcribed as  
25 polyadenylated segments in the untranslated portion of the mRNA encoding tissue factor protein. The 3' untranslated regions also include transcription termination sites. It is preferred that the transcription unit also contain a polyadenylation region. One benefit of this region is that it increases the likelihood that the transcribed unit will be processed and transported like mRNA. The identification and use of polyadenylation signals in  
30 expression constructs is well established. It is preferred that homologous polyadenylation signals be used in the transgene constructs. In certain transcription units, the polyadenylation region is derived from the SV40 early polyadenylation signal and consists

of about 400 bases. It is also preferred that the transcribed units contain other standard sequences alone or in combination with the above sequences improve expression from, or stability of, the construct.

[103] In certain embodiments the promoters are constitutive promoters. This can be any promoter that causes transcription regulation in the absence of the addition of other factors. Examples of this type of promoter are the CMV promoter and the beta actin promoter, as well as others discussed herein. In certain embodiments the promoter can consist of fusions of one or more different types of promoters. For example, the regulatory regions of the CMV promoter and the beta actin promoter are well known and understood, examples, of which are disclosed herein. Parts of these promoters can be fused together to, for example, produce a CMV-beta actin fusion promoter, such as the one shown in SEQ ID NO:23. It is understood that this type of promoter has a CMV component and a beta actin component. These components can function independently as promoters, and thus, are themselves considered beta actin promoters and CMV promoters. A promoter can be any portion of a known promoter that causes promoter activity. It is well understood that many promoters, including the CMV and Beta Actin promoters have functional domains which are understood and that these can be used as a beta actin promoter or CMV promoter. Furthermore, these domains can be determined. For example, SEQ ID NO:s 21-41 display a number of CMV promoters, beta actin promoters, and fusion promoters. These promoters can be compared, and for example, functional regions delineated, as described herein. Furthermore, each of these sequences can function independently or together in any combination to provide a promoter region for the disclosed nucleic acids.

#### b) Markers

[104] The viral vectors can include nucleic acid sequence encoding a marker product. This marker product is used to determine if the gene has been delivered to the cell and once delivered is being expressed. Preferred marker genes are the *E. Coli lacZ* gene, which encodes  $\beta$ -galactosidase, and green fluorescent protein.

[105] In some embodiments the marker may be a selectable marker. Examples of suitable selectable markers for mammalian cells are dihydrofolate reductase (DHFR), thymidine kinase, neomycin, neomycin analog G418, hydromycin, and puromycin. When such selectable markers are successfully transferred into a mammalian host cell, the transformed mammalian host cell can survive if placed under selective pressure. There are

two widely used distinct categories of selective regimes. The first category is based on a cell's metabolism and the use of a mutant cell line which lacks the ability to grow independent of a supplemented media. Two examples are: CHO DHFR- cells and mouse LTK- cells. These cells lack the ability to grow without the addition of such nutrients as thymidine or hypoxanthine. Because these cells lack certain genes necessary for a complete nucleotide synthesis pathway, they cannot survive unless the missing nucleotides are provided in a supplemented media. An alternative to supplementing the media is to introduce an intact DHFR or TK gene into cells lacking the respective genes, thus altering their growth requirements. Individual cells which were not transformed with the DHFR or TK gene will not be capable of survival in non-supplemented media.

[106] The second category is dominant selection which refers to a selection scheme used in any cell type and does not require the use of a mutant cell line. These schemes typically use a drug to arrest growth of a host cell. Those cells which have a novel gene would express a protein conveying drug resistance and would survive the selection. Examples of such dominant selection use the drugs neomycin, (Southern P. and Berg, P., J. Molec. Appl. Genet. 1: 327 (1982)), mycophenolic acid, (Mulligan, R.C. and Berg, P. Science 209: 1422 (1980)) or hygromycin, (Sugden, B. et al., Mol. Cell. Biol. 5: 410-413 (1985)). The three examples employ bacterial genes under eukaryotic control to convey resistance to the appropriate drug G418 or neomycin (geneticin), xgpt (mycophenolic acid) or hygromycin, respectively. Others include the neomycin analog G418 and puramycin.

#### c) Post transcriptional regulatory elements

[107] The disclosed vectors can also contain post-transcriptional regulatory elements. Post-transcriptional regulatory elements can enhance mRNA stability or enhance translation of the transcribed mRNA. An exemplary post-transcriptional regulatory sequence is the WPRE sequence isolated from the woodchuck hepatitis virus. (Zufferey R, et al., "Woodchuck hepatitis virus post-transcriptional regulatory element enhances expression of transgenes delivered by retroviral vectors," J Virol; 73:2886-92 (1999)). Post-transcriptional regulatory elements can be positioned both 3' and 5' to the exogenous gene, but it is preferred that they are positioned 3' to the exogenous gene.

#### d) Transduction efficiency elements

[108] Transduction efficiency elements are sequences that enhance the packaging and transduction of the vector. These elements typically contain polypurine sequences. An

example of a transduction efficiency element is the ppt-cts sequence that contains the central polypurine tract (ppt) and central terminal site (cts) from the HIV-1 pSG3 molecular clone (SEQ ID NO:1 bp 4327 to 4483 of HIV-1 pSG3 clone).

e) 3' untranslated regions

5 [109] Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human or nucleated cells) may also contain sequences necessary for the termination of transcription which may affect mRNA expression. These 3' untranslated regions are transcribed as polyadenylated segments in the untranslated portion of the mRNA encoding the exogenous gene. The 3' untranslated regions also include transcription termination sites.  
10 The transcription unit also can contain a polyadenylation region. One benefit of this region is that it increases the likelihood that the transcribed unit will be processed and transported like mRNA. The identification and use of polyadenylation signals in expression constructs is well established. Homologous polyadenylation signals can be used in the transgene constructs. In an embodiment of the transcription unit, the polyadenylation region is  
15 derived from the SV40 early polyadenylation signal and consists of about 400 bases. Transcribed units can contain other standard sequences alone or in combination with the above sequences improve expression from, or stability of, the construct.

4. Sequence similarities

[110] It is understood that as discussed herein the use of the terms homology and  
20 identity mean the same thing as similarity. Thus, for example, if the use of the word homology is used between two non-natural sequences it is understood that this is not necessarily indicating an evolutionary relationship between these two sequences, but rather is looking at the similarity or relatedness between their nucleic acid sequences. Many of the methods for determining homology between two evolutionarily related molecules are  
25 routinely applied to any two or more nucleic acids or proteins for the purpose of measuring sequence similarity regardless of whether they are evolutionarily related or not.

[111] In general, it is understood that one way to define any known variants and derivatives or those that might arise, of the disclosed genes and proteins herein, is through defining the variants and derivatives in terms of homology to specific known sequences.  
30 This identity of particular sequences disclosed herein is also discussed elsewhere herein. In general, variants of genes and proteins herein disclosed typically have at least, about 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95,

96, 97, 98, or 99 percent homology to the stated sequence or the native sequence. Those of skill in the art readily understand how to determine the homology of two proteins or nucleic acids, such as genes. For example, the homology can be calculated after aligning the two sequences so that the homology is at its highest level.

5           [112] Another way of calculating homology can be performed by published algorithms. Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman *Adv. Appl. Math.* 2: 482 (1981), by the homology alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48: 443 (1970), by the search for similarity method of Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85: 10 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by inspection.

          [113] The same types of homology can be obtained for nucleic acids by for example the algorithms disclosed in Zuker, M. *Science* 244:48-52, 1989, Jaeger et al. *Proc.* 15 *Natl. Acad. Sci. USA* 86:7706-7710, 1989, Jaeger et al. *Methods Enzymol.* 183:281-306, 1989 which are herein incorporated by reference for at least material related to nucleic acid alignment. It is understood that any of the methods typically can be used and that in certain instances the results of these various methods may differ, but the skilled artisan understands if identity is found with at least one of these methods, the sequences would be said to have 20 the stated identity, and be disclosed herein.

          [114] For example, as used herein, a sequence recited as having a particular percent homology to another sequence refers to sequences that have the recited homology as calculated by any one or more of the calculation methods described above. For example, a first sequence has 80 percent homology, as defined herein, to a second sequence if the 25 first sequence is calculated to have 80 percent homology to the second sequence using the Zuker calculation method even if the first sequence does not have 80 percent homology to the second sequence as calculated by any of the other calculation methods. As another example, a first sequence has 80 percent homology, as defined herein, to a second sequence if the first sequence is calculated to have 80 percent homology to the second sequence using 30 both the Zuker calculation method and the Pearson and Lipman calculation method even if the first sequence does not have 80 percent homology to the second sequence as calculated by the Smith and Waterman calculation method, the Needleman and Wunsch calculation method, the Jaeger calculation methods, or any of the other calculation methods. As yet



another example, a first sequence has 80 percent homology, as defined herein, to a second sequence if the first sequence is calculated to have 80 percent homology to the second sequence using each of calculation methods (although, in practice, the different calculation methods will often result in different calculated homology percentages).

5                   **5. Hybridization/selective hybridization**

[115] The term hybridization typically means a sequence driven interaction between at least two nucleic acid molecules, such as a primer or a probe and a gene. Sequence driven interaction means an interaction that occurs between two nucleotides or nucleotide analogs or nucleotide derivatives in a nucleotide specific manner. For example, 10 G interacting with C or A interacting with T are sequence driven interactions. Typically sequence driven interactions occur on the Watson-Crick face or Hoogsteen face of the nucleotide. The hybridization of two nucleic acids is affected by a number of conditions and parameters known to those of skill in the art. For example, the salt concentrations, pH, and temperature of the reaction all affect whether two nucleic acid molecules will hybridize.

15

[116] Parameters for selective hybridization between two nucleic acid molecules are well known to those of skill in the art. For example, in some embodiments selective hybridization conditions can be defined as stringent hybridization conditions. For example, stringency of hybridization is controlled by both temperature and salt concentration of 20 either or both of the hybridization and washing steps. For example, the conditions of hybridization to achieve selective hybridization may involve hybridization in high ionic strength solution (6X SSC or 6X SSPE) at a temperature that is about 12-25°C below the T<sub>m</sub> (the melting temperature at which half of the molecules dissociate from their hybridization partners) followed by washing at a combination of temperature and salt 25 concentration chosen so that the washing temperature is about 5°C to 20°C below the T<sub>m</sub>. The temperature and salt conditions are readily determined empirically in preliminary experiments in which samples of reference DNA immobilized on filters are hybridized to a labeled nucleic acid of interest and then washed under conditions of different stringencies. Hybridization temperatures are typically higher for DNA-RNA and RNA-RNA 30 hybridizations. The conditions can be used as described above to achieve stringency, or as is known in the art. (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989; Kunkel et al. Methods Enzymol. 1987:154:367, 1987 which is herein incorporated by reference for

material at least related to hybridization of nucleic acids). A preferable stringent hybridization condition for a DNA:DNA hybridization can be at about 68°C (in aqueous solution) in 6X SSC or 6X SSPE followed by washing at 68°C. Stringency of hybridization and washing, if desired, can be reduced accordingly as the degree of complementarity  
5 desired is decreased, and further, depending upon the G-C or A-T richness of any area wherein variability is searched for. Likewise, stringency of hybridization and washing, if desired, can be increased accordingly as homology desired is increased, and further, depending upon the G-C or A-T richness of any area wherein high homology is desired, all as known in the art.

10 [117] Another way to define selective hybridization is by looking at the amount (percentage) of one of the nucleic acids bound to the other nucleic acid. For example, in some embodiments selective hybridization conditions would be when at least about, 60, 65, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 percent of the limiting nucleic acid is bound to the non-limiting  
15 nucleic acid. Typically, the non-limiting primer is in for example, 10 or 100 or 1000 fold excess. This type of assay can be performed at under conditions where both the limiting and non-limiting primer are for example, 10 fold or 100 fold or 1000 fold below their  $k_d$ , or where only one of the nucleic acid molecules is 10 fold or 100 fold or 1000 fold or where one or both nucleic acid molecules are above their  $k_d$ .

20 [118] Another way to define selective hybridization is by looking at the percentage of primer that gets enzymatically manipulated under conditions where hybridization is required to promote the desired enzymatic manipulation. For example, in some embodiments selective hybridization conditions would be when at least about, 60, 65, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94,  
25 95, 96, 97, 98, 99, 100 percent of the primer is enzymatically manipulated under conditions which promote the enzymatic manipulation, for example if the enzymatic manipulation is DNA extension, then selective hybridization conditions would be when at least about 60, 65, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 percent of the primer molecules are extended. Preferred  
30 conditions also include those suggested by the manufacturer or indicated in the art as being appropriate for the enzyme performing the manipulation.

[119] Just as with homology, it is understood that there are a variety of methods herein disclosed for determining the level of hybridization between two nucleic acid

molecules. It is understood that these methods and conditions may provide different percentages of hybridization between two nucleic acid molecules, but unless otherwise indicated meeting the parameters of any of the methods would be sufficient. For example if 80% hybridization was required and as long as hybridization occurs within the required parameters in any one of these methods it is considered disclosed herein.

[120] It is understood that those of skill in the art understand that if a composition or method meets any one of these criteria for determining hybridization either collectively or singly it is a composition or method that is disclosed herein.

#### 6. Nucleic acids

[121] There are a variety of molecules disclosed herein that are nucleic acid based, including for example the nucleic acids that encode, for example HexA and HexB, or functional nucleic acids. The disclosed nucleic acids can be made up of for example, nucleotides, nucleotide analogs, or nucleotide substitutes. Non-limiting examples of these and other molecules are discussed herein. It is understood that for example, when a vector is expressed in a cell, that the expressed mRNA will typically be made up of A, C, G, and U. Likewise, it is understood that if, for example, an antisense molecule is introduced into a cell or cell environment through for example exogenous delivery, it is advantageous that the antisense molecule be made up of nucleotide analogs that reduce the degradation of the antisense molecule in the cellular environment.

[122] A nucleotide is a molecule that contains a base moiety, a sugar moiety and a phosphate moiety. Nucleotides can be linked together through their phosphate moieties and sugar moieties creating an internucleoside linkage. The base moiety of a nucleotide can be adenin-9-yl (A), cytosin-1-yl (C), guanin-9-yl (G), uracil-1-yl (U), and thymin-1-yl (T). The sugar moiety of a nucleotide is a ribose or a deoxyribose. The phosphate moiety of a nucleotide is pentavalent phosphate. An non-limiting example of a nucleotide would be 3'-AMP (3'-adenosine monophosphate) or 5'-GMP (5'-guanosine monophosphate).

[123] A nucleotide analog is a nucleotide which contains some type of modification to either the base, sugar, or phosphate moieties. Modifications to nucleotides are well known in the art and would include for example, 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, and 2-aminoadenine as well as modifications at the sugar or phosphate moieties.

[124] Nucleotide substitutes are molecules having similar functional properties to nucleotides, but which do not contain a phosphate moiety, such as peptide nucleic acid (PNA). Nucleotide substitutes are molecules that will recognize nucleic acids in a Watson-Crick or Hoogsteen manner, but which are linked together through a moiety other than a phosphate moiety. Nucleotide substitutes are able to conform to a double helix type structure when interacting with the appropriate target nucleic acid.

[125] It is also possible to link other types of molecules (conjugates) to nucleotides or nucleotide analogs to enhance for example, cellular uptake. Conjugates can be chemically linked to the nucleotide or nucleotide analogs. Such conjugates include but are not limited to lipid moieties such as a cholesterol moiety. (Letsinger et al., Proc. Natl. Acad. Sci. USA, 1989,86, 6553-6556),

[126] A Watson-Crick interaction is at least one interaction with the Watson-Crick face of a nucleotide, nucleotide analog, or nucleotide substitute. The Watson-Crick face of a nucleotide, nucleotide analog, or nucleotide substitute includes the C2, N1, and C6 positions of a purine based nucleotide, nucleotide analog, or nucleotide substitute and the C2, N3, C4 positions of a pyrimidine based nucleotide, nucleotide analog, or nucleotide substitute.

[127] A Hoogsteen interaction is the interaction that takes place on the Hoogsteen face of a nucleotide or nucleotide analog, which is exposed in the major groove of duplex DNA. The Hoogsteen face includes the N7 position and reactive groups (NH2 or O) at the C6 position of purine nucleotides.

#### a) Sequences

[128] There are a variety of sequences related to the HexA, HexB, IRES sequences, and promoter sequences. For example, the HexA and hexB genes have the following Genbank Accession Numbers: M16411 and NM\_000520 for HexA and NM\_000521 for HexB, these sequences and others are herein incorporated by reference in their entireties as well as for individual subsequences contained therein. It is understood that there are numerous Genbank accession sequences related to HexA and HexB, all of which are incorporated by reference herein.

[129] One particular sequence set forth in SEQ ID NO:4 and having Genbank accession number NM\_000521, which is a sequence for human HexB cDNA, is used herein, as an example, to exemplify the disclosed compositions and methods. It is

understood that the description related to this sequence is applicable to any sequence related to HexA or HexB unless specifically indicated otherwise. Those of skill in the art understand how to resolve sequence discrepancies and differences and to adjust the compositions and methods relating to a particular sequence to other related sequences.

- 5 Primers and/or probes can be designed for any of the sequences disclosed herein given the information disclosed herein and that known in the art.

[130] It is also understood for example that there are numerous bicistronic vectors that can be used to create the  $\beta$ -Hex construct nucleic acids See for example, Genbank accession no Y11035 and Y11034.

10

**b) Primers and probes**

[131] Disclosed are compositions including primers and probes, which are capable of interacting with, for example, the  $\beta$ -Hex construct nucleic acids, as disclosed herein. In certain embodiments the primers are used to support DNA amplification reactions.

Typically the primers will be capable of being extended in a sequence specific manner.

- 15 Extension of a primer in a sequence specific manner includes any methods wherein the sequence and/or composition of the nucleic acid molecule to which the primer is hybridized or otherwise associated directs or influences the composition or sequence of the product produced by the extension of the primer. Extension of the primer in a sequence specific manner therefore includes, but is not limited to, PCR, DNA sequencing, DNA extension,
- 20 DNA polymerization, RNA transcription, or reverse transcription. Techniques and conditions that amplify the primer in a sequence specific manner are preferred. In certain embodiments the primers are used for the DNA amplification reactions, such as PCR or direct sequencing. It is understood that in certain embodiments the primers can also be extended using non-enzymatic techniques, where for example, the nucleotides or
- 25 oligonucleotides used to extend the primer are modified such that they will chemically react to extend the primer in a sequence specific manner. Typically the disclosed primers hybridize with, for example, the  $\beta$ -Hex construct nucleic acid, or region of the  $\beta$ -Hex construct nucleic acids or they hybridize with the complement of the  $\beta$ -Hex construct nucleic acids or complement of a region of the  $\beta$ -Hex construct nucleic acids.

## 7. Peptides

### a) Protein variants

[132] As discussed herein there are numerous variants of the HEX- $\alpha$  and HEX- $\beta$  proteins that are known and herein contemplated. In addition, to the known functional species and allelic variants of HEX- $\alpha$  and HEX- $\beta$  there are derivatives of the HEX- $\alpha$  and HEX- $\beta$  proteins which also function in the disclosed methods and compositions. Protein variants and derivatives are well understood to those of skill in the art and in can involve amino acid sequence modifications. For example, amino acid sequence modifications typically fall into one or more of three classes: substitutional, insertional or deletional variants. Insertions include amino and/or carboxyl terminal fusions as well as intrasequence insertions of single or multiple amino acid residues. Insertions ordinarily will be smaller insertions than those of amino or carboxyl terminal fusions, for example, on the order of one to four residues. Immunogenic fusion protein derivatives, such as those described in the examples, are made by fusing a polypeptide sufficiently large to confer immunogenicity to the target sequence by cross-linking in vitro or by recombinant cell culture transformed with DNA encoding the fusion. Deletions are characterized by the removal of one or more amino acid residues from the protein sequence. Typically, no more than about from 2 to 6 residues are deleted at any one site within the protein molecule. These variants ordinarily are prepared by site specific mutagenesis of nucleotides in the DNA encoding the protein, thereby producing DNA encoding the variant, and thereafter expressing the DNA in recombinant cell culture. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence are well known, for example M13 primer mutagenesis and PCR mutagenesis. Amino acid substitutions are typically of single residues, but can occur at a number of different locations at once; insertions usually will be on the order of about from 1 to 10 amino acid residues; and deletions will range about from 1 to 30 residues. Deletions or insertions preferably are made in adjacent pairs, i.e. a deletion of 2 residues or insertion of 2 residues. Substitutions, deletions, insertions or any combination thereof may be combined to arrive at a final construct. The mutations must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary mRNA structure. Substitutional variants are those in which at least one residue has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the following Tables 1 and 2 and are referred to as conservative substitutions.

[133] TABLE 1: Amino Acid Abbreviations

Amino Acid	Abbreviations
alanine	AlaA
allosoleucine	Alle
arginine	ArgR
asparagine	AsnN
aspartic acid	AspD
cysteine	CysC
glutamic acid	GluE
glutamine	GlnK
glycine	GlyG
histidine	HisH
isoleucine	IleI
leucine	LeuL
lysine	LysK
phenylalanine	PheF
proline	ProP
pyroglutamic acidp	Glu
serine	SerS
threonine	ThrT
tyrosine	TyrY
tryptophan	TrpW
valine	ValV

TABLE 2: Amino Acid Substitutions

Original Residue Exemplary Conservative Substitutions, others are known in the art.	
Ala	ser
Arg	lys, gln
Asn	gln; his
Asp	glu
Cys	ser
Gln	asn, lys
Glu	asp
Gly	pro
His	asn; gln
Ile	leu; val
Leu	ile; val
Lys	arg; gln;
Met	Leu; ile
Phe	met; leu; tyr
Ser	thr
Thr	ser
Trp	tyr
Tyr	trp; phe
Val	ile; leu

[134] Substantial changes in function or immunological identity are made by selecting substitutions that are less conservative than those in Table 2, i.e., selecting residues that differ more significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site or (c) the

bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in the protein properties will be those in which (a) a hydrophilic residue, e.g. seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g. leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other  
5 residue; (c) a residue having an electropositive side chain, e.g., lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine, in this case, (e) by increasing the number of sites for sulfation and/or glycosylation.

10 [135] For example, the replacement of one amino acid residue with another that is biologically and/or chemically similar is known to those skilled in the art as a conservative substitution. For example, a conservative substitution would be replacing one hydrophobic residue for another, or one polar residue for another. The substitutions include combinations such as, for example, Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr;  
15 Lys, Arg; and Phe, Tyr. Such conservatively substituted variations of each explicitly disclosed sequence are included within the mosaic polypeptides provided herein.

[136] Substitutional or deletional mutagenesis can be employed to insert sites for N-glycosylation (Asn-X-Thr/Ser) or O-glycosylation (Ser or Thr). Deletions of cysteine or other labile residues also may be desirable. Deletions or substitutions of potential  
20 proteolysis sites, e.g. Arg, is accomplished for example by deleting one of the basic residues or substituting one by glutaminyl or histidyl residues.

[137] Certain post-translational derivatizations are the result of the action of recombinant host cells on the expressed polypeptide. Glutaminyl and asparaginyl residues are frequently post-translationally deamidated to the corresponding glutamyl and asparyl  
25 residues. Alternatively, these residues are deamidated under mildly acidic conditions. Other post-translational modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the o-amino groups of lysine, arginine, and histidine side chains (T.E. Creighton, Proteins: Structure and Molecular Properties, W. H. Freeman & Co., San Francisco pp 79-86 [1983]),  
30 acetylation of the N-terminal amine and, in some instances, amidation of the C-terminal carboxyl.



[138] It is understood that one way to define the variants and derivatives of the disclosed proteins herein is through defining the variants and derivatives in terms of homology/identity to specific known sequences. For example, SEQ ID NO:1 sets forth a particular sequence of HEX- $\alpha$  and SEQ ID NO:3 sets forth a particular sequence of a HEX-  
5  $\beta$  protein. Specifically disclosed are variants of these and other proteins herein disclosed which have at least, 70% or 75% or 80% or 85% or 90% or 95% homology to the stated sequence. Those of skill in the art readily understand how to determine the homology of two proteins. For example, the homology can be calculated after aligning the two sequences so that the homology is at its highest level.

10 [139] Another way of calculating homology can be performed by published algorithms. Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman *Adv. Appl. Math.* 2: 482 (1981), by the homology alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48: 443 (1970), by the search for similarity method of Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:  
15 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by inspection.

[140] The same types of homology can be obtained for nucleic acids by for example the algorithms disclosed in Zuker, M. *Science* 244:48-52, 1989, Jaeger et al. *Proc.*  
20 *Natl. Acad. Sci. USA* 86:7706-7710, 1989, Jaeger et al. *Methods Enzymol.* 183:281-306, 1989 which are herein incorporated by reference for at least material related to nucleic acid alignment.

[141] It is understood that the description of conservative mutations and homology can be combined together in any combination, such as embodiments that have at least 70%  
25 homology to a particular sequence wherein the variants are conservative mutations.

[142] As this specification discusses various proteins and protein sequences it is understood that the nucleic acids that can encode those protein sequences are also disclosed. This would include all degenerate sequences related to a specific protein sequence, i.e. all nucleic acids having a sequence that encodes one particular protein sequence as well as all  
30 nucleic acids, including degenerate nucleic acids, encoding the disclosed variants and derivatives of the protein sequences. Thus, while each particular nucleic acid sequence may not be written out herein, it is understood that each and every sequence is in fact disclosed

and described herein through the disclosed protein sequence. For example, one of the many nucleic acid sequences that can encode the protein sequence set forth in SEQ ID NO:3 is set forth in SEQ ID NO:4. Another nucleic acid sequence that encodes the same protein sequence set forth in SEQ ID NO:3 is set forth in SEQ ID NO:11. In addition, for example, 5 a disclosed conservative derivative of SEQ ID NO:3 is shown in SEQ ID NO: 12, where the valine (V) at position 21 is changed to a isoleucine (I). It is understood that for this mutation all of the nucleic acid sequences that encode this particular derivative of the SEQ ID NO:3 polypeptide are also disclosed. It is also understood that while no amino acid sequence indicates what particular DNA sequence encodes that protein within an organism, 10 where particular variants of a disclosed protein are disclosed herein, the known nucleic acid sequence that encodes that protein in the particular organism from which that protein arises is also known and herein disclosed and described.

#### 8. Pharmaceutical carriers/Delivery of pharmaceutical products

[143] As described above, the compositions can also be administered *in vivo* in a 15 pharmaceutically acceptable carrier. By "pharmaceutically acceptable" is meant a material that is not biologically or otherwise undesirable, i.e., the material may be administered to a subject, along with the nucleic acid or vector, without causing any undesirable biological effects or interacting in a deleterious manner with any of the other components of the pharmaceutical composition in which it is contained. The carrier would naturally be 20 selected to minimize any degradation of the active ingredient and to minimize any adverse side effects in the subject, as would be well known to one of skill in the art.

[144] The compositions may be administered orally, parenterally (e.g., intravenously), by intramuscular injection, by intraperitoneal injection, transdermally, extracorporeally, topically or the like, including topical intranasal administration or 25 administration by inhalant. As used herein, "topical intranasal administration" means delivery of the compositions into the nose and nasal passages through one or both of the nares and can comprise delivery by a spraying mechanism or droplet mechanism, or through aerosolization of the nucleic acid or vector. Administration of the compositions by inhalant can be through the nose or mouth via delivery by a spraying or droplet mechanism. 30 Delivery can also be directly to any area of the respiratory system (e.g., lungs) via intubation. The exact amount of the compositions required will vary from subject to subject, depending on the species, age, weight and general condition of the subject, the severity of the allergic disorder being treated, the particular nucleic acid or vector used, its

mode of administration and the like. Thus, it is not possible to specify an exact amount for every composition. However, an appropriate amount can be determined by one of ordinary skill in the art using only routine experimentation given the teachings herein.

[145] Parenteral administration of the composition, if used, is generally  
5 characterized by injection. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. A more recently revised approach for parenteral administration involves use of a slow release or sustained release system such that a constant dosage is maintained. See, e.g., U.S. Patent No. 3,610,795, which is incorporated  
10 by reference herein.

[146] The materials may be in solution, suspension (for example, incorporated into microparticles, liposomes, or cells). These may be targeted to a particular cell type via antibodies, receptors, or receptor ligands. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Senter, et al., Bioconjugate Chem., 2:447-451, (1991); Bagshawe, K.D., Br. J. Cancer, 60:275-281, (1989); Bagshawe, et al., Br. J. Cancer, 58:700-703, (1988); Senter, et al., Bioconjugate Chem., 4:3-9, (1993); Battelli, et al., Cancer Immunol. Immunother., 35:421-425, (1992); Pietersz and McKenzie, Immunolog. Reviews, 129:57-80, (1992); and Roffler, et al., Biochem. Pharmacol., 42:2062-2065, (1991)). Vehicles such as "stealth" and other antibody conjugated liposomes  
20 (including lipid mediated drug targeting to colonic carcinoma), receptor mediated targeting of DNA through cell specific ligands, lymphocyte directed tumor targeting, and highly specific therapeutic retroviral targeting of murine glioma cells *in vivo*. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Hughes et al., Cancer Research, 49:6214-6220, (1989); and Litzinger and Huang,  
25 Biochimica et Biophysica Acta, 1104:179-187, (1992)). In general, receptors are involved in pathways of endocytosis, either constitutive or ligand induced. These receptors cluster in clathrin-coated pits, enter the cell via clathrin-coated vesicles, pass through an acidified endosome in which the receptors are sorted, and then either recycle to the cell surface, become stored intracellularly, or are degraded in lysosomes. The internalization pathways  
30 serve a variety of functions, such as nutrient uptake, removal of activated proteins, clearance of macromolecules, opportunistic entry of viruses and toxins, dissociation and degradation of ligand, and receptor-level regulation. Many receptors follow more than one intracellular pathway, depending on the cell type, receptor concentration, type of ligand,

ligand valency, and ligand concentration. Molecular and cellular mechanisms of receptor-mediated endocytosis has been reviewed (Brown and Greene, DNA and Cell Biology 10:6, 399-409 (1991)).

**a) Pharmaceutically Acceptable Carriers**

5           [147] The compositions, including antibodies, can be used therapeutically in combination with a pharmaceutically acceptable carrier.

          [148] Suitable carriers and their formulations are described in *Remington: The Science and Practice of Pharmacy* (19th ed.) ed. A.R. Gennaro, Mack Publishing Company, Easton, PA 1995. Typically, an appropriate amount of a pharmaceutically-acceptable salt is  
10       used in the formulation to render the formulation isotonic. Examples of the pharmaceutically-acceptable carrier include, but are not limited to, saline, Ringer's solution and dextrose solution. The pH of the solution is preferably from about 5 to about 8, and more preferably from about 7 to about 7.5. Further carriers include sustained release preparations such as semipermeable matrices of solid hydrophobic polymers containing the  
15       antibody, which matrices are in the form of shaped articles, e.g., films, liposomes or microparticles. It will be apparent to those persons skilled in the art that certain carriers may be more preferable depending upon, for instance, the route of administration and concentration of composition being administered.

          [149] Pharmaceutical carriers are known to those skilled in the art. These most  
20       typically would be standard carriers for administration of drugs to humans, including solutions such as sterile water, saline, and buffered solutions at physiological pH. The compositions can be administered intramuscularly or subcutaneously. Other compounds will be administered according to standard procedures used by those skilled in the art.

          [150] Pharmaceutical compositions may include carriers, thickeners, diluents, buffers,  
25       preservatives, surface active agents and the like in addition to the molecule of choice. Pharmaceutical compositions may also include one or more active ingredients such as antimicrobial agents, antiinflammatory agents, anesthetics, and the like.

          [151] The pharmaceutical composition may be administered in a number of ways depending on whether local or systemic treatment is desired, and on the area to be treated.  
30       Administration may be topically (including ophthalmically, vaginally, rectally, intranasally), orally, by inhalation, or parenterally, for example by intravenous drip, subcutaneous, intraperitoneal or intramuscular injection. The disclosed antibodies can be administered

intravenously, intraperitoneally, intramuscularly, subcutaneously, intracavity, or transdermally.

[152] Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

[153] Formulations for topical administration may include ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable.

[154] Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets, or tablets. Thickeners, flavorings, diluents, emulsifiers, dispersing aids or binders may be desirable.

[155] Some of the compositions may potentially be administered as a pharmaceutically acceptable acid- or base- addition salt, formed by reaction with inorganic acids such as hydrochloric acid, hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, and phosphoric acid, and organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, and fumaric acid, or by reaction with an inorganic base such as sodium hydroxide, ammonium hydroxide, potassium hydroxide, and organic bases such as mono-, di-, trialkyl and aryl amines and substituted ethanolamines.

#### 9. Chips and micro arrays

[156] Disclosed are chips where at least one address is the sequences or part of the sequences set forth in any of the nucleic acid sequences disclosed herein. Also disclosed are chips where at least one address is the sequences or portion of sequences set forth in any of the peptide sequences disclosed herein.

[157] Also disclosed are chips where at least one address is a variant of the sequences or part of the sequences set forth in any of the nucleic acid sequences disclosed herein. Also disclosed are chips where at least one address is a variant of the sequences or portion of sequences set forth in any of the peptide sequences disclosed herein.

5                   **10. Computer readable mediums**

[158] It is understood that the disclosed nucleic acids and proteins can be represented as a sequence consisting of the nucleotides of amino acids. There are a variety of ways to display these sequences, for example the nucleotide guanosine can be represented by G or g. Likewise the amino acid valine can be represented by Val or V.

10 Those of skill in the art understand how to display and express any nucleic acid or protein sequence in any of the variety of ways that exist, each of which is considered herein disclosed. Specifically contemplated herein is the display of these sequences on computer readable mediums, such as, commercially available floppy disks, tapes, chips, hard drives, compact disks, and video disks, or other computer readable mediums. Also disclosed are

15 the binary code representations of the disclosed sequences. Those of skill in the art understand what computer readable mediums. Thus, computer readable mediums on which the nucleic acids or protein sequences are recorded, stored, or saved.

[159] Disclosed are computer readable mediums comprising the sequences and information regarding the sequences set forth herein.

20                   **11. Kits**

[160] Disclosed herein are kits that are drawn to reagents that can be used in practicing the methods disclosed herein. The kits can include any reagent or combination of reagent discussed herein or that would be understood to be required or beneficial in the practice of the disclosed methods. For example, the kits could include primers to perform

25 the amplification reactions discussed in certain embodiments of the methods, as well as the buffers and enzymes required to use the primers as intended.

**D. Methods of making the compositions**

[161] The compositions disclosed herein and the compositions necessary to perform the disclosed methods can be made using any method known to those of skill in the

30 art for that particular reagent or compound unless otherwise specifically noted.

[162] The disclosed viral vectors can be made using standard recombinant molecular biology techniques. Many of these techniques are illustrated in Maniatis

(Maniatis et al., "*Molecular Cloning--A Laboratory Manual*," (Cold Spring Harbor Laboratory, Latest edition) and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989.

#### 1. Nucleic acid synthesis

5           [163] For example, the nucleic acids, such as, the oligonucleotides to be used as primers can be made using standard chemical synthesis methods or can be produced using enzymatic methods or any other known method. Such methods can range from standard enzymatic digestion followed by nucleotide fragment isolation (see for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edition (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989) Chapters 5, 6) to purely synthetic methods, for example, by the cyanoethyl phosphoramidite method using a Milligen or Beckman System Plus DNA synthesizer (for example, Model 8700 automated synthesizer of Milligen-Bioscience, Burlington, MA or ABI Model 380B). Synthetic methods useful for making oligonucleotides are also described by Ikuta et al., *Ann. Rev. Biochem.* 53:323-356  
10           (1984), (phosphotriester and phosphite-triester methods), and Narang et al., *Methods Enzymol.*, 65:610-620 (1980), (phosphotriester method). Protein nucleic acid molecules can be made using known methods such as those described by Nielsen et al., *Bioconjug. Chem.* 5:3-7 (1994).

#### 2. Peptide synthesis

20           [164] One method of producing the disclosed proteins is to link two or more peptides or polypeptides together by protein chemistry techniques. For example, peptides or polypeptides can be chemically synthesized using currently available laboratory equipment using either Fmoc (9-fluorenylmethyloxycarbonyl) or Boc (*tert*-butyloxycarbonyl) chemistry. (Applied Biosystems, Inc., Foster City, CA). One skilled  
25           in the art can readily appreciate that a peptide or polypeptide corresponding to the disclosed proteins, for example, can be synthesized by standard chemical reactions. For example, a peptide or polypeptide can be synthesized and not cleaved from its synthesis resin whereas the other fragment of a peptide or protein can be synthesized and subsequently cleaved from the resin, thereby exposing a terminal group which is functionally blocked on the other  
30           fragment. By peptide condensation reactions, these two fragments can be covalently joined via a peptide bond at their carboxyl and amino termini, respectively, to form an antibody, or fragment thereof. (Grant GA (1992) *Synthetic Peptides: A User Guide*. W.H. Freeman and Co., N.Y. (1992); Bodansky M and Trost B., Ed. (1993) *Principles of Peptide Synthesis*.

Springer-Verlag Inc., NY (which is herein incorporated by reference at least for material related to peptide synthesis). Alternatively, the peptide or polypeptide is independently synthesized *in vivo* as described herein. Once isolated, these independent peptides or polypeptides may be linked to form a peptide or fragment thereof via similar peptide condensation reactions.

[165] For example, enzymatic ligation of cloned or synthetic peptide segments allow relatively short peptide fragments to be joined to produce larger peptide fragments, polypeptides or whole protein domains (Abrahmsen L et al., Biochemistry, 30:4151 (1991)). Alternatively, native chemical ligation of synthetic peptides can be utilized to synthetically construct large peptides or polypeptides from shorter peptide fragments. This method consists of a two step chemical reaction (Dawson et al. Synthesis of Proteins by Native Chemical Ligation. Science, 266:776-779 (1994)). The first step is the chemoselective reaction of an unprotected synthetic peptide--thioester with another unprotected peptide segment containing an amino-terminal Cys residue to give a thioester-linked intermediate as the initial covalent product. Without a change in the reaction conditions, this intermediate undergoes spontaneous, rapid intramolecular reaction to form a native peptide bond at the ligation site (Baggiolini M et al. (1992) FEBS Lett. 307:97-101; Clark-Lewis I et al., J.Biol.Chem., 269:16075 (1994); Clark-Lewis I et al., Biochemistry, 30:3128 (1991); Rajarathnam K et al., Biochemistry 33:6623-30 (1994)).

[166] Alternatively, unprotected peptide segments are chemically linked where the bond formed between the peptide segments as a result of the chemical ligation is an unnatural (non-peptide) bond (Schnolzer, M et al. Science, 256:221 (1992)). This technique has been used to synthesize analogs of protein domains as well as large amounts of relatively pure proteins with full biological activity (deLisle Milton RC et al., Techniques in Protein Chemistry IV. Academic Press, New York, pp. 257-267 (1992)).

### 3. Processes for making the compositions

[167] Disclosed are processes for making the compositions as well as making the intermediates leading to the compositions. There are a variety of methods that can be used for making these compositions, such as synthetic chemical methods and standard molecular biology methods. It is understood that the methods of making these and the other disclosed compositions are specifically disclosed.



[168] Disclosed are nucleic acid molecules produced by the process comprising linking in an operative way a promoter element, a HexB element, a IRES element, and a HexA element.

5 [169] Disclosed are nucleic acid molecules produced by the process comprising linking in an operative way nucleic acid molecules comprising sequences set forth in SEQ ID NO:10 and SEQ ID NO:4.

[170] Also disclosed are nucleic acid molecules produced by the process comprising linking in an operative way nucleic acid molecules comprising sequences having 80% identity to sequences set forth in SEQ ID NO:10 and SEQ ID NO:4.

10 [171] Also disclosed are nucleic acid molecules produced by the process comprising linking in an operative way nucleic acid molecules comprising sequences that hybridizes under stringent hybridization conditions to sequences set forth in SEQ ID NO:10 and SEQ ID NO:4.

[172] Disclosed are nucleic acid molecules produced by the process comprising  
15 linking in an operative way a nucleic acid molecule comprising a sequence encoding HEX- $\beta$  and HEX- $\alpha$  peptides and a sequence controlling an expression of the sequence encoding HEX- $\beta$  and HEX- $\alpha$ .

[173] Disclosed are nucleic acid molecules produced by the process comprising  
20 linking in an operative way a nucleic acid molecule comprising a sequence encoding HEX- $\beta$  and HEX- $\alpha$  peptides wherein the HEX- $\beta$  and HEX- $\alpha$  peptides have 80% identity to the peptides set forth in SEQ ID NO:1 and SEQ ID NO:3 and a sequence controlling expression of the sequences encoding the peptides.

[174] Disclosed are nucleic acid molecules produced by the process comprising  
25 linking in an operative way a nucleic acid molecule comprising a sequence encoding HEX- $\beta$  and HEX- $\alpha$  peptides wherein the HEX- $\beta$  and HEX- $\alpha$  peptides have 80% identity to the peptides set forth in SEQ ID NO:1 and SEQ ID NO:3, wherein any change from the sequences set forth in SEQ ID NO:1 and SEQ ID NO:3 are conservative changes and a sequence controlling expression of the sequences encoding the peptides.

[175] Disclosed are cells produced by the process of transforming the cell with any  
30 of the disclosed nucleic acids. Disclosed are cells produced by the process of transforming the cell with any of the non-naturally occurring disclosed nucleic acids.

[176] Disclosed are any of the disclosed peptides produced by the process of expressing any of the disclosed nucleic acids. Disclosed are any of the non-naturally occurring disclosed peptides produced by the process of expressing any of the disclosed nucleic acids. Disclosed are any of the disclosed peptides produced by the process of  
5 expressing any of the non-naturally disclosed nucleic acids.

[177] Disclosed are animals produced by the process of transfecting a cell within the animal with any of the nucleic acid molecules disclosed herein. Disclosed are animals produced by the process of transfecting a cell within the animal any of the nucleic acid molecules disclosed herein, wherein the animal is a mammal. Also disclosed are animals  
10 produced by the process of transfecting a cell within the animal any of the nucleic acid molecules disclosed herein, wherein the mammal is mouse, rat, rabbit, cow, sheep, pig, or primate. Also disclosed are mammals wherein mammal is a murine, ungulate, or non-human primate.

[178] Also disclose are animals produced by the process of adding to the animal  
15 any of the cells disclosed herein.

#### **E. Methods of using the compositions**

##### **1. Methods of using the compositions as research tools**

[179] The disclosed compositions can be used in a variety of ways as research tools. For example, the disclosed compositions, the  $\beta$ -Hex constructs, and other nucleic  
20 acids, such as SEQ ID NOs:10 and 4 can be used to produce organisms, such as transgenic or knockout mice, which can be used as model systems for the study of Tay Sachs and Sandoffs disease.

##### **2. Methods of gene modification and gene disruption**

[180] The disclosed compositions and methods can be used for targeted gene  
25 disruption and modification in any animal that can undergo these events. Gene modification and gene disruption refer to the methods, techniques, and compositions that surround the selective removal or alteration of a gene or stretch of chromosome in an animal, such as a mammal, in a way that propagates the modification through the germ line of the mammal. In general, a cell is transformed with a vector which is designed to  
30 homologously recombine with a region of a particular chromosome contained within the cell, as for example, described herein. This homologous recombination event can produce a chromosome which has exogenous DNA introduced, for example in frame, with the

surrounding DNA. This type of protocol allows for very specific mutations, such as point mutations, to be introduced into the genome contained within the cell. Methods for performing this type of homologous recombination are disclosed herein.

[181] One of the preferred characteristics of performing homologous recombination in mammalian cells is that the cells should be able to be cultured, because the desired recombination event occurs at a low frequency.

[182] Once the cell is produced through the methods described herein, an animal can be produced from this cell through either stem cell technology or cloning technology. For example, if the cell into which the nucleic acid was transfected was a stem cell for the organism, then this cell, after transfection and culturing, can be used to produce an organism which will contain the gene modification or disruption in germ line cells, which can then in turn be used to produce another animal that possesses the gene modification or disruption in all of its cells. In other methods for production of an animal containing the gene modification or disruption in all of its cells, cloning technologies can be used. These technologies generally take the nucleus of the transfected cell and either through fusion or replacement fuse the transfected nucleus with an oocyte which can then be manipulated to produce an animal. The advantage of procedures that use cloning instead of ES technology is that cells other than ES cells can be transfected. For example, a fibroblast cell, which is very easy to culture can be used as the cell which is transfected and has a gene modification or disruption event take place, and then cells derived from this cell can be used to clone a whole animal.

### 3. Therapeutic Uses

[183] Effective dosages and schedules for administering the compositions may be determined empirically, and making such determinations is within the skill in the art. The dosage ranges for the administration of the compositions are those large enough to produce the desired effect in which the symptoms disorder are effected. The dosage should not be so large as to cause adverse side effects, such as unwanted cross-reactions, anaphylactic reactions, and the like. Generally, the dosage will vary with the age, condition, sex and extent of the disease in the patient, route of administration, or whether other drugs are included in the regimen, and can be determined by one of skill in the art. The dosage can be adjusted by the individual physician in the event of any counterindications. Dosage can vary, and can be administered in one or more dose administrations daily, for one or several

days. Guidance can be found in the literature for appropriate dosages for given classes of pharmaceutical products.

[184] Following administration of a disclosed composition, such as the disclosed constructs, for treating, inhibiting, or preventing Tay Sachs or Sandoffs disease, the efficacy of the therapeutic construct can be assessed in various ways well known to the skilled practitioner. For instance, one of ordinary skill in the art will understand that a composition, such as the disclosed constructs, disclosed herein is efficacious in treating Tay Sachs or Sandoffs disease or inhibiting or reducing the effects of Tay Sachs or Sandoffs disease in a subject by observing that the composition reduces the onset of the conditions associated with these diseases. Furthermore, the amount of protein or transcript produced from the constructs can be analyzed using any diagnostic method. For example, it can be measured using polymerase chain reaction assays to detect the presence of construct nucleic acid or antibody assays to detect the presence of protein produced from the construct in a sample (e.g., but not limited to, blood or other cells, such as neural cells) from a subject or patient.

#### F. Examples

[185] It will be apparent to those skilled in the art that various modifications and variations can be made in the present invention without departing from the scope or spirit of the invention. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

[186] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compounds, compositions, articles, devices and/or methods claimed herein are made and evaluated, and are intended to be purely exemplary of the invention and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.), but some errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in °C or is at ambient temperature, and pressure is at or near atmospheric.

## 1. Example 1 Making $\beta$ -Hex constructs

### a) Construction of bicistronic $\beta$ -Hex construct

[187] A bicistronic construct encoding for both isoforms of human  $\beta$ -hexosaminidase, hHexA and hHexB was made (Figure 1). hHexB cDNA was isolated following *Xho* I digestion of pHexB43 (ATCC, Manassas VA) and cloned into the *Xho* I site of pIRES (Clonotech Laboratories, Palo Alto CA) downstream of the vector's cytomegalovirus (CMV) promoter sequence. The HexA cDNA was isolated from pBHA-5' (ATCC, Manassas VA) by *Xho* I digestion and was subsequently inserted into the *Xba* I site of pIRES(HexB) downstream of the vectors IRES cassette by blunt ligation. In this construct, the cytomegalovirus promoter (CMV) drives transgene expression, and the translation of the second open reading frame, *HexB*, is facilitated by an internal ribosomal entry sequence (IRES).

### b) Results

[188] The HEX*lacZ* encodes for both isoforms of human  $\beta$ -hexosaminidase, HexA & HexB. (Figure 1) The vector pHEX*lacZ* is shown in Figure 1(A). BHK<sup>Hex*lacZ*</sup> are developed by stable *HexlacZ* transduction. Figure 1(B) shows that the cells transfected with the pHEX*lacZ* vector stain positively by X-gal histochemistry. Furthermore, HexA & HexB mRNA was detected by RT-PCR in total RNA extracts (Figure 1(C)). Likewise, not only was transcript of pHEX*lacZ* vector identified, human HEXA and human HEXB proteins were detected in the transfected BHK<sup>Hex*lacZ*</sup> cells by immunocytochemistry. (Figure 1(D<sub>1</sub>) and 1(E<sub>1</sub>)). This data indicates that the disclosed constructs can be expressed in target cells and that sufficient levels of protein are produced within these cells.

[189] The  $\beta$ -Hex therapeutic gene is capable of correcting deficiencies in cells that are not transfected through cross-correction. (Figure 2) An important property of the  $\beta$ -Hex transgene is the products hHEXA & hHEXB have the ability to cross-correct, specifically, to be released extracellularly and then to be absorbed via paracrine pathways by other cells whereby they contribute to  $\beta$ -hexosaminidase activity. BHK<sup>Hex*lacZ*</sup> cells were cultured and the supernatant was collected (conditioned medium), filtered (.45mm) and applied on normal mouse kidney fibroblasts in culture. Forty-eight hours later, the cells were washed thoroughly with phosphate buffered saline, and briefly treated with a trypsin solution to remove extracellular proteins from the cell surfaces. Following trypsin inactivation with Tris/EDTA buffer, the cells were fixed with 4% paraformaldehyde

solution and processed by Fast Garnet histochemistry for  $\beta$ -hexosaminidase activity. Fast Garnet histochemistry of murine fibroblasts exposed to (Figure 2A) conditioned medium collected from BHK<sup>HexlacZ</sup> cells compared to cells exposed to medium from normal parent BHK-21 cells (Figure 2B). These results demonstrate that hHEXA & hHEXB, products of the  $\beta$ -Hex transgene, are released into the extracellular medium and can be absorbed by other cells via paracrine pathways resulting in induction of the cellular  $\beta$ -hexosaminidase.

## 2. Example 2 Transfecting constructs

### a) Construction of the tricistronic $\beta$ -Hex construct

[190] A tricistronic construct encoding for both isoforms of human  $\beta$ -hexosaminidase, hHexA & hHexB, as well as the  $\beta$ -galactosidase reporter gene (*lacZ*) was also made. hHexB cDNA was isolated following *Xho* I digestion of pHexB43 (ATCC, Manassas VA) and cloned into the *Xho* I site of pIRES (Clontech Laboratories, Palo Alto CA) downstream of the vector's cytomegalovirus (CMV) promoter sequence. The HexA cDNA was isolated from pBHA-5 (ATCC, Manassas VA) by *Xho* I digestion and was subsequently inserted into the *Xba* I site of pIRES(HexB) downstream of the vector's IRES cassette by blunt ligation. A *IRES-lacZ* cassette was obtained from Dr. Howard J. Federoff, University of Rochester School of Medicine and Dentistry, but can be produced using standard recombinant techniques with known reagents and was inserted downstream to HexA into the *Sal* I site of pHexB-IRES-HexA by blunt ligation. In this construct, the cytomegalovirus promoter (CMV) drives transgene expression, and the translation of the second and third open reading frames (ORF), *HexB* and *lacZ*, respectively, are facilitated by an internal ribosomal entry sequence (IRES). The FIV(Hex) vector was constructed by isolating the HexB-IRES-HexA ( $\beta$ -Hex) fragment of pHexlacZ with *Nhe*I - *Not*I digestion is present and it was cloned into the FIV backbone (Poeschla et al., 1998), derived after excising the *lacZ* cassette from pFIV(*lacZ*) with *Bpu*1102I, leading to the successful construction of pFIV(Hex) (See Figures 3 and 4). Restriction fragment analysis indicated that pFIV(Hex) was constructed as designed. (Figure 5).

[191] The viral derived IRES sequence can effectively drive the expression of second genes in bicistronic constructs *in vitro* and *in vivo*, (Gurtu et al., 1996; Geschwind et al., 1996; Havenga et al. 1998). Nevertheless, IRES-mediated transcription in bicistronic constructs has been shown to reduce the levels of expression of the second ORF by about 40-50%. Hence, since HexB is necessary in the synthesis of both HEXA ( $\alpha/\beta$ ) and HEXB

( $\alpha/\alpha$ ), it was cloned first in our tricistronic construct. Confirmation of the construct has been achieved by multiple restriction enzyme digestions as well as direct DNA sequencing.

#### b) Results

[192] The FIV backbone vector was isolated from the FIV(lacZ) vector following  
5 *Sst* II & *Not* I digestion. The bicistronic transgene *HexB*-IRES-*HexA* was extracted from the  
pHexlacZ vector following *Nhe* I & *Not* I digestion, and was cloned into the FIV backbone  
by blunt ligation. FIV(Hex) digestion with the restriction enzymes *Xho* I and *Sal* I  
confirmed the cloning. (Figure 6) FIV(Hex) virus was prepared using established methods  
and was tested *in vitro* as follows. Cultured murine fibroblasts (CrFK cell line) were exposed  
10 to FIV(Hex) for 12 hours, followed fresh media change. After 48 hours, cellular DNA and  
RNA extracts were collected. The presence of viral DNA was assessed by PCR with primers  
sets specifically designed for HexB (Figure 6A). HexB expression was assessed by RT-PCR  
(Figure 6B). These results demonstrate the ability of FIV(Hex) to transduce mouse  
fibroblasts with  $\beta$ -Hex, resulting in transgene mRNA expression. (Figure 6).

15 [193] The tricistronic vector pHEXlacZ was stably expressed in embryonic  
hamster kidney fibroblasts (BHK-21; ATCC) following standard transfection laboratory  
techniques using the LIPOFECTAMINE® reagent (Gibco BRL) per manufacturer's  
instructions. Forty-eight hours post-transfection, the cells were treated with 800 $\mu$ g/mL  
G418 (Gibco BRL) for 10 days, and cell lines were selected, expanded and analyzed for  
20 expression of our tricistronic gene as follows. Analysis of the transfected cells showed that  
cell lines (Crfk, spleen, brain, liver, and kidney) stained positively for X-gal, indicating  
expression of and translation of the expressed product from the tricistronic vector. (Figure  
6)

#### 3. Example 4 In vivo use of FIV HEX vectors

25 [194] FIV(Hex) was constructed by inserting the bicistronic gene HexB-IRES-  
HexA in the place of the reporter gene lacZ in the FIV backbone vector using standard  
molecular biology techniques. FIV(Hex) was prepared *in vitro* by transient co-transfection  
of the transfer vector along with the packaging and envelop plasmids into 293H cells. The  
virus-rich supernatant was centrifuged and the viral pellet was reconstituted in normal  
30 saline, and was then titered in CrFK cells by the X-Hex histochemical method ( $10^7$ - $10^8$   
infectious particles/ml). The viral solution was injected intraperitoneally to 2 days old  
HexB<sup>-/-</sup> knockout mouse pups, which were allowed to reach the critical age of 16 weeks,

when they displayed full signs of the lysosomal storage disease. For control, littermates were injected with the FIV(lacZ) virus, which is identical to FIV(Hex), but instead of carrying the HexB-IRES-HexA gene it carries the reporter gene lacZ. Locomotive performance was evaluated by placing the mice on a wire mesh attached on a clear plexiglass cylinder, and turning the wire mesh up-side-down. The lapse time until the mice fell off the wire mesh was recorded on weekly basis until the mice were terminated. It is important to state that at the critical time point of 16 weeks, the FIV(Hex) injected mice showed statistically better locomotive performance compared to FIV(lacZ) injected mice (controls). Furthermore, the FIV(Hex) mice had an extended life span for at least 2-3 additional weeks, at which point they were also terminated because they were showing signs of the disease.

#### 4. Example 3 HIV HEX vectors

[195] The HexB-IRES-HexA therapeutic gene was cloned into the Lenti6/V5D-TOPO vector commercially available by Invitrogen (Carlsbad, CA), whereby the cytomegalovirus promoter CMV drives gene expression [in a manner similar to FIV(Hex)]. A virus was constructed whereby the expression of HexB-IRES-HexA is driven by a promoter, such as that show in SEQ ID NO:23, which consists of a beta-actin portion and a CMV portion. This type of promoter has high expression in mammalian cells.

#### G. References

[196] Akli S, Guidotti J-E, Vigne E, Perricaudet M, Sandhoff K, Kahn A, Poenaru L (1996) Restoration of hexosaminidase A activity in human Tay-Sachs fibroblasts via adenoviral vector mediated gene transfer. *Gene Therapy* 3: 769-774

[197] Bowers WJ, Howard DF, Federoff HJ (2000). Discordance between expression and genome transfer titering of HSV amplicon vectors: recommendation for standardized enumeration. *Mol Ther* 1:294-9.

[198] Burns JC, Friedmann T, Driever W, Burrascano M, Yee J-K (1993) Vesicular stomatitis virus G-glycoprotein pseudotyped retroviral vectors Concentration to very high titer and efficient gene transfer into mammalian and nonmammalian cells. *Proc Natl Acad Scie USA* 90: 8033-8037.

[199] Cannon PM, Anderson WF (2000) Retroviral vectors for gene therapy. *In* *Gene Therapy: therapeutic mechanisms and strategies*. NS Templeton and DD Lasic, Editors. Marcel Dekker Inc, New York, pp 1-16.



- [200] Daly TM, Okuyama T, Vogler C, Haskins ME, Muzyczka N, Sands MS (1999a) Intracranial injection of recombinant adeno-associated virus improves cognitive function in a murine model of mucopolysaccharidosis type VII. *Hum Gene Ther* 10:85-94.
- [201] Daly TM, Vogler C, Levy B, Haskins ME, Sands MS (1999b) Neonatal gene transfer leads to widespread correction of pathology in a murine model of lysosomal disease. *Proc Natl Acad Sci USA* 96: 2296-2300.
- [202] Enlow DH, Hans MG (1996) *Essentials of facial growth*. W.B.Saunders Co, New York.
- [203] Goldstein GW, Betz AL, Bowman PD, Dorovini-Zis K (1986) In vitro studies of the blood-brain barrier using isolated brain capillaries and cultured endothelial cells. *Ann NY Acad Sci* 481:202-13.
- [204] Gorlin RJ, Cohen MM, Levin LS (1990). *SYNDROMES OF THE HEAD AND NECK*, 3<sup>rd</sup> Edition. Oxford university Press, New York.
- [205] Guidotti JE, M. A., Haase G, Caillaud C, McDonell N, Kahn A, Poenaru L (1999) Adenoviral gene therapy of the Tay-Sachs disease in hexosaminidase A deficient knockout mice. *Hum Mol Genet* 8: 831-838.
- [206] Kjaer I, Keeling JW, Hansen BF (1999) The prenatal human cranium-Normal and pathologic development. Munksgaard, Copenhagen, pp 20-51.
- [207] Kordower JH, Bloch J, Ma SY, Chu Y, Palfi S, Roitberg BZ, Emborg M, Hantraye P, Deglon N, Aebischer P (1999) Lentiviral gene transfer to the nonhuman primate brain. *Exp Neurol* 160: 1-16.
- [208] Kyrkanides S, Kjaer I, Fisher-Hansen B (1993) The basilar part of the occipital bone in normal and pathological fetuses. *J Cranio Genet Develop Biol* 13: 184-192.
- [209] Kyrkanides S, Bellohusen R, Subtelny JD (1995) Skeletal asymmetries of the nasomaxillary complex in non-cleft and post-surgical unilateral cleft lip and palate individuals. *Cleft Pal Craniofac J* 32: 428-32.
- [210] Kyrkanides S, Bellohusen R, Subtelny JD (1996) Asymmetries of the upper lip and nose in non-cleft and post-surgical unilateral cleft lip and palate individuals. *Cleft Palate-Craniofacial Journal*, 33:306-10.

- [211] Kyrkanides S, Olschowka JA, Williams JP, Hansen JT, O'Banion MK (1999). TNF $\alpha$  and IL-1 $\beta$  mediate ICAM-1 induction via microglia-astrocyte interaction in CNS radiation injury. *J Neuroimmunol* 95:95-106.
- [212] Kyrkanides S, Klambani M, Subtelny JD (2000). Cranial base and facial skeleton asymmetries in unilateral cleft lip & palate individuals. *Cleft Palate-Craniofacial Journal* 37: 556-561.
- [213] Kyrkanides S, Olschowka JA, Whitley P, O'Banion MK (2001). Enhanced glial activation and expression of specific CNS inflammation-related molecules in aged versus young rats following cortical stab injury. *In Press*, *J Neuroimmunol*.
- 10 [214] Lipschutz GS, Gruber CA, Cao Y, Hardy J, Contag CH, Gaensler KML (2001). *In utero* delivery of adeno-associated viral vectors: Intraperitoneal gene transfer produces long term expression. *Mol Ther* 3: 284-92.
- [215] McCormack JE, Edwards W, Sensintaffer J, Lillegren L, Kozloski M, Brumm D, Karavodin L, Jolly DJ, Greengard J (2001) *Mol Ther* 3: 516-525.
- 15 [216] Phaneuf D, Wakamatsu N, Huang J-Q, Borowski A, Peterson AC, Fortunato SR, Ritter G, Igoudra SA, Morales CR, Benoit G, Akerman BR, Leclerc D, Hanai N, Marth JD, Trasler JM, Gravel RA (1996) Dramatically different phenotypes in mouse models of human Tay-Sachs and Sandhoff diseases. *Hum Mol Genet* 5: 1-14.
- [217] Poeschla EM, Wong-Stall F, Looney DL (1998) Efficient transduction of  
20 nondividing human cells by feline immunodeficiency virus lentiviral vectors. *Nature Medicine* 4: 354-357.
- [218] Proia RL, d'Azzo A, Neufeld EF (1984). Association of a- and b- subunits during the biosynthesis of b-hexosaminidase in cultured human fibroblasts. *J Biol Chem* 259: 3350-3354.
- [219] Risau W, Hallmann R, Albrecht U (1986) Differentiation-dependent expression of  
25 proteins in brain endothelium during development of the blood-brain barrier. *Devel Biol* 117: 537-545.
- [220] Risau W, Wolburg H (1990) Development of blood-brain barrier. *TINS* 13:174-178.
- [221] Rosenberg A (1999) Bones, joints and soft tissue tumors. *In* Robbins Pathologic Basis of Disease, 6<sup>th</sup> edition; Cotran, Kumar & Collins (Ed), W.B.Saunders Co, New York.
- 30 [222] Sakurada H, Itoh K, Kuroki Y, Kase R, Shimmoto M, Utsumi K, Ozawa H, Tai T, Hara A, Uyama E (1997) Immunocytochemical detection of accumulated substrates

in cultured fibroblasts from patients with the infantile and adult forms of Sandhoff disease.  
Clin Chim Acta 265: 263-266.

[223] Sango K, McDonald MP, Crawley JN, Mack ML, Tift CJ, Skop E, Starr CM, Hoffmann A, Sandhoff K, Suzuki K, Proia RL (1996) Mice lacking both subunits of lysosomal  $\beta$ -hexosaminidase display gangliosidosis and mucopolysaccharidosis. Nature Genet 14: 348-352.

[224] Sango K, Yamanaka S, Hoffman A, Okuda Y, Grinberg A, Westphal H, McDonald MP, Crawley JN, Sandhoff K, Suzuki K, Proia RL (1995) Mouse models of Tay-Sachs and Sandhoff diseases differ in neurologic phenotype and ganglioside metabolism. Nature Genet 11: 170-176.

10 [225] Schroeder WH, Mortari F, Shiokawa S, Kirkham PM, Elgavish RA, Bertrand FE III (1995) Developmental regulation of the human antibody repertoire. Ann NY Acad Scie 764: 242-260.

[226] Suzuki K, Sango K, Proia RL, Langman C (1997) Mice deficient in all forms of lysosomal  $\beta$ -hexosaminidase show mucopolysaccharosis-like pathology. J Neuropath Exp Neurol 15 56: 693-703.

[227] Tarantal AF, O'Rourke JP, Case SS, Newbound GC, Li J, Lee CI, Baskin CR, Kohn DB, Bunnell BA (2001) Rhesus monkey model for fetal gene transfer: studies with retroviral-based vector systems. Mol Ther 3: 128-138.

[228] Wolff JA, Harding CO (2000) Principles of gene therapy for inborn errors of metabolism. In Gene Therapy: therapeutic mechanisms and strategies. NS Templeton and DD Lasic, Editors. Marcel Dekker Inc, New York, pp 507-533.

## H. Sequences

1. SEQ ID NO:1 Homo sapiens hexosaminidase A (alpha polypeptide) (HEXA),  
Genbank Accession No. XM\_037778
2. SEQ ID NO:2 Homo sapiens hexosaminidase A (alpha polypeptide) (HEXA),  
Genbank Accession No. XM\_037778
3. SEQ ID NO:3 Homo sapiens hexosaminidase B (beta polypeptide) (HEXB),  
protein Genbank Accession No XM\_032554
4. SEQ ID NO:4 Homo sapiens hexosaminidase B (beta polypeptide) (HEXB),  
mRNA Genbank Accession No XM\_032554
5. SEQ ID NO:5 IRES sequence United States Patent No. 4,937,190 herein  
incorporated by reference covers entire Vector, and is cited at least for material  
relating to the pIRES vector)
6. SEQ ID NO:6 Mus musculus hexosaminidase A (Hexa), protein Genbank  
Accession No, NM\_010421
7. SEQ ID NO:7 Mus musculus hexosaminidase A (Hexa), mRNA Genbank  
Accession No, NM\_010421
8. SEQ ID NO:8 FIV(LacZ) construct 12750 bp
9. SEQ ID NO:9: HEX- $\alpha$  polypeptide Genbank accession number NM\_000520  
(Proia) beta-hexosaminidase A alpha-subunit to human chromosomal region  
15q23—q24
10. SEQ ID NO:10 HexA gene Genbank accession number NM\_000520 (Proia)
11. SEQ ID NO:11 HexB degenerate cDNA G to A change at position 6
12. SEQ ID NO:12: HEX- $\beta$  polypeptide conservative substitution of Val21 to I21
13. SEQ ID NO:13 HEX- $\alpha$  polypeptide Genbank accession number M16411  
(Tissue sample from ATCC)
14. SEQ ID NO:14 HexA gene Genbank accession number M16411
15. SEQ ID NO:15: HEX- $\beta$  polypeptide Genbank accession number NM\_000521  
(Proia) beta-hexosaminidase A alpha-subunit to human chromosomal region  
chromosome 5 map="5q13"
16. SEQ ID NO:16 HexB gene Genbank accession number NM\_000521 Proia
17. SEQ ID NO:17 Mus musculus hexosaminidase B (Hexb), protein. Genbank  
Accession No. NM\_010422

18. SEQ ID NO:18 *Mus musculus* hexosaminidase B (Hexb), mRNA. Genbank  
Accession No. NM\_010422
19. SEQ ID NO:19 Bactin Hex sequence
20. SEQ ID NO:20 HIV Hex vector sequence
- 5 21. SEQ ID NO:21 E02199 DNA encoding chicken beta actin gene promoter.
22. SEQ ID NO:22 Chicken Beta Actin promoter
23. SEQ ID NO:23 CMV-Beta actin promoter
24. SEQ ID NO:24 Fusion promoter-CMV portion
25. SEQ ID NO:25 Fusion promoter – beta actin portion
- 10 26. SEQ ID NO:26 Chicken beta actin promoter
27. SEQ ID NO:27 Accession # BD136067. promoter element for sustained gene  
expression from CMV promoter.
28. SEQ ID NO:28 BD136066 Accession # promoter element for sustained gene  
expression from CMV promoter.
- 15 29. SEQ ID NO:29 BD136065 Accession # promoter element for sustained gene  
expression from CMV promoter.
30. SEQ ID NO:30 BD136064 Accession # promoter element for sustained gene  
expression from CMV promoter.
- 20 31. SEQ ID NO:31 L77202 Accession # Murine Cytomegalovirus early (E1) gene,  
promoter region.
32. SEQ ID NO:32 X03922 Accession # Human cytomegalovirus (HCMV) IE1  
gene promoter region.
33. SEQ ID NO:33 E06566 Accession # Promoter gene of human beta-actin gene.
- 25 34. SEQ ID NO:34 E02198 Accession # Dna encoding 3'end region of beta-actin  
gene promoter
35. SEQ ID NO:35 E02197 Accession # DNA encoding 3'end region of beta-actin  
gene promoter.
36. SEQ ID NO:36 E02196 Accession # DNA encoding 3'end region of beta-actin  
gene promoter.
- 30 37. SEQ ID NO:37 E02195 Accession # DNA encoding 3'end region of beta-actin  
gene promoter.

38. EQ ID NO:38 E02194 Accession # DNA encoding chicken beta-actin gene promoter.
39. SEQ ID NO:39 E01452 Accession # Genomic DNA of promoter of human beta-actin.
- 5 40. SEQ ID NO E03011 Accession # DNA encoding hybrid promoter that is composed of chicken beta-actin gene promoter and rabbit beta-globin gene promoter.
41. SEQ ID NO:41 BD015377 Accession # Baculovirus containing minimum CMV promoter.
- 10 42. Other cytomegalovirus promoter regions
- [229] Other human cytomegalovirus promoter regions can be found in accession numbers M64940, Human cytomegalovirus IE-1 promoter region, M64944 Human cytomegalovirus IE-1 promoter region, M64943 Human cytomegalovirus IE-1 promoter region, M64942 Human cytomegalovirus IE-1 promoter region, M64941 Human
- 15 cytomegalovirus IE-1 promoter region (All of which are herein incorporated by reference at least for their sequence and information)

## VI. CLAIMS

What is claimed is:

1. A composition comprising a nucleic acid wherein the nucleic acid comprises a sequence encoding a HEX- $\alpha$  and a sequence encoding a HEX- $\beta$ .
2. The composition of claim 1, wherein the sequence encoding the HEX- $\beta$  is orientated 5' to the sequence encoding HEX- $\alpha$ .
3. The composition of claim 1, further comprising a promoter.
4. The composition of claim 1, further comprising an integrated ribosomal entry site (IRES).
5. The composition of claim 4, wherein the sequence encoding the HEX- $\beta$  is orientated 5' to the IRES sequence and the IRES sequence is located 5' to the sequence encoding HEX- $\alpha$ .
6. The composition of claim 4, further comprising a promoter.
7. The composition of claim 6, wherein the promoter is located 5' to the sequence encoding the HEX- $\beta$  and the sequence encoding the HEX- $\beta$  is orientated 5' to the IRES sequence and the IRES sequence is located 5' to the sequence encoding HEX- $\alpha$ .
8. The composition of claim 6, wherein the parts are oriented 5'-promoter- HEX- $\beta$  encoding sequence-IRES- HEX- $\alpha$  encoding sequence-3'.
9. The composition of claim 6, wherein the parts are oriented 5'-promoter- HEX- $\alpha$  encoding sequence -IRES- HEX- $\beta$  encoding sequence -3'.
10. The composition of claim 6, wherein the nucleic acid comprises a second IRES sequence.
11. The composition of claim 10, wherein the second IRES sequence is located 3' to the other parts.
12. The composition of claim 6, wherein the HEX- $\beta$  has at least 70%, 75%, 80%, 85%, 90%, or 95% identity to the sequence set forth in SEQ ID NO:3 and the HEX- $\alpha$  has at least 70%, 75%, 80%, 85%, 90%, or 95% identity to the sequence set forth in SEQ ID NO:1.
13. The composition of claim 12, wherein any change from SEQ ID NO:3 or SEQ

ID NO:1 is a conservative change.

14. The composition of claim 13 wherein the HEX- $\beta$  has the sequence set forth in SEQ ID NO:3 and the HEX- $\alpha$  has the sequence set forth in SEQ ID NO:1.

15. The composition of claim 6, wherein the sequence encoding HEX- $\beta$  hybridizes to SEQ ID NO:2 under stringent conditions and wherein the HEX- $\alpha$  element hybridizes to SEQ ID NO:4 under stringent conditions.

16. The composition of claim 12, wherein the IRES sequence comprises a sequence having at least 70%, 75%, 80%, 85%, 90%, or 95% identity to the sequence set forth in SEQ ID NO:5.

17. The composition of claim 16, wherein the promoter sequence comprises a constitutive promoter.

18. The composition of claim 17, wherein the promoter sequence comprises a CMV promoter.

19. The composition of claim 18, wherein the CMV promoter comprises the sequence set forth in SEQ ID NO:32.

20. The composition of claim 16, wherein the promoter sequence comprises a beta actin promoter.

21. The composition of claim 20, wherein the beta actin promoter sequence comprises an avian beta actin promoter sequence.

22. The composition of claim 21, wherein the beta actin promoter sequence comprises a mammalian beta actin promoter sequence.

23. The composition of claim 21, wherein the beta actin promoter comprises the sequence set forth in SEQ ID NO:26.

24. The composition of claim 16, wherein the promoter sequence comprises an inducible promoter.

25. The composition of claim 18, wherein the promoter sequence further comprises a beta actin promoter.

26. The composition of claim 6, wherein the composition produces a functional HEXB product.



27. The composition of claim 6, wherein the composition produces a functional HEXA product.
28. The composition of claim 6, wherein the composition produces a functional HEXS product.
29. The composition of claim 26, wherein the composition is capable of cross correcting.
30. The composition of claim 26, wherein the function is the catabolism of GM2 gangliosides in mammalian cells. Same for HEXB, the homodimer of HexB/HexB.
31. The composition of claim 6, wherein the nucleic acid further comprises a reporter gene.
32. The composition of claim 31, wherein the reporter gene is a lacZ gene.
33. The composition of claim 31, wherein the reporter gene is flanked by recombinase sites.
34. The composition of claim 33, wherein the recombinase sites are for the cre recombinase.
35. The composition of claim 6, wherein the nucleic acid further comprises a transcription termination site.
36. The composition of claim 35, wherein the transcription termination site is oriented 5' to the promoter sequence.
37. The composition of claim 36, wherein the transcription termination site is flanked by recombinase sites.
38. The composition of claim 37, wherein the recombinase sites are for the cre recombinase.
39. The composition of claim 6, further comprising a vector.
40. The composition of claim 39, wherein the vector comprises a lentiviral vector.
41. The composition of claim 40, wherein the lentiviral vector comprises a feline immunodeficiency virus.
42. The composition of claim 40, wherein the lentiviral vector comprises a human immunodeficiency virus.

43. The composition of claim 39, wherein the vector can be stably integrated for at least three months.

44. A composition comprising a cell wherein the cell comprises the nucleic acid of claim 6.

45. A composition comprising a cell wherein the cell comprises the vector of claim 39.

46. The composition of claim 47, wherein the cell comprises a neuron, glia cell, fibroblast, chondrocyte, osteocyte, endothelial cell, or hepatocyte.

47. The composition of claims 6, wherein the composition is in pharmaceutically acceptable form.

48. The composition of claims 6, wherein the composition is in an effective dosage.

49. The composition of claim 48, wherein the effective dosage is determined as a dosage that reduces the effects of Tay Sachs or Sandoff's disease.

50. A composition comprising an animal wherein the animal comprises the vector of claim 39.

51. A composition comprising an animal wherein the animal comprises the nucleic acid of claim 6.

52. A composition comprising an animal wherein the animal comprises the cell of claim 45.

53. The composition of claim 50, wherein the animal is mammal.

54. The composition of claim 53, wherein the mammal is a murine, ungulate, or non-human primate.

55. The method of claim 54, wherein the mammal is a mouse, rat, rabbit, cow, sheep, or pig.

56. The composition of claim 54, wherein the mammal is mouse.

57. The composition of claim 56, wherein the mouse comprises a HexB knockout.

58. The composition of claim 56, wherein the mouse comprises a HexA knockout.

59. The composition of claim 58, wherein the mouse further comprises a HexB

knockout.

60. The composition of claim 54, wherein the mammal is a non-human primate.

61. A method of providing HEXA in a cell comprising transfecting the cell with the nucleic acids of claims 6.

62. A method of providing HEXB in a cell comprising transfecting the cell with the nucleic acids of claims 6.

63. A method of providing HEX- $\alpha$  and HEX- $\beta$  in a cell comprising transfecting the cell with the nucleic acid of claims 6.

64. The method of claim 63, wherein the step of transfecting occurs in vitro.

65. The method of claim 63, wherein the step of transfecting occurs in vivo.

66. A method of providing HEXS in a cell comprising transfecting the cell with the nucleic acids of claims 6.

67. A method of making a transgenic organism comprising administering the nucleic acid of claims 6.

68. A method of making a transgenic organism comprising administering the vector of claim 39.

69. A method of making a transgenic organism comprising administering the cell of claims 45.

70. A method of making a transgenic organism comprising transfecting a lentiviral vector to the organism at during a perinatal stage of the organism's development.

71. A method of treating a subject having Tay Sachs disease and/or Sandoff disease comprising administering the composition of claim 47.

72. A method of making a composition, the composition comprising a nucleic acid molecule, wherein the nucleic acid molecule is produced by the process comprising linking in an operative way a promoter element, an element comprising sequence encoding HEX- $\beta$ , a IRES element, and an element encoding HEX- $\alpha$ .

73. The method of claim 72 wherein the HEX- $\beta$  element comprises a sequence having at least 80% SEQ ID NO:1 and the HEX- $\alpha$  element comprises a sequence having at least 80% to SEQ ID NO:3.

74. The method of claim 73, wherein any change in SEQ ID NO:1 or SEQ ID NO:3 is a conservative change.

75. The method of claim 72, wherein the sequence encoding HEX- $\beta$  hybridizes to SEQ ID NO:2 under stringent conditions and wherein the sequence encoding the HEX- $\alpha$  hybridizes to SEQ ID NO:4 under stringent conditions.

76. A method of producing a composition, the composition comprising a cell, the method comprising administering the nucleic acid of claim 6 to the cell.

77. A method of producing a composition, the composition comprising a peptide, the method comprising expressing the nucleic acid of claim 6.

78. The method of claim 77, further comprising isolating the peptide.

79. A method of producing a composition, the composition comprising an animal, the method comprising administering the nucleic acid of claim 6 to the animal.

80. The method of claim 79, wherein the animal is a mammal.

81. Wherein the mammal is a murine, ungulate, or non-human primate.

82. The method of claim 81, wherein the mammal is a mouse, rat, rabbit, cow, sheep, or pig.

83. A nucleic acid comprising a sequence encoding HEX- $\beta$  wherein the HEX- $\beta$  has the sequence set forth in SEQ ID NO:3, a sequence encoding HEX- $\alpha$ , wherein the HEX- $\alpha$  has the sequence set forth in SEQ ID NO:1, a promoter, and an IRES sequence, wherein the promoter is located 5' to the sequence encoding the HEX- $\beta$  and the sequence encoding the HEX- $\beta$  is orientated 5' to the IRES sequence and the IRES sequence is located 5' to the sequence encoding HEX- $\alpha$ .

84. A composition comprising a nucleic acid wherein the nucleic acid comprises a sequence encoding a first HEX- $\beta$  and a sequence encoding a second HEX- $\beta$ .

85. A composition comprising a nucleic acid wherein the nucleic acid comprises a sequence encoding a first HEX- $\alpha$  and a sequence encoding a second HEX- $\alpha$ .

86. A composition comprising four parts: 1) a promoter, 2) a sequence encoding a HEX- $\alpha$ , 3) a sequence encoding a HEX- $\beta$ , and 4) an integrated ribosomal entry site (IRES).

1/27




FIG.1A




FIG.1F

2/27



FIG.1B




FIG.1C

3/27



FIG. 1D1




FIG. 1D2




FIG. 1E1



FIG. 1E2



FIG. 1G1







FIG. 1G2

4/27



control plain medium

FIG.2B



conditioned medium

FIG.2A



5/27




FIG.3

6/27

FIV (Hex)




FIG.4

7/27

FIV (Hex) 6.2 maxi prep      Total 1mg  
1kb<sup>+</sup> uncut Sca I Not I Sal I Xho I 1kb<sup>+</sup>      (1μg/μl)




FIG.5

8/27




FIG.6

9/27




FIG.7

10/27




FIG.8

11/27






FIG.9

12/27



**FIG. 10**



13/27




FIG.11

14/27




FIG. 11A




FIG. 11B

15/27




FIG.12

16/27




FIG.13




FIG.14

18/27




FIG.15

Hex<sup>XAT</sup>: Excisionally activated  $\beta$ -hexosaminidase gene




FIG.16A




FIG.16B




FIG.16C

20/27


Hex<sup>XAT</sup>: EXCISIONALLY ACTIVATED  $\beta$ -HEXOSAMINIDASE GENE

FIG.17A


Hex<sup>XAT</sup>: EXCISIONALLY ACTIVATED  $\beta$ -HEXOSAMINIDASE GENE

FIG.17B



21/27




FIG.17C

22/27




FIG.18

23/27

GM<sub>2</sub> ganglioside immunocytochemistry




FIG.19

24/27




FIG. 20

25/27

PERINATAL GENE THERAPY IN  $\text{hexB}^{-/-}$  MICE




FIG.21

26/27




FIG.22

27/27




FIG.23

Met 1	Met	Thr	Ser	Val 5	Tyr	Ser	Ser	Leu	Arg 10	Leu	Ser	Gly	Glu	Leu 15	Ser
Glu	Val	Trp	Arg 20	Leu	Leu	Ala	Ser	Leu 25	Phe	Gly	Asn	Leu 30	Leu	Arg	Ala
Gln	Phe	Phe 35	Ile	Asn	Lys	Thr	Glu 40	Ile	Glu	Asp	Phe	Pro 45	Arg	Phe	Pro
His	Arg 50	Gly	Leu	Leu	Leu	Asp 55	Thr	Ser	Arg	His	Tyr 60	Leu	Pro	Leu	Ser
Ser 65	Ile	Leu	Asp	Thr	Leu 70	Asp	Val	Met	Ala	Tyr 75	Asn	Lys	Leu	Asn 80	Val
Phe	His	Trp	His 85	Leu	Val	Asp	Asp	Pro 90	Ser	Phe	Pro	Tyr	Glu 95	Ser	Phe
Thr	Phe	Pro	Glu 100	Leu	Met	Arg	Lys	Gly 105	Ser	Tyr	Asn	Pro 110	Val	Thr	His
Ile	Tyr	Thr 115	Ala	Gln	Asp	Val	Lys 120	Glu	Val	Ile	Glu	Tyr 125	Ala	Arg	Leu
Arg	Gly 130	Ile	Arg	Val	Leu	Ala 135	Glu	Phe	Asp	Thr	Pro 140	Gly	His	Thr	Leu
Ser 145	Trp	Gly	Pro	Gly 150	Ile	Pro	Gly	Leu	Leu	Thr 155	Pro	Cys	Tyr	Ser	Gly 160
Ser	Glu	Pro	Ser 165	Gly	Thr	Phe	Gly	Pro	Val 170	Asn	Pro	Ser	Leu 175	Asn	Asn
Thr	Tyr	Glu 180	Phe	Met	Ser	Thr	Phe 185	Phe	Leu	Glu	Val	Ser 190	Ser	Val	Phe
Pro	Asp	Phe 195	Tyr	Leu	His	Leu	Gly 200	Gly	Asp	Glu	Val	Asp 205	Phe	Thr	Cys
Trp	Lys 210	Ser	Asn	Pro	Glu	Ile 215	Gln	Asp	Phe	Met	Arg 220	Lys	Lys	Gly	Phe
Gly 225	Glu	Asp	Phe	Lys 230	Gln	Leu	Glu	Ser	Phe	Tyr 235	Ile	Gln	Thr	Leu 240	Leu



```

Asp Ile Val Ser Ser Tyr Gly Lys Gly Tyr Val Val Trp Gln Glu Val
                245                250                255
Phe Asp Asn Lys Val Lys Ile Gln Pro Asp Thr Ile Ile Gln Val Trp
                260                265                270
Arg Glu Asp Ile Pro Val Asn Tyr Met Lys Glu Leu Glu Leu Val Thr
                275                280                285
Lys Ala Gly Phe Arg Ala Leu Leu Ser Ala Pro Trp Tyr Leu Asn Arg
                290                295                300
Ile Ser Tyr Gly Pro Asp Trp Lys Asp Phe Tyr Ile Val Glu Pro Leu
305                310                315                320
Ala Phe Glu Gly Thr Pro Glu Gln Lys Ala Leu Val Ile Gly Gly Glu
                325                330                335
Ala Cys Met Trp Gly Glu Tyr Val Asp Asn Thr Asn Leu Val Pro Arg
                340                345                350
Leu Trp Pro Arg Ala Gly Ala Val Ala Glu Arg Leu Trp Ser Asn Lys
                355                360                365
Leu Thr Ser Asp Leu Thr Phe Ala Tyr Glu Arg Leu Ser His Phe Arg
370                375                380
Cys Glu Leu Leu Arg Arg Gly Val Gln Ala Gln Pro Leu Asn Val Gly
385                390                395                400
Phe Cys Glu Gln Glu Phe Glu Gln Thr
                405

```

&lt;210&gt; 2

&lt;211&gt; 2256

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 2

```

cctccgagag gggagaccag cgggccatga caagctccag gctttgggtt tcgctgctgc      60
tggcggcagc gttcgcagga cgggcgacgg ccctctggcc ctggcctcag aacttccaaa      120
cctccgacca gcgctaagtc ctttaccgga acaactttca attccagtac gatgtcagct      180
cggccgcgca gcccggtctg tcagtcctcg acgaggcctt ccagcgctat cgtgacctgc      240
ttttcggttc cgggtcttgg ccccgctcct acctcacagg gaaacggcat aactgggaga      300
agaatgtgtt ggttgtctct gtagtcacac ctggatgtaa ccagcttctt actttggagt      360
cagtggagaa ttataccctg accataaatg atgaccagtg ttactcctc tctgagactg      420
tctggggagc tctccgaggt ctggagactt ttagccagct tgtttggaaa tctgctgagg      480
gcacagttct ttatcaacaa gactgagatt gaggactttc cccgctttcc tcaccggggc      540
ttgctgttgg atacatctcg ccattacctg ccactctcta gcatcctgga cactctggat      600
gtcatggcgt acaataaatt gaacgtgttc cactggcatc tggtagatga tccttccttc      660
ccatatgaga gcttcacttt tccagagctc atgagaaagg ggtcctacaa ccctgtcacc      720
cacatctaca cagcacagga tgtgaaggag gtcattgaat acgcacggct ccgggggtatc      780
cgtgtgcttg cagagtttga cactcctggc cacactttgt cctggggacc aggtatccct      840
ggattactga ctcccttgct ctctgggtct gagccctctg gcacctttgg accagtgaat      900
cccagttctc ataataacct tgagttcatg agcacattct tcttagaagt cagctctgtc      960
ttcccagatt tttatcttca tcttgaggga gatgaggttg atttcacctg ctggaagtcc     1020
aaccagaga tccaggactt tatgaggaag aaaggcttcg gtgaggactt caagcagctg     1080
gagtccttct acatccagac gctgctggac atcgtctctt cttatggcaa gggctatgtg     1140
gtgtggcagg aggtgtttga taataaagta aagattcagc cagacacaaat catacaggtg     1200
tggcgagagg atattccagt gaactatatg aaggagctgg aactggtcac caaggccggc     1260
ttccgggccc ttctctctgc cccctgttac ctgaaccgta tatcctatgg ccctgactgg     1320
aaggatttct acatagtgga acccctggca tttgaaggta cccctgagca gaaggctctg     1380
gtgattggtg gagaggcttg tatgtgggga gaatatgtgg acaacacaaa cctggtcccc     1440
aggctctggc ccagagcagg ggctgttgcc gaaaggctgt ggagcaacaa gttgacatct     1500
gacctgacat ttgcctatga acgtttgtca cacttccgct gtgaattgct gaggcgaggt     1560

```

```

gtccaggccc aaccctcaa tgtaggcttc tgtgagcagg agtttgaaca gacctgagcc 1620
ccaggcaccg aggagggtgc tggctgtagg tgaatggtag tggagccagg cttccactgc 1680
atcctggcca ggggacggag ccccttgccct tcgtgccctc tgcctgcgtg cccctgtgct 1740
tggagagaaa ggggcccgtg ctggcgctcg cattcaataa agagtaatgt ggcatttttc 1800
tataataaac atggattacc tgtgtttaaa aaaaaaagtg tgaatggcgt tagggtaagg 1860
gcacagccag gctggagtca gtgtctgccc ctgaggtctt ttaagttgag ggctgggaat 1920
gaaacctata gcctttgtgc tgttctgcct tgccgtgag ctatgtcact cccctccac 1980
tcctgacatc attccagaca cctgccctaa tcctcagcct gctcacttca cttctgcatt 2040
atatctccaa ggcgttggtg tatggaaaaa gatgtagggg cttggagggtg ttctggacag 2100
tggggagggc tccagaccca acctggtcac agaagagcct ctcccccattg catactcatc 2160
cactccctc ccctagagct attctcctt gggtttcttg ctgcttcaat tttatacaac 2220
cattatttaa atattattaa acacatattg ttctct 2256

```

&lt;210&gt; 3

&lt;211&gt; 544

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 3

```

Met Leu Leu Ala Leu Leu Leu Ala Thr Leu Leu Ala Ala Met Leu Ala
 1          5          10          15
Leu Leu Thr Gln Val Ala Leu Val Val Gln Val Ala Glu Ala Ala Arg
 20          25          30
Ala Pro Ser Val Ser Ala Lys Pro Gly Pro Ala Leu Trp Pro Leu Pro
 35          40          45
Leu Leu Val Lys Met Thr Pro Asn Leu Leu His Leu Ala Pro Glu Asn
 50          55          60
Phe Tyr Ile Ser His Ser Pro Asn Ser Thr Ala Gly Pro Ser Cys Thr
 65          70          75          80
Leu Leu Glu Glu Ala Phe Arg Arg Tyr His Gly Tyr Ile Phe Gly Phe
 85          90          95
Tyr Lys Trp His His Glu Pro Ala Glu Phe Gln Ala Lys Thr Gln Val
100          105          110
Gln Gln Leu Leu Val Ser Ile Thr Leu Gln Ser Glu Cys Asp Ala Phe
115          120          125
Pro Asn Ile Ser Ser Asp Glu Ser Tyr Thr Leu Leu Val Lys Glu Pro
130          135          140
Val Ala Val Leu Lys Ala Asn Arg Val Trp Gly Ala Leu Arg Gly Leu
145          150          155          160
Glu Thr Phe Ser Gln Leu Val Tyr Gln Asp Ser Tyr Gly Thr Phe Thr
165          170          175
Ile Asn Glu Ser Thr Ile Ile Asp Ser Pro Arg Phe Ser His Arg Gly
180          185          190
Ile Leu Ile Asp Thr Ser Arg His Tyr Leu Pro Val Lys Ile Ile Leu
195          200          205
Lys Thr Leu Asp Ala Met Ala Phe Asn Lys Phe Asn Val Leu His Trp
210          215          220
His Ile Val Asp Asp Gln Ser Phe Pro Tyr Gln Ser Ile Thr Phe Pro
225          230          235          240
Glu Leu Ser Asn Lys Gly Ser Tyr Ser Leu Ser His Val Tyr Thr Pro
245          250          255
Asn Asp Val Arg Met Val Ile Glu Tyr Ala Arg Leu Arg Gly Ile Arg
260          265          270
Val Leu Pro Glu Phe Asp Thr Pro Gly His Thr Leu Ser Trp Gly Lys
275          280          285

```

Gly Gln Lys Asp Leu Leu Thr Pro Cys Tyr Ser Arg Gln Asn Lys Leu  
 290 295 300  
 Asp Ser Phe Gly Pro Ile Asn Pro Thr Leu Asn Thr Thr Tyr Ser Phe  
 305 310 315 320  
 Leu Thr Thr Phe Phe Lys Glu Ile Ser Glu Val Phe Pro Asp Gln Phe  
 325 330 335  
 Ile His Leu Gly Gly Asp Glu Val Glu Phe Lys Cys Trp Glu Ser Asn  
 340 345 350  
 Pro Lys Ile Gln Asp Phe Met Arg Gln Lys Gly Phe Gly Thr Asp Phe  
 355 360 365  
 Lys Lys Leu Glu Ser Phe Tyr Ile Gln Lys Val Leu Asp Ile Ile Ala  
 370 375 380  
 Thr Ile Asn Lys Gly Ser Ile Val Trp Gln Glu Val Phe Asp Asp Lys  
 385 390 395 400  
 Ala Lys Leu Ala Pro Gly Thr Ile Val Glu Val Trp Lys Asp Ser Ala  
 405 410 415  
 Tyr Pro Glu Glu Leu Ser Arg Val Thr Ala Ser Gly Phe Pro Val Ile  
 420 425 430  
 Leu Ser Ala Pro Trp Tyr Leu Asp Leu Ile Ser Tyr Gly Gln Asp Trp  
 435 440 445  
 Arg Lys Tyr Tyr Lys Val Glu Pro Leu Asp Phe Gly Gly Thr Gln Lys  
 450 455 460  
 Gln Lys Gln Leu Phe Ile Gly Gly Glu Ala Cys Leu Trp Gly Glu Tyr  
 465 470 475 480  
 Val Asp Ala Thr Asn Leu Thr Pro Arg Leu Trp Pro Arg Ala Ser Ala  
 485 490 495  
 Val Gly Glu Arg Leu Trp Ser Ser Lys Asp Val Arg Asp Met Asp Asp  
 500 505 510  
 Ala Tyr Asp Arg Leu Thr Arg His Arg Cys Arg Met Val Glu Arg Gly  
 515 520 525  
 Ile Ala Ala Gln Pro Leu Tyr Ala Gly Tyr Cys Asn His Glu Asn Met  
 530 535 540

&lt;210&gt; 4

&lt;211&gt; 1635

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

 <223> Description of Artificial Sequence:/Note =  
 Synthetic Construct

&lt;400&gt; 4

atgctgctgg	cgctgctggt	ggcgacactg	ctggcggcga	tggtggcgct	gctgactcag	60
gtggcgctgg	tggtgcaggt	ggcggaggcg	gctcggggccc	cgagcgtctc	ggccaagccg	120
gggcccggcg	tggtggccct	gccgctcttg	gtgaagatga	ccccgaacct	gctgcatctc	180
gccccggaga	acttctacat	cagccacagc	cccaattcca	cggcggggccc	ctcctgcacc	240
ctgctggagg	aagcgtttcg	acgatatcat	ggctatatatt	ttggtttcta	caagtggcat	300
catgaacctg	ctgaattcca	ggctaaaacc	caggttcagc	aacttcttgt	ctcaatcacc	360
cttcagtcag	agtgtgatgc	tttccccaac	atatcttcag	atgagtctta	tactttactt	420
gtgaaagaac	cagtggctgt	ccttaaggcc	aacagagttt	ggggagcatt	acgagggtta	480
gagaccttta	gccagttagt	ttatcaagat	tcttatggaa	ctttcaccat	caatgaatcc	540
accattattg	attctccaag	gttttctcac	agaggaattt	tgattgatac	atccagacat	600
tatctgccag	ttaagattat	tcttaaaact	ctggatgcc	tggtttttaa	taagttta	660
gttcttcact	ggcacatagt	tgatgaccag	tctttcccat	atcagagcat	cacttttcct	720
gagttaagca	ataaaggaag	ctattctttg	tctcatgttt	atacaccaaa	tgatgtccgt	780
atgggtgattg	aatatgccag	attacgagga	attcagatcc	tgccagaatt	tgatacccct	840
gggcatacac	tatcttgggg	aaaaggtcag	aaagacctcc	tgactccatg	ttacagtaga	900
caaaacaagt	tggtactctt	tggtacctata	aaccctactc	tgaatacaac	atacagcttc	960

```

cttactacat ttttcaaaga aattagtgag gtgtttccag atcaattcat tcatttggga 1020
ggagatgaag tggaaatttaa atgttgggaa tcaaattccaa aaattcaaga tttcatgagg 1080
caaaaaggct ttggcacaga ttttaagaaa ctagaatctt tctacattca aaaggttttg 1140
gatattattg caaccataaa caagggatcc attgtctggc aggagggttt tgatgataaa 1200
gcaaagcttg cgccgggcac aatagttgaa gtatggaaa acagcgcata tcctgaggaa 1260
ctcagtagag tcacagcatc tggcttccct gtaatccttt ctgctccttg gtacttagat 1320
ttgattagct atggacaaga ttggaggaaa tactataaag tggaacctct tgattttggc 1380
ggtactcaga aacagaaaca acttttcatt ggtggagaag cttgtctatg gggagaatat 1440
gtggatgcaa ctaacctcac tccaagatta tggcctcggg caagtgtctg ttggtagaga 1500
ctctggagtt ccaaagatgt cagagatatg gatgacgcct atgacagact gacaaggcac 1560
cgctgcagga tggtcgaacg tggaatagct gcacaacctc tttatgctgg atattgtaac 1620
catgagaaca tgtaa 1635

```

&lt;210&gt; 5

&lt;211&gt; 581

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 5

```

aattccgccc ctctccctcc ccccccccta acgttactgg ccgaagccgc ttggaataag 60
gccggtgtgc gtttgtctat atgtgatttt ccaccatatt gccgtctttt ggcaatgtga 120
gggcccggaa acctggccct gtcttcttga cgagcattcc taggggtctt tcccctctcg 180
ccaaaggaat gcaaggctctg ttgaatgtcg tgaaggaagc agttcctctg gaagcttctt 240
gaagacaaac aacgtctgta gcgacccttt gcaggcagcg gaacccccca cctggcgaca 300
ggtgcctctg cggccaaaag ccacgtgtat aagatacacc tgcaaaggcg gcacaacccc 360
agtgccacgt tgtgagttgg atagttgtgg aaagagtcaa atggctctcc tcaagcgtat 420
tcaacaaggg gctgaaggat gcccagaagg taccctattg tatgggatct gatctggggc 480
ctcgggtcac atgctttaca tgtgtttagt cgagggttaa aaaacgtcta ggccccccga 540
accacgggga cgtgggttttc ctttgaaaaa cacgatgata a 581

```

&lt;210&gt; 6

&lt;211&gt; 528

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 6

```

Met Ala Gly Cys Arg Leu Trp Val Ser Leu Leu Leu Ala Ala Ala Leu
1           5           10           15
Ala Cys Leu Ala Thr Ala Leu Trp Pro Trp Pro Gln Tyr Ile Gln Thr
20           25           30
Tyr His Arg Arg Tyr Thr Leu Tyr Pro Asn Asn Phe Gln Phe Arg Tyr
35           40           45
His Val Ser Ser Ala Ala Gln Gly Gly Cys Val Val Leu Asp Glu Ala
50           55           60
Phe Arg Arg Tyr Arg Asn Leu Leu Phe Gly Ser Gly Ser Trp Pro Arg
65           70           75           80
Pro Ser Phe Ser Asn Lys Gln Gln Thr Leu Gly Lys Asn Ile Leu Val
85           90           95
Val Ser Val Val Thr Ala Glu Cys Asn Glu Phe Pro Asn Leu Glu Ser
100          105          110

```

```

Val Glu Asn Tyr Thr Leu Thr Ile Asn Asp Asp Gln Cys Leu Leu Ala
      115      120      125
Ser Glu Thr Val Trp Gly Ala Leu Arg Gly Leu Glu Thr Phe Ser Gln
      130      135      140
Leu Val Trp Lys Ser Ala Glu Gly Thr Phe Phe Ile Asn Lys Thr Lys
      145      150      155      160
Ile Lys Asp Phe Pro Arg Phe Pro His Arg Gly Val Leu Leu Asp Thr
      165      170      175
Ser Arg His Tyr Leu Pro Leu Ser Ser Ile Leu Asp Thr Leu Asp Val
      180      185      190
Met Ala Tyr Asn Lys Phe Asn Val Phe His Trp His Leu Val Asp Asp
      195      200      205
Ser Ser Phe Pro Tyr Glu Ser Phe Thr Phe Pro Glu Leu Thr Arg Lys
      210      215      220
Gly Ser Phe Asn Pro Val Thr His Ile Tyr Thr Ala Gln Asp Val Lys
      225      230      235      240
Glu Val Ile Glu Tyr Ala Arg Leu Arg Gly Ile Arg Val Leu Ala Glu
      245      250      255
Phe Asp Thr Pro Gly His Thr Leu Ser Trp Gly Pro Gly Ala Pro Gly
      260      265      270
Leu Leu Thr Pro Cys Tyr Ser Gly Ser His Leu Ser Gly Thr Phe Gly
      275      280      285
Pro Val Asn Pro Ser Leu Asn Ser Thr Tyr Asp Phe Met Ser Thr Leu
      290      295      300
Phe Leu Glu Ile Ser Ser Val Phe Pro Asp Phe Tyr Leu His Leu Gly
      305      310      315      320
Gly Asp Glu Val Asp Phe Thr Cys Trp Lys Ser Asn Pro Asn Ile Gln
      325      330      335
Ala Phe Met Lys Lys Lys Gly Phe Thr Asp Phe Lys Gln Leu Glu Ser
      340      345      350
Phe Tyr Ile Gln Thr Leu Leu Asp Ile Val Ser Asp Tyr Asp Lys Gly
      355      360      365
Tyr Val Val Trp Gln Glu Val Phe Asp Asn Lys Val Lys Val Arg Pro
      370      375      380
Asp Thr Ile Ile Gln Val Trp Arg Glu Glu Met Pro Val Glu Tyr Met
      385      390      395      400
Leu Glu Met Gln Asp Ile Thr Arg Ala Gly Phe Arg Ala Leu Leu Ser
      405      410      415
Ala Pro Trp Tyr Leu Asn Arg Val Lys Tyr Gly Pro Asp Trp Lys Asp
      420      425      430
Met Tyr Lys Val Glu Pro Leu Ala Phe His Gly Thr Pro Glu Gln Lys
      435      440      445
Ala Leu Val Ile Gly Gly Glu Ala Cys Met Trp Gly Glu Tyr Val Asp
      450      455      460
Ser Thr Asn Leu Val Pro Arg Leu Trp Pro Arg Ala Gly Ala Val Ala
      465      470      475      480
Glu Arg Leu Trp Ser Ser Asn Leu Thr Thr Asn Ile Asp Phe Ala Phe
      485      490      495
Lys Arg Leu Ser His Phe Arg Cys Glu Leu Val Arg Arg Gly Ile Gln
      500      505      510
Ala Gln Pro Ile Ser Val Gly Tyr Cys Glu Gln Glu Phe Glu Gln Thr
      515      520      525

```

&lt;210&gt; 7

&lt;211&gt; 1960

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 7

ctgcagaatc	ctttgcttac	ggatctctga	gatcgagccg	ccttgcttcc	ctcccggtca	60
cgtagaccctc	cgattgtcac	gcgggctgcc	gctcagctga	ccggggctca	cgtagggctca	120
gcctgctggc	cgaggagctg	gccggtgggc	atggccggct	gcaggtctg	ggtttcgctg	180
ctgctggcgg	cggcgttgcc	ttgcttgccc	acggcactgt	ggccgtggcc	ccagtagatc	240
caaacctacc	accggcgcta	cacctgtac	cccaacaact	tccagttccg	gtaccatgtc	300
agttcggccg	cgcaggcgcg	ctgcgtcgtc	ctcgacgagg	cctttcgacg	ctaccgtaac	360
ctgctcttcc	gttcgggctc	ttggccccga	cccagcttct	caaataaaca	gcaaacgttg	420
gggaagaaca	ttctgggtgt	ctccgtcgtc	acagctgaat	gtaatgaatt	tcctaatttg	480
gagtcggtag	aaaattacac	cctaaccatt	aatgatgacc	agtgtttact	cgctctgag	540
actgtctggg	gcgctctccg	aggtctggag	actttcagtc	agcttggttg	gaaatcagct	600
gagggcacgt	tctttatcaa	caagacaaaag	attaaagact	ttcctcgatt	ccctcaccgg	660
ggcgtactgc	tggatacatc	tgcgccattac	ctgccattgt	ctagcatcct	ggatacactg	720
gatgtcatgg	catacaataa	attcaacgtg	ttccactggc	acttggtgga	cgactcttcc	780
ttcccatatg	agagcttcac	tttcccagag	ctcaccagaa	aggggtcctt	caaccctgtc	840
actcacatct	acacagcaca	ggatgtgaag	gaggtcattg	aatacgcaag	gcttcggggg	900
atccgtgtgc	tggcagaatt	tgacactcct	ggccacactt	tgtcctgggg	gccagggtgcc	960
cctgggttat	taacaccttg	ctactctggg	tctcatctct	ctggcacatt	tggaccggtg	1020
aacccagtc	tcaacagcac	ctatgacttc	atgagcacac	tcttcttgga	gatcagctca	1080
gtcttcccg	acttttatct	ccacctggga	ggggatgaag	tcgacttcac	ctgctggaag	1140
tccaacccca	acatccaggc	cttcatgaag	aaaaagggt	ttactgactt	caagcagctg	1200
gagtccttct	acatccagac	gctgctggac	atcgtctctg	attatgacaa	gggctatgtg	1260
gtgtggcagg	aggtatttga	taataaagtg	aaggttcggc	cagatacaat	catacagggtg	1320
tggcggaagg	aaatgccagt	agagtacatg	ttggagatgc	aagatatcac	cagggctggc	1380
ttccgggccc	tgctgtctgc	tccctggtac	ctgaaccgtg	taaagtatgg	ccctgactgg	1440
aaggacatgt	acaaagtggg	gcccctggca	tttcatggta	cgctgaaca	gaaggctctg	1500
gtcattggag	gggaggcctg	tatgtgggga	gagtatgtgg	acagcaccaa	cctggtcccc	1560
agactctggc	ccagagcggg	tgcctcgtct	gagagactgt	ggagcagtaa	cctgacaact	1620
aatatagact	ttgcctttaa	acgtttgtcg	catttccgtt	gtgagctggg	gaggagagga	1680
atccaggccc	agcccatcag	tgtaggctac	tgtgagcagg	agtttgagca	gacttgagcc	1740
accagcgtcg	aacacccagg	aggttgctgt	cctttgagtc	agctgcgtcg	agcaccagg	1800
agggtgctgg	ccttaagaga	gcaggtcccg	ggcagggtct	aatctttcac	tgcctcccg	1860
caaggggaga	gcacccttg	cccgtgtgcc	cctgtgacta	cagagaagga	ggctggtgct	1920
ggcactggtg	ttcaataaag	atctatgtgg	cattttctct			1960

<210> 8

<211> 12745

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 8

atgcgggtttt	ggcagtagat	caatgggcgt	ggatagcggg	ttgactcacg	gggatttcca	60
agtctccacc	ccattgacgt	caatgggagt	ttgttttggc	acaaaaatca	acgggacttt	120
ccaaaatgtc	gtaacaactc	cgccccattg	acgcaaatgg	gcggtaggcg	tgtacgggtg	180
gaggtctata	taagcagagc	tctgtgaaac	ttcgaggagt	ctctttgttg	aggacttttg	240
agttctccct	tgaggctccc	acagatacaa	taaatatattg	agattgaacc	ctgtcgagta	300
tctgtgtaat	cttttttacc	tgtgaggtct	cggaatccgg	gccgagaact	tcgcagttgg	360
cgcccgaaaca	gggacttgat	tgagagtgat	tgaggaagtg	aagctagagc	aatagaaaagc	420
tgtaagcag	aactcctgct	gacctaaata	gggaagcagt	agcagacgct	gctaacagtg	480
agtatctcta	gtgaagcggg	ctcgagctca	taatcaagtc	attgtttaaa	ggccagata	540

aattacatct	ggtgactcct	cgcggacctt	caagccagga	gattcgccga	gggacagtca	600
acaaggtagg	agagattcta	cagcaacatg	gggaatggac	aggggcgaga	ttggaaaatg	660
gccattaaga	gatgtagtaa	tggtgctgta	ggagtagggg	ggaagagtaa	aaaatttgga	720
gaagggaatt	tcagatgggc	cattagaatg	gctaattgat	ctacaggacg	agaacctggg	780
gatataccag	agactttaga	tcaactaagg	ttgggttattt	gcgatttaca	agaaagaaga	840
gaaaaatttg	gatctagcaa	agaaattgat	atggcaattg	tgacattaaa	agtctttgcg	900
gtagcaggac	ttttaaatat	gacgggtgtc	tactgtgct	gcagctgaaa	atatgtattc	960
tcaaatggga	ttagacacta	ggccatctat	gaaggaagca	ggtggaaaaag	aggaaggccc	1020
tccacaggca	tatcctatto	aaacagtaaa	tgagtagcca	caatatgtag	cacttgaccc	1080
aaaaatggtg	tccattttta	tggaaaaggc	aagagaagga	ctaggagggtg	aggaagttca	1140
actatggttt	actgccttct	ctgcaaattt	aacacctact	gacatggcca	cattaataat	1200
ggccgcacca	gggtgcgctg	cagataaaga	aatattggat	gaaagcttaa	agcaactgac	1260
agcagaatat	gatcgcacac	atccccctga	tgctcccaga	ccattaccct	attttactgc	1320
agcagaaatt	atgggtatag	gattaactca	agaacaacaa	gcagaagcaa	gatttgcacc	1380
agctaggatg	catgtatag	catggtatct	cgaggcatta	ggaaaattgg	ctgccataaa	1440
agctaagtct	cctcgagctg	tgcaagttag	acaaggagct	aaggaagatt	attcatcctt	1500
tatagacaga	ttgtttgccc	aaatagatca	agaacaaaat	acagctgaag	ttaagttata	1560
tttaaaacag	tcattgagca	tagctaatgc	taatgcagac	tgtaaaaagg	caatgagcca	1620
ccttaagcca	gaaagtaccc	tagaagaaaa	gttgagagct	tgtcaagaaa	taggctcacc	1680
aggatataaa	atgcaactct	tggcagaagc	tcttacaaaa	gttcaagtag	tgcaatcaaa	1740
aggatcagga	ccagtgtgtt	ttaattgtaa	aaaaccagga	catctagcaa	gacaatgtag	1800
agaagtgaaa	aaatgtaata	aatgtggaaa	acctggtcat	gtagctgccca	aatgttgcca	1860
aggaaatgat	aagaattgta	caagggaaga	aagggtatca	acaattacaa	aagtgggaag	1920
attgggtagg	atggatagga	aatattccac	aatattttaa	gggactattg	ggaggtatct	1980
tgggaaatagg	attaggagtg	ttattattga	ttttatgttt	acctacattg	gttgattgta	2040
taagaaattg	tatccacaag	atactaggat	acacagtaat	tgcaatgcct	gaagtagaag	2100
gagaagaaat	acaaccacaa	atggaattga	ggagaaatgg	taggcaatgt	ggcatgtctg	2160
aaaaagagga	ggaatgatga	agtatctcag	acttatttta	taaggagat	actgtgctga	2220
gttcttcctt	ttgaggaagg	tatgtcatat	gaatccattt	cgaatcaaat	caaactaata	2280
aagtatgtat	tgtaaagtga	caaggaaaga	caaagaaaga	gaagaaagaa	gaaagccttc	2340
agtacattta	tattggctca	tgtccaatat	gaccgccatg	ttgacattga	ttattgacta	2400
gttattaata	gtaatcaatt	acgggggtcat	tagttcatag	cccataatag	gagttccgcg	2460
ttacataact	tacggttaatt	ggccgcgctg	ctgaccgccc	aacgaccccc	gcccattgac	2520
gtcaataatg	acgtatgttc	ccatagtaac	gccaataggg	actttccatt	gacgtcaatg	2580
ggtggagtat	ttacggtaaa	ctgcccactt	ggcagtagat	caagtgtatc	atatgccaaag	2640
tccggccccc	tattgacgtc	aatgacggta	aatggccgcg	ctggcattat	gccagtagca	2700
tgaccttacg	ggactttggt	acttggcagt	ccatctacgt	attagtcatc	gctattacca	2760
tggtgatgag	gtttttggcag	tacaccaatg	ggcgtggata	gcggtttgac	tcacggggat	2820
ttccaagtct	ccaccccat	gacgtcaatg	ggagtttggt	ttggcaccaa	aatcaacggg	2880
actttccaaa	atgtcgtaat	aaccccgccc	cgttgacgca	aatgggcggt	aggcgtgtac	2940
ggtgggaggt	ctatataagc	agagctcggt	tagtgaaccg	tcagatcgcc	tggagacgcc	3000
atccacgctg	ttttgacctc	catagaagac	accgggaccg	atccagcctc	cgcggccggg	3060
aacggtgcat	tggaaacgcg	attccccgtg	ccaagagtga	cgtaagtacc	gcctatagac	3120
tctataggca	cacccttttg	gtctttatgc	atgtataact	gtttttggct	tggggcctat	3180
acaccccgcc	tccttatgct	ataggtgatg	gtatagctta	gcctataggt	gtgggttatt	3240
gaccattatt	gaccactccc	ctatttggtga	cgatactttc	cattactaat	ccataacatg	3300
gctctttgcc	acaactatct	ctattggcta	tatgccataa	ctctgtcctt	cagagactga	3360
cacggactct	gtatttttac	aggatggggg	cccatthatt	atttacaat	tcacatatac	3420
aacaacgcgg	tccccgtg	ccgcagtttt	tattaaacat	agcgtgggat	ctccacgcga	3480
atctcgggta	cgtgttcggg	acatgggctc	ttctccggta	gcggcgagc	ttccacatcc	3540
gagccctggt	cccatgcctc	cagcggctca	tggtcgctcg	gcagctcctt	gtcctaaca	3600
gtggagggcca	gacttaggca	cagcacaatg	cccaccacca	ccagtgtgcc	gcacaaggcc	3660
gtggcggtag	ggtatgtgtc	tgaaaatgag	ctcggagatt	gggctcgac	cgtgacgcag	3720
atggaagact	taaggcagcg	gcagaagaag	atgcaggcag	ctgagttggt	gtattctgat	3780
aagagtcaga	ggtaactccc	gttgcggttc	tgtaaacggg	ggagggcagt	gtagtctgag	3840
cagtactcgt	tgctgccg	cgcgccacca	gacataatag	ctgacagact	aacagactgt	3900
tcctttccat	gggtcttttc	tgcagtcacc	gtcgtcgaag	cttatgacca	tgattacgga	3960
ttcactggcc	gtcgttttac	aacgtcgtga	ctgggaaaa	cctggcggtta	cccaacttaa	4020

tcgccttgca	gcacatcccc	ctttcgccag	ctggcgtaat	agcgaagagg	cccgaccgga	4080
tcgcccttcc	caacagttgc	gcagcctgaa	tggcgaatgg	cgctttgcct	ggtttccggc	4140
accagaagcg	gtgccggaaa	gctggctgga	gtgcgatctt	cctgaggccg	atactgtcgt	4200
cgtcccctca	aactggcaga	tgcacggtta	cgatgcgccc	atctacacca	acgtaaccta	4260
tcccattacg	gtcaatccgc	cgtttgttcc	cacggagaat	ccgacggggt	gttactcgct	4320
cacatttaat	gttgatgaaa	gctggctaca	ggaaggccag	acgcgaatta	tttttgatgg	4380
cgtaaactcg	gcgtttcatc	tgtggtgcaa	cgggcgctgg	gtcggttacg	gccaggacag	4440
tcgtttgccg	tctgaatttg	acctgagcgc	atttttacgc	gccggagaaa	accgcctcgc	4500
ggtgatgggtg	ctgcgttgga	gtgacggcag	ttatctggaa	gatcaggata	tgtggcggat	4560
gagcggcatt	ttccgtgacg	tctcgttgct	gcataaaccg	actacacaaa	tcagcgattt	4620
ccatgttgcc	actcgcttta	atgatgattt	cagccgcgct	gtactggagg	ctgaagtcca	4680
gatgtgcggc	gagttgcgtg	actacctacg	ggtaacagtt	tctttatggc	agggtgaaac	4740
gcaggtcgcc	agcggcaccg	cgcctttcgg	cggtgaaatt	atcgatgagc	gtggtgggta	4800
tgccgatcgc	gtcacactac	gtctgaacgt	cgaaaaccgc	aaactgtgga	gcgccgaaat	4860
ccggaatctc	tatcgtgcgg	tggttgaaat	gcacaccgcc	gacggcacgc	tgattgaaag	4920
agaagcctgc	gatgtcgggt	tccgcgaggt	cgggattgaa	aatggctctgc	tgctgctgaa	4980
cggcaagccg	ttgctgattc	gaggcgtaaa	ccgtcacgag	catcatcctc	tgcatgggtca	5040
ggtcatggat	gagcagacga	tggtgcagga	tatcctgctg	atgaagcaga	acaactttaa	5100
cgccgtgcgc	tgttcgcatt	atccgaacca	tccgctgtgg	tacacgctgt	gcgaccgcta	5160
cggcctgtat	gtggtgggatg	aagccaatat	tgaaaaccac	ggcatggtgc	caatgaatcg	5220
tctgaccgat	gatccgcgct	ggctaccggc	gatgagcgaa	cgcgtaacgc	gaatggtgca	5280
gcgcgatcgt	aatcacccga	gtgtgatcat	ctggtcgtcg	gggaatgaat	caggccacgg	5340
cgctaatac	gacgcgctgt	atcgctggat	caaactgtgc	gatccttccc	gcccgggtga	5400
gtatgaaggc	ggcggagccg	acaccacggc	caccgatatt	atttgcccga	tgtacgcgcg	5460
cgtggatgaa	gaccagccct	tccgggctgt	gccgaaatgg	tccatcaaaa	aatggctttc	5520
gctacctgga	gagacgcgcc	cgctgatcct	ttgcgaatac	gcccacgcga	tgggtaacag	5580
tcttgccggt	ttcgctaaat	actggcaggc	gtttcgtcag	tatccccgtt	tacagggcgg	5640
cttcgtcttg	gactgggtgg	atcagtcgct	gattaaatat	gatgaaaacg	gcaaccctgt	5700
gtcggcttac	ggcgtgtgatt	ttggcgatac	gccgaacgat	cgccagttct	gtatgaacgg	5760
tctggtcttt	gccgaccgca	cgcgcgcatc	atggctgacg	gaagcaaaac	accagcagca	5820
gtttttccag	ttccggtttat	ccgggcaaac	catcgaagtg	accagcgaat	acctgttccg	5880
tcatagcgat	aacgagctcc	tgcactggat	ggtggcgctg	gatggtaagc	cgctggcaag	5940
cgggtgaagtg	cctctggatg	tcgctccaca	aggtaaacag	ttgattgaac	tgcttgaact	6000
accgcagccg	gagagcgccg	ggcaactctg	gctcacagta	cgcgtagtgc	aaccgaacgc	6060
gaccgcatgg	tcagaagccg	ggcacatcag	cgcctggcag	cagtggcgctc	tggcggaaaa	6120
cctcagttgt	acgctccccg	ccgcgtccca	cgccatcccg	catctgacca	ccagcgaat	6180
ggattttttt	atcgagttgg	gtaataagcg	taagctattt	aaccgccagt	caggctttct	6240
ttcacagatg	tggattggcg	ataaaaaaca	actgctgacg	ccgctgcgcg	atcagttcac	6300
ccgtgcaccg	ctggataacg	acattggcgt	aagtgaagcg	acccgcattg	accctaacgc	6360
ctgggtcgaa	cgctggaagg	cggcgggcca	ttaccaggcc	gaagcagcgt	tggtgcagtg	6420
cacggcagat	acacttgctg	atgcggtgct	gattacgacc	gctcacgcgt	ggcagcatca	6480
ggggaaaaacc	ttattttatca	gccggaaaac	ctaccggatt	gatggtagtg	gtcaaatggc	6540
gattaccggt	gatgttgaag	tggcgagcga	tacaccgcac	ccggcgcgga	ttggcctgaa	6600
ctgccagctg	gcgcaggtag	cagagcgggt	aaactggctc	ggattagggc	cgcaagaaaa	6660
ctatcccgcg	cgcttactg	ccgcctgttt	tgaccgctgg	gatctgccat	tgtcagacat	6720
gtataccccg	tacgtcttcc	cgagcgaaaa	cggctctgcgc	tgccggacgc	gcgaattgaa	6780
ttatggccca	caccagtggc	gcggcgactt	ccagttcaac	atcagccgct	acagtcaaca	6840
gcaactgatg	gaaaccagcc	atcgccatct	gctgcacgcg	gaagaaggca	catggctgaa	6900
tatcgacggt	ttccatatgg	ggattggtgg	cgacgactcc	tggagcccgt	cagtatcggc	6960
ggaattccag	ctgagcgccg	gtcgtctacca	ttaccagttg	gtctggtgtc	aaaaataact	7020
cgatcgacca	gagctgagat	cctacaggag	tccagggtcg	gagagaaaaa	ctctgaagag	7080
gatgatgaca	gagttagaag	atcgcttcag	gaagctattt	ggcacgactt	ctacaacggg	7140
agacagcaca	gtagattctg	aagatgaacc	tcctaaaaaa	gaaaaaagg	tggactggga	7200
tgagtatttg	aaccctgaag	aaatagaaag	aatgcttatg	gactagggac	tgtttacgaa	7260
caaatgataa	aaggaaatag	ctgagcatga	ctcatagtta	aagcgctagc	agctgcctaa	7320
ccgcaaaacc	acatcctatg	gaaagcttgc	taatgacgta	taagttgttc	cattgtaaga	7380
gtatataacc	agtgtcttgt	gaaacttcga	ggagtctctt	tgttgaggac	ttttgagttc	7440
tcctttgagg	ctcccacaga	tacaataaat	atttgagatt	gaaccctgtc	gagtatctgt	7500
gtaatctttt	ttacctgtga	ggtctcggaa	tccgggcccga	gaacttcgca	gcggccgctc	7560



gagcatgcat	ctagagggcc	ctattctata	gtgtcaccta	aatgctagag	ctcgtgatac	7620
agcctcgact	gtgccttcta	gttgccagcc	atctgttggt	tgccctccc	ccgtgccttc	7680
cttgaccctg	gaaggtgcca	ctccactgt	cctttcctaa	taaaatgagg	aaattgcatc	7740
gcattgtctg	agtaggtgtc	attctattct	ggggggtggg	gtggggcagg	acagcaaggg	7800
ggaggattgg	gaagacaata	gcaggcatgc	tggggatgcg	gtgggtctta	tggcttctga	7860
ggcggaaga	accagctggg	gctcgagggg	ggatccccac	gcgccctgta	gcggcgcatc	7920
aagcgcgcg	ggtgtggtgg	ttacgcgcag	cgtgaccgct	acacttgcca	gcgccctagc	7980
gcccgctcct	tctcgcttct	tcccttccct	tctcgccacg	ttcgccggct	ttccccgtca	8040
agctctaaat	cggggcatcc	ctttagggtt	ccgatttagt	gctttacggc	acctcgaccc	8100
caaaaaactt	gattagggtg	atggttcacg	tagtgggcca	tcgccctgat	agacggtttt	8160
tcgccctttg	acgttgaggt	ccacgttctt	taatagtggg	ctcttggtcc	aaactggaac	8220
aacactcaac	cctatctcgg	tctattcttt	tgatttataa	gggatttttg	ggatttcggc	8280
ctatttggtta	aaaaatgagc	tgatttaaca	aaaatttaac	gcgaatttta	acaaaatatt	8340
aacgtttaca	atttaaatat	ttgcttatac	aatcttccct	tttttggggc	ttttctgatt	8400
atcaaccggg	gtgggtacog	agctcgaaat	ctgtggaatg	tgtgtcagtt	aggggtgtgga	8460
aagtccccag	gctccccagg	caggcagaag	tatgcaaagc	atgcatctca	attagtacgc	8520
aaccaggtgt	ggaaagtccc	caggctcccc	agcaggcaga	agtagcaaaa	gcagtcacat	8580
caattagtca	gcaaccatag	tcccgcacct	aactccgccc	atcccgcccc	taactccgcc	8640
cagttccgcc	cattctccgc	cccatggctg	actaatTTTT	tttatttatg	cagaggccga	8700
ggccgcctcg	gcctctgagc	tattccagaa	gtagtggagg	ggcttttttg	gaggcctagg	8760
cttttgcaaa	aagctcccg	gagcttggat	atccattttc	ggatctgata	aagagacagg	8820
atgaggatcg	tttcgcatga	ttgaacaaga	tggattgcac	gcaggttctc	cggccgcttg	8880
ggtggagagg	ctattcggct	atgactgggc	acaacagaca	atcggtgct	ctgatgccgc	8940
cgtgttccgg	ctgtcagcgc	aggggcgccc	ggttcttttt	gtcaagaccg	acctgtccgg	9000
tgccctgaat	gaactgcagg	acgaggcagc	gcggctatcg	tggctggcca	cgacgggcgt	9060
tccttgccga	gctgtgctcg	acgttgtcac	tgaagcggga	agggactggc	tgctattggg	9120
cgaagtgccg	gggcaggatc	tccgtgcac	tcaccttgct	cctgccgaga	aagtatccat	9180
catggctgat	gcaatgcggc	ggctgcatac	gcttgatccg	gctacctgcc	cattcgacca	9240
ccaagcgaaa	catcgcatcg	agcagcacgc	tactcgatg	gaagccggtc	ttgtcgatca	9300
ggatgatctg	gacgaagagc	atcaggggct	cgcgccagcc	gaactgttcg	ccaggctcaa	9360
ggcgcgcag	cccagcggcg	aggatctcgt	cgtgacccat	ggcgatgcct	gcttgccgaa	9420
tatcatgggt	gaaaatggcc	gcttttctgg	attcatcgac	tgtggccggc	tgggtgtggc	9480
ggaccgctat	caggacatag	cgttggctac	ccgtgatatt	gctgaagagc	ttggcggcga	9540
atgggctgac	cgttccctcg	tgttttacgg	tatcgccgct	cccgattcgc	agcgcacgc	9600
cttctatcgc	cttcttgacg	agttcttctg	agcgggactc	tggggttcga	aatgaccgac	9660
caagcgacgc	ccaactggcc	atcacgagat	tactgattcca	ccgcgcctt	ctatgaaagg	9720
ttgggcttcg	gaatcgttt	cgggagcgc	ggctggatga	tcctccagcg	cggggatctc	9780
atgctggagt	tcttcgcccc	ccccaaactg	tttattgcag	cttataatgg	ttacaaataa	9840
agcaatagca	tcacaaattt	cacaaataaa	gcattttttt	cactgcattc	tagttgtggt	9900
ttgtccaaac	tcataaatgt	atcttatcat	gtctggatcc	cgtcgacctc	gagagcttgg	9960
cgtaatcatg	gtcatagctg	tttccctgtg	gaaattgtta	tccgctcaca	attccacaca	10020
acatacgagc	cgggaagcata	aagtgtaaaag	cctgggggtgc	ctaagtgtg	agctaactca	10080
cattaatttg	gttgcgtcga	ctgcccgctt	tccagtcggg	aaacctgtcg	tgccagctgc	10140
attaatgaat	ggccaacgc	gcggggagag	gcggtttgcg	tattgggcgc	tcttccgctt	10200
cctcgctcac	tgactcgctg	cgtcgggtcg	ttcggtgctg	gcgagcggtg	tcagctcact	10260
caaaggcggt	aatacggtta	tccacagaat	caggggataa	cgcaggaaag	aacatgtgag	10320
caaaaaggcca	gcaaaaggcc	aggaaccgta	aaaaggccgc	gttgctggcg	ttttcccata	10380
ggctccgccc	ccctgacgag	catcacaaaa	atcgacgctc	aagttagagg	tgccgaaacc	10440
cgacaggact	ataaagatac	caggcgtttc	cccctggaag	ctccctcggt	cgctctcctg	10500
ttccgacctt	gccgcttacc	ggatacctgt	ccgcctttct	cccttcggga	agcgtggcgc	10560
tttctcaatg	ctcacgctgt	aggtatctca	gttcggtgta	ggtcgttcgc	tccaagctgg	10620
gctgtgtgca	cgaacccccc	gttcagcccc	accgctgcgc	cttatccggt	aactatcgct	10680
ttgagtccaa	cccggtaaga	cacgacttat	cgccactggc	agcagccact	ggtaacagga	10740
ttagcagagc	gaggtatgta	ggcgggtgta	cagagttctt	gaagtgggtg	cctaactacg	10800
gctacactag	aaggacagta	tttggtatct	gcgctctgct	gaagccagtt	accttcggaa	10860
aaagagttgg	tagctcttga	tccggcaaac	aaaccaccgc	tggtagcggt	ggtttttttg	10920
tttgcaagca	gcagattacg	cgcagaaaaa	aaggatctca	agaagatcct	ttgatctttt	10980
ctacggggtc	tgacgctcag	tggaaacgaa	actcacgtta	agggattttg	gtcatgagat	11040
tatcaaaaag	gatcttcacc	tagatccttt	taaattaaaa	atgaagtttt	aaatcaatct	11100

```

aaagtatata tgagtaaact tggctctgaca gttaccaatg cttaatcagt gaggcaccta 11160
tctcagcgat ctgtctatatt cgttcatcca tagttgcctg actccccgtc gtgtagataa 11220
ctacgatacg ggagggttta ccatctggcc ccagtgtctg aatgataccg cgagaccac 11280
gtccaccggc tccagattta tcagcaataa accagccagc cggaagggcc gagcgagaa 11340
gtggtcctgc aactttatcc gcctccatcc agtctattaa ttggtgccgg gaagctagag 11400
taagtagttc gccagttaat agtttgcgca acgttggtgc cattgctaca ggcatcgtgg 11460
tgtcacgtc gtcgtttggg atggcttcat tcagctccgg ttcccaacga tcaaggcgag 11520
ttacatgatc ccccatggtg tgcaaaaaag cggttagctc cttcggtcct ccgatcgttg 11580
tcagaagtaa gttggccgca gtgttatcac tcatggttat ggcagcactg cataattctc 11640
ttactgtcat gccatccgta agatgctttt ctgtgactgg tgagtactca accaagtcac 11700
tctgagaata gtgtatgcgg cgaccgagtt gctcttgccc ggcgtaataa cgggataata 11760
ccgcgccaca tagcagaact ttaaaagtgc tcatcattgg aaaacgttct tcggggcgaa 11820
aactctcaag gatcttaccg ctggtgagat ccagttcgat gtaaccact cgtgcaccca 11880
actgatcttc agcatctttt actttcacca gcgtttctgg gtgagcaaaa acaggaaggc 11940
aaaatgccgc aaaaaaggga ataaggcgca caccgaaatg ttgaatactc atactcttcc 12000
tttttcaata ttattgaagc atttatcagg gttattgtct catgagcgga tacatatattg 12060
aatgtattta gaaaaataaa caaatagggg ttccgcgcac atttccccga aaagtgccac 12120
ctgacgtcga cggatcggga gatctcccga tcccctatgg tcgactctca gtacaatctg 12180
ctctgatgcc gcatagttaa gccagtatct gctccctgct tgtgtgttg aggtcgctga 12240
gtagtgcgcy agcaaaattht aagctacaac aaggcaaggc ttgaccgaca attgcatgaa 12300
gaatctgctt aggggttaggc gttttgcgct gcttcgcgat gtacgggcca gatatacgcy 12360
ttgacattga ttattgacta gttattaata gtaatcaatt acgggggtcat tagttcatag 12420
cccatatatg gagttccgcy ttacataact tacggtaaat ggccgcctg gctgaccgcc 12480
caacgacccc cgccattga cgtcaataat gacgtatgtt cccatagtaa cgccaatagg 12540
gactttccat tgacgtcaat ggggtgacta tttacggtaa actgccact tggcagtaca 12600
tcaagtgtat catatgcca gtacgcccc tattgacgtc aatgacggt aatggccgc 12660
ctggcattat gccagtaga tgaccttat ggactttcct acttggcagt acatctacgt 12720
attagtcac gctattacca tgggtg 12745

```

&lt;210&gt; 9

&lt;211&gt; 529

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 9

```

Met Thr Ser Ser Arg Leu Trp Phe Ser Leu Leu Leu Ala Ala Ala Phe
1          5          10          15
Ala Gly Arg Ala Thr Ala Leu Trp Pro Trp Pro Gln Asn Phe Gln Thr
20          25          30
Ser Asp Gln Arg Tyr Val Leu Tyr Pro Asn Asn Phe Gln Phe Gln Tyr
35          40          45
Asp Val Ser Ser Ala Ala Gln Pro Gly Cys Ser Val Leu Asp Glu Ala
50          55          60
Phe Gln Arg Tyr Arg Asp Leu Leu Phe Gly Ser Gly Ser Trp Pro Arg
65          70          75          80
Pro Tyr Leu Thr Gly Lys Arg His Thr Leu Glu Lys Asn Val Leu Val
85          90          95
Val Ser Val Val Thr Pro Gly Cys Asn Gln Leu Pro Thr Leu Glu Ser
100         105         110
Val Glu Asn Tyr Thr Leu Thr Ile Asn Asp Asp Gln Cys Leu Leu Leu
115         120         125
Ser Glu Thr Val Trp Gly Ala Leu Arg Gly Leu Glu Thr Phe Ser Gln
130         135         140

```

12/37

Leu Val Trp Lys Ser Ala Glu Gly Thr Phe Phe Ile Asn Lys Thr Glu  
 145 150 155 160  
 Ile Glu Asp Phe Pro Arg Phe Pro His Arg Gly Leu Leu Leu Asp Thr  
 165 170 175  
 Ser Arg His Tyr Leu Pro Leu Ser Ser Ile Leu Asp Thr Leu Asp Val  
 180 185 190  
 Met Ala Tyr Asn Lys Leu Asn Val Phe His Trp His Leu Val Asp Asp  
 195 200 205  
 Pro Ser Phe Pro Tyr Glu Ser Phe Thr Phe Pro Glu Leu Met Arg Lys  
 210 215 220  
 Gly Ser Tyr Asn Pro Val Thr His Ile Tyr Thr Ala Gln Asp Val Lys  
 225 230 235 240  
 Glu Val Ile Glu Tyr Ala Arg Leu Arg Gly Ile Arg Val Leu Ala Glu  
 245 250 255  
 Phe Asp Thr Pro Gly His Thr Leu Ser Trp Gly Pro Gly Ile Pro Gly  
 260 265 270  
 Leu Leu Thr Pro Cys Tyr Ser Gly Ser Glu Pro Ser Gly Thr Phe Gly  
 275 280 285  
 Pro Val Asn Pro Ser Leu Asn Asn Thr Tyr Glu Phe Met Ser Thr Phe  
 290 295 300  
 Phe Leu Glu Val Ser Ser Val Phe Pro Asp Phe Tyr Leu His Leu Gly  
 305 310 315 320  
 Gly Asp Glu Val Asp Phe Thr Cys Trp Lys Ser Asn Pro Glu Ile Gln  
 325 330 335  
 Asp Phe Met Arg Lys Lys Gly Phe Gly Glu Asp Phe Lys Gln Leu Glu  
 340 345 350  
 Ser Phe Tyr Ile Gln Thr Leu Leu Asp Ile Val Ser Ser Tyr Gly Lys  
 355 360 365  
 Gly Tyr Val Val Trp Gln Glu Val Phe Asp Asn Lys Val Lys Ile Gln  
 370 375 380  
 Pro Asp Thr Ile Ile Gln Val Trp Arg Glu Asp Ile Pro Val Asn Tyr  
 385 390 395 400  
 Met Lys Glu Leu Glu Leu Val Thr Lys Ala Gly Phe Arg Ala Leu Leu  
 405 410 415  
 Ser Ala Pro Trp Tyr Leu Asn Arg Ile Ser Tyr Gly Pro Asp Trp Lys  
 420 425 430  
 Asp Phe Tyr Val Val Glu Pro Leu Ala Phe Glu Gly Thr Pro Glu Gln  
 435 440 445  
 Lys Ala Leu Val Ile Gly Gly Glu Ala Cys Met Trp Gly Glu Tyr Val  
 450 455 460  
 Asp Asn Thr Asn Leu Val Pro Arg Leu Trp Pro Arg Ala Gly Ala Val  
 465 470 475 480  
 Ala Glu Arg Leu Trp Ser Asn Lys Leu Thr Ser Asp Leu Thr Phe Ala  
 485 490 495  
 Tyr Glu Arg Leu Ser His Phe Arg Cys Glu Leu Leu Arg Arg Gly Val  
 500 505 510  
 Gln Ala Gln Pro Leu Asn Val Gly Phe Cys Glu Gln Glu Phe Glu Gln  
 515 520 525  
 Thr

&lt;210&gt; 10

&lt;211&gt; 2255

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 10  
 cctccgagag gggagaccag cgggccatga caagctccag gctttggttt tcgctgctgc 60  
 tggcggcagc gttcgagga cggcgacgg ccctctggcc ctggcctcag aacttccaaa 120  
 cctccgacca gcgctacgtc ctttaccoga acaactttca attccagtac gatgtcagct 180  
 cggccgcgca gcccggtgc tcagtcctcg acgaggcctt ccagcgctat cgtgacctgc 240  
 ttttcggttc cgggtcttgg ccccgctcctt acctcacagg gaaacggcat aacttgga 300  
 agaatgtgtt ggttgtctct gtagtcacac ctggatgtaa ccagcttcct actttggagt 360  
 cagtggagaa ttataccctg accataaatg atgaccagt tttactcctc tctgagactg 420  
 tctggggagc tctccgaggt ctggagactt ttaggcagct tgtttggaaa tctgctgagg 480  
 gcacattctt tatcaacaag actgagattg aggactttcc ccgctttcct caccggggct 540  
 tgctgttggg tacatctcgc cattacctgc cactctctag catcctggac actctggatg 600  
 tcatggcgta caataaattg aacgtgttcc actggcatct ggtagatgat ccttctctcc 660  
 catatgagag cttcactttt ccagagctca tgagaaagg gtcctacaac cctgtcacc 720  
 acatctacac agcacaggat gtgaaggagg tcattgaata cgacaggctc cggggtatcc 780  
 gtgtgcttgc agagtttgac actcctggcc acactttgtc ctggggacca ggtatccctg 840  
 gattactgac tcttctgtac tctgggtctg agccctctgg cacctttgga ccagtgaatc 900  
 ccagtctcaa taatacctat gagttcatga gcacattctt cttagaagtc agctctgtct 960  
 tcccagattt ttatcttcat cttggaggag atgaggttga tttcacctgc tggaggtcca 1020  
 acccagagat ccaggacttt atgaggaaga aaggcttcgg tgaggacttc aagcagctgg 1080  
 agtcttcta catccagacg ctgctggaca tctctcttcc ttatggcaag ggctatgtgg 1140  
 tgtggcagga ggtgtttgat aataaagtaa agattcagcc agacacaatc atacagggtg 1200  
 ggcgagagga tattccagtg aactatatga aggagctgga actggtcacc aaggccggct 1260  
 tccgggccct tctctctgcc ccctgggtacc tgaaccgtat atcctatggc cctgactgga 1320  
 aggatttcta cgtagtggaa cccctggcat ttgaaggtag ccctgagcag aaggctctgg 1380  
 tgattggtgg agaggcttgt atgtggggag aatatgtgga caacacaaac ctggtcccca 1440  
 ggctctggcc cagagcaggg gctgttgccg aaaggctgtg gagcaacaag ttgacatctg 1500  
 acctgacatt tgcctatgaa cgtttgtcac acttccgctg tgagttgctg aggcgaggtg 1560  
 tccaggccca acccctcaat gtaggcttct gtgagcagga gtttgaacag acctgagccc 1620  
 caggcaccga ggagggtgct ggctgtaggt gaatgtagt ggagccaggc ttccactgca 1680  
 tcttgccag gggacggagc cccttgccct cgtgcccctt gcctgctgct cctgtgctt 1740  
 ggagagaaaag gggccggtgc tggcgctcgc attcaataaa gagtaatgtg gcattttct 1800  
 ataataaaca tggattacct gtgtttaaaa aaaaaagtggt gaatggcggt agggtaagg 1860  
 cacagccagg ctggagtcag tgtctgccc tgaggtcttt taagttgagg gctgggaatg 1920  
 aaacctatag cctttgtgct gttctgcctt gcctgtgagc tatgtcactc cctcccact 1980  
 cctgaccata ttccagacac ctgccctaact cctcagcctg ctacttcac ttctgcatta 2040  
 tatctccaag gcgttggtat atggaaaaag atgtaggggc ttggagggtg tctggacagt 2100  
 ggggagggct ccagaccaca cctggtcaca aaagagcctc tcccccatgc atactcatcc 2160  
 acctccctcc cctagagcta ttctcctttg ggtttcttgc tgctgcaatt ttatacaacc 2220  
 attatttaaa tattattaaa cacatattgt tctct 2255

<210> 11

<211> 1635

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 11  
 atgctactgg cgctgctgtt ggcgacactg ctggcggcga tgttggcgct gctgactcag 60  
 gtggcgctgg tgggtgcagg ggcggaggcg gctcgggccc cgagcgtctc ggccaagccg 120  
 gggccggcgc tgtggcccct gccgctcttg gtgaagatga ccccgaaact gctgcacttc 180  
 gccccggaga acttctacat cagccacagc cccaattcca cggcgggccc ctctgcacc 240  
 ctgctggagg aagcgtttcg acgatatcat ggctatattt ttggtttcta caagtggcat 300  
 catgaacctg ctgaattcca ggctaaaacc caggttcagc aacttcttgt ctcaatcacc 360

```

cttcagtcag agtgtgatgc tttccccaac atatcttcag atgagtcctta tactttactt 420
gtgaaagaac cagtggctgt ccttaaggcc aacagagttt ggggagcatt acgagggtta 480
gagaccttta gccagttagt ttatcaagat tcttatggaa ctttcaccat caatgaatcc 540
accattattg attctccaag gttttctcac agaggaattt tgattgatac atccagacat 600
tatctgccag ttaagattat tcttaaaact ctggatgcca tggcttttaa taagttaa 660
gttcttcaact ggcacatagt tgatgaccag tctttcccat atcagagcat cacttttcct 720
gagttaagca ataaaggaag ctattctttg tctcatgttt atacacaaa tgatgtccgt 780
atggtgattg aatatgccag attacgagga attcgagtc tgcagaatt tgatacccct 840
gggcatacac tatcttgggg aaaaggtcag aaagacctcc tgactccatg ttacagtaga 900
caaaacaagt tggactcttt tggacctata aaccctactc tgaatacaac atacagcttc 960
cttactacat ttttcaaaga aattagttag gtgtttccag atcaattcat tcatttgggg 1020
ggagatgaag tggaatttaa atgttgggaa tcaaatccaa aaattcaaga tttcatgagg 1080
caaaaaggct ttggcacaga ttttaagaaa ctagaatctt tctacattca aaaggttttg 1140
gatattattg caaccataaa caagggatcc attgtctggc aggaggtttt tgatgataaa 1200
gcaaagcttg cgccgggcac aatagttaga gtatggaaa acagcgcata tcctgaggaa 1260
ctcagtagag tcacagcatc tggcttccct gtaatccttt ctgctccttg gtacttagat 1320
ttgattagct atggacaaga ttggaggaaa tactataaag tggaaacctct tgattttggc 1380
ggtactcaga aacagaaaca acttttcatt ggtggagaag cttgtctatg gggagaatat 1440
gtggatgcaa ctaacctcac tccaagatta tggcctcggg caagtgtctg tggtagaga 1500
ctctggagtt ccaaagatgt cagagatatg gatgacgct atgacagact gacaaggcac 1560
cgctgcagga tggtcgaacg tggaatagct gcacaacctc tttatgctgg atattgtaac 1620
catgagaaca tgtaa 1635

```

&lt;210&gt; 12

&lt;211&gt; 544

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 12

```

Met Leu Leu Ala Leu Leu Ala Thr Leu Leu Ala Ala Met Leu Ala
1          5          10          15
Leu Leu Thr Gln Ile Ala Leu Val Val Gln Val Ala Glu Ala Ala Arg
20          25          30
Ala Pro Ser Val Ser Ala Lys Pro Gly Pro Ala Leu Trp Pro Leu Pro
35          40          45
Leu Leu Val Lys Met Thr Pro Asn Leu Leu His Leu Ala Pro Glu Asn
50          55          60
Phe Tyr Ile Ser His Ser Pro Asn Ser Thr Ala Gly Pro Ser Cys Thr
65          70          75          80
Leu Leu Glu Glu Ala Phe Arg Arg Tyr His Gly Tyr Ile Phe Gly Phe
85          90          95
Tyr Lys Trp His His Glu Pro Ala Glu Phe Gln Ala Lys Thr Gln Val
100         105         110
Gln Gln Leu Leu Val Ser Ile Thr Leu Gln Ser Glu Cys Asp Ala Phe
115         120         125
Pro Asn Ile Ser Ser Asp Glu Ser Tyr Thr Leu Leu Val Lys Glu Pro
130         135         140
Val Ala Val Leu Lys Ala Asn Arg Val Trp Gly Ala Leu Arg Gly Leu
145         150         155         160
Glu Thr Phe Ser Gln Leu Val Tyr Gln Asp Ser Tyr Gly Thr Phe Thr
165         170         175
Ile Asn Glu Ser Thr Ile Ile Asp Ser Pro Arg Phe Ser His Arg Gly
180         185         190
Ile Leu Ile Asp Thr Ser Arg His Tyr Leu Pro Val Lys Ile Ile Leu

```

195	200	205
Lys Thr Leu Asp Ala Met	Ala Phe Asn Lys Phe Asn Val Leu His Trp	
210	215	220
His Ile Val Asp Asp Gln Ser Phe Pro Tyr Gln Ser Ile Thr Phe Pro		240
225	230	235
Glu Leu Ser Asn Lys Gly Ser Tyr Ser Leu Ser His Val Tyr Thr Pro		255
245	250	255
Asn Asp Val Arg Met Val Ile Glu Tyr Ala Arg Leu Arg Gly Ile Arg		270
260	265	270
Val Leu Pro Glu Phe Asp Thr Pro Gly His Thr Leu Ser Trp Gly Lys		285
275	280	285
Gly Gln Lys Asp Leu Leu Thr Pro Cys Tyr Ser Arg Gln Asn Lys Leu		300
290	295	300
Asp Ser Phe Gly Pro Ile Asn Pro Thr Leu Asn Thr Thr Tyr Ser Phe		320
305	310	315
Leu Thr Thr Phe Phe Lys Glu Ile Ser Glu Val Phe Pro Asp Gln Phe		335
325	330	335
Ile His Leu Gly Gly Asp Glu Val Glu Phe Lys Cys Trp Glu Ser Asn		350
340	345	350
Pro Lys Ile Gln Asp Phe Met Arg Gln Lys Gly Phe Gly Thr Asp Phe		365
355	360	365
Lys Lys Leu Glu Ser Phe Tyr Ile Gln Lys Val Leu Asp Ile Ile Ala		380
370	375	380
Thr Ile Asn Lys Gly Ser Ile Val Trp Gln Glu Val Phe Asp Asp Lys		400
385	390	395
Ala Lys Leu Ala Pro Gly Thr Ile Val Glu Val Trp Lys Asp Ser Ala		415
405	410	415
Tyr Pro Glu Glu Leu Ser Arg Val Thr Ala Ser Gly Phe Pro Val Ile		430
420	425	430
Leu Ser Ala Pro Trp Tyr Leu Asp Leu Ile Ser Tyr Gly Gln Asp Trp		445
435	440	445
Arg Lys Tyr Tyr Lys Val Glu Pro Leu Asp Phe Gly Gly Thr Gln Lys		460
450	455	460
Gln Lys Gln Leu Phe Ile Gly Gly Glu Ala Cys Leu Trp Gly Glu Tyr		480
465	470	475
Val Asp Ala Thr Asn Leu Thr Pro Arg Leu Trp Pro Arg Ala Ser Ala		495
485	490	495
Val Gly Glu Arg Leu Trp Ser Ser Lys Asp Val Arg Asp Met Asp Asp		510
500	505	510
Ala Tyr Asp Arg Leu Thr Arg His Arg Cys Arg Met Val Glu Arg Gly		525
515	520	525
Ile Ala Ala Gln Pro Leu Tyr Ala Gly Tyr Cys Asn His Glu Asn Met		540
530	535	540

&lt;210&gt; 13

&lt;211&gt; 529

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 13

Met Thr Ser Ser Arg Leu Trp Phe Ser Leu Leu Leu Ala Ala Ala Phe	
1	5
Ala Gly Arg Ala Thr Ala Leu Trp Pro Trp Pro Gln Asn Phe Gln Thr	10
20	25
Ser Asp Gln Arg Tyr Val Leu Tyr Pro Asn Asn Phe Gln Phe Gln Tyr	30



500                      505                      510  
 Gln Ala Gln Pro Leu Asn Val Gly Phe Cys Glu Gln Glu Phe Glu Gln  
           515                      520                      525  
 Thr

&lt;210&gt; 14

&lt;211&gt; 739

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
 Synthetic Construct

&lt;400&gt; 14

ttttaatcct	ccgtttttct	gcttctgaag	ttacttcagc	ctggcaagtc	ctttacctcc	60
ccgtaggcct	ggcgagctgc	atcacaacat	tcaagattca	ccctagagcc	atctgggaaa	120
ctttcttctc	caggtcgccc	tgcgtcctcg	cctccccacc	ccgttcttct	cgagtcgggt	180
gagctgtcta	gttccatcac	ggccggcacg	gccgcagggg	tggccgggta	tttactgctc	240
tactggggcc	gtgagcagtc	tggcgagccg	agcagttgcc	gacgcccggc	acaatccgct	300
gcacgtagca	ggagcctcag	gtccaggccg	gaagtgaag	ggcaggggtg	gggtcctcct	360
ggggtcgcag	gcgcagagcc	gcctctggtc	acgtgattcg	ccgataagtc	acgggggcgc	420
cgctcacctg	accaggggtc	cacgtggcca	gccccctccg	agaggggaga	ccagcgggcc	480
atgacaagct	ccaggctttg	gttttcgctg	ctgctggcgg	cagcgttcgc	aggacgggcg	540
acggccctct	ggccctggcc	tcagaacttc	caaacctccg	accagcgcta	cgtcctttac	600
ccgaacaact	ttcaattcca	gtacgatgtc	agctcggccg	cgcagcccgg	ctgctcagtc	660
ctcgacgagg	ccttccagcg	ctatcgtgac	ctgcttttcg	gttccgggtc	ttggccccgt	720
ccttacctca	cagtgagtg					739

&lt;210&gt; 15

&lt;211&gt; 556

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
 Synthetic Construct

&lt;400&gt; 15

Met	Glu	Leu	Cys	Gly	Leu	Gly	Leu	Pro	Arg	Pro	Pro	Met	Leu	Leu	Ala
1				5					10					15	
Leu	Leu	Leu	Ala	Thr	Leu	Leu	Ala	Ala	Met	Leu	Ala	Leu	Leu	Thr	Gln
			20					25					30		
Val	Ala	Leu	Val	Val	Gln	Val	Ala	Glu	Ala	Ala	Arg	Ala	Pro	Ser	Val
		35					40					45			
Ser	Ala	Lys	Pro	Gly	Pro	Ala	Leu	Trp	Pro	Leu	Pro	Leu	Ser	Val	Lys
	50					55					60				
Met	Thr	Pro	Asn	Leu	Leu	His	Leu	Ala	Pro	Glu	Asn	Phe	Tyr	Ile	Ser
65					70					75				80	
His	Ser	Pro	Asn	Ser	Thr	Ala	Gly	Pro	Ser	Cys	Thr	Leu	Leu	Glu	Glu
				85					90					95	
Ala	Phe	Arg	Arg	Tyr	His	Gly	Tyr	Ile	Phe	Gly	Phe	Tyr	Lys	Trp	His
			100					105					110		
His	Glu	Pro	Ala	Glu	Phe	Gln	Ala	Lys	Thr	Gln	Val	Gln	Gln	Leu	Leu
	115						120					125			
Val	Ser	Ile	Thr	Leu	Gln	Ser	Glu	Cys	Asp	Ala	Phe	Pro	Asn	Ile	Ser
	130					135					140				
Ser	Asp	Glu	Ser	Tyr	Thr	Leu	Leu	Val	Lys	Glu	Pro	Val	Ala	Val	Leu



145					150					155				160
Lys	Ala	Asn	Arg	Val	Trp	Gly	Ala	Leu	Arg	Gly	Leu	Glu	Thr	Phe
				165					170					175
Gln	Leu	Val	Tyr	Gln	Asp	Ser	Tyr	Gly	Thr	Phe	Thr	Ile	Asn	Glu
			180					185					190	
Thr	Ile	Ile	Asp	Ser	Pro	Arg	Phe	Ser	His	Arg	Gly	Ile	Leu	Ile
		195					200					205		Asp
Thr	Ser	Arg	His	Tyr	Leu	Pro	Val	Lys	Ile	Ile	Leu	Lys	Thr	Leu
	210					215					220			Asp
Ala	Met	Ala	Phe	Asn	Lys	Phe	Asn	Val	Leu	His	Trp	His	Ile	Val
225					230					235				240
Asp	Gln	Ser	Phe	Pro	Tyr	Gln	Ser	Ile	Thr	Phe	Pro	Glu	Leu	Ser
			245						250					255
Lys	Gly	Ser	Tyr	Ser	Leu	Ser	His	Val	Tyr	Thr	Pro	Asn	Asp	Val
		260					265						270	Arg
Met	Val	Ile	Glu	Tyr	Ala	Arg	Leu	Arg	Gly	Ile	Arg	Val	Leu	Pro
	275					280						285		Glu
Phe	Asp	Thr	Pro	Gly	His	Thr	Leu	Ser	Trp	Gly	Lys	Gly	Gln	Lys
	290					295					300			Asp
Leu	Leu	Thr	Pro	Cys	Tyr	Ser	Arg	Gln	Asn	Lys	Leu	Asp	Ser	Phe
305					310					315				320
Pro	Ile	Asn	Pro	Thr	Leu	Asn	Thr	Thr	Tyr	Ser	Phe	Leu	Thr	Thr
			325					330						335
Phe	Lys	Glu	Ile	Ser	Glu	Val	Phe	Pro	Asp	Gln	Phe	Ile	His	Leu
		340					345						350	Gly
Gly	Asp	Glu	Val	Glu	Phe	Lys	Cys	Trp	Glu	Ser	Asn	Pro	Lys	Ile
	355					360						365		Gln
Asp	Phe	Met	Arg	Gln	Lys	Gly	Phe	Gly	Thr	Asp	Phe	Lys	Lys	Leu
	370					375				380				Glu
Ser	Phe	Tyr	Ile	Gln	Lys	Val	Leu	Asp	Ile	Ile	Ala	Thr	Ile	Asn
385					390				395					Lys
Gly	Ser	Ile	Val	Trp	Gln	Glu	Val	Phe	Asp	Asp	Lys	Ala	Lys	Leu
			405					410						415
Pro	Gly	Thr	Ile	Val	Glu	Val	Trp	Lys	Asp	Ser	Ala	Tyr	Pro	Glu
		420					425						430	Glu
Leu	Ser	Arg	Val	Thr	Ala	Ser	Gly	Phe	Pro	Val	Ile	Leu	Ser	Ala
	435					440						445		Pro
Trp	Tyr	Leu	Asp	Leu	Ile	Ser	Tyr	Gly	Gln	Asp	Trp	Arg	Lys	Tyr
	450					455				460				Tyr
Lys	Val	Glu	Pro	Leu	Asp	Phe	Gly	Gly	Thr	Gln	Lys	Gln	Lys	Gln
465					470				475					480
Phe	Ile	Gly	Gly	Glu	Ala	Cys	Leu	Trp	Gly	Glu	Tyr	Val	Asp	Ala
			485					490						495
Asn	Leu	Thr	Pro	Arg	Leu	Trp	Pro	Arg	Ala	Ser	Ala	Val	Gly	Glu
		500					505						510	Arg
Leu	Trp	Ser	Ser	Lys	Asp	Val	Arg	Asp	Met	Asp	Asp	Ala	Tyr	Asp
	515					520						525		Arg
Leu	Thr	Arg	His	Arg	Cys	Arg	Met	Val	Glu	Arg	Gly	Ile	Ala	Ala
	530				535						540			Gln
Pro	Leu	Tyr	Ala	Gly	Tyr	Cys	Asn	His	Glu	Asn	Met			
545					550					555				

&lt;210&gt; 16

&lt;211&gt; 1857

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

```

<400> 16
ctgatccggg ccgggcgggg agtcgggtcc cgaggctccg gctcggcaga ccgggcgga 60
agcagccgag cggccatgga gctgtgcggg ctggggctgc ccgggccgcc catgctgctg 120
gcgctgctgt tggcgacact gctggcgggc atgttggcgc tgctgactca ggtggcgctg 180
gtggtgcagg tggcgaggc ggctcgggcc ccgagcgtct cggccaagcc ggggcccggc 240
ctgtggcccc tgccgctctc ggtgaagatg accccgaacc tgctgcatct cggcccggag 300
aacttctaca tcagccacag cccaattcc acggcgggcc cctcctgcac cctgctggag 360
gaagcgtttc gacgatata tggctatatt tttggtttct acaagtggca tcatgaacct 420
gctgaattcc aggctaaaac ccaggttcag caacttcttg tctcaatcac ccttcagtca 480
gagtgatgat ctttcccaaa catatcttca gatgagcttt atactttact tgtgaaagaa 540
ccagtggtctg tccttaaggc caacagagtt tggggagcat tacgaggttt agagaccttt 600
agccagttag tttatcaaga ttcttatgga actttcacca tcaatgaatc caccattatt 660
gatttctcaa ggttttctca cagaggaatt ttgattgata catccagaca ttatctgcca 720
gttaagatta ttcttaaaac tctggatgcc atggctttta ataagtttaa tgttcttcac 780
tggcacatag ttgatgacca gtctttccca tatcagagca tcacttttcc tgagttaagc 840
aataaaggaa gctattcttt gtctcatggt tatacaccaa atgatgtccg tatggtgatt 900
gaatatgcca gattacgagg aattcgagtc ctgccagaat ttgatacccc tgggcataca 960
ctatcttggg gaaaaggta gaaagacctc ctgactccat gttacagtag acaaaaacaag 1020
ttggactctt ttggacctat aaacctact ctgaatacaa catacagctt ccttactaca 1080
tttttcaaag aaattagtga ggtgtttcca gatcaattca ttcatattggg aggagatgaa 1140
gtggaattta aatgttggga atcaaatcca aaaattcaag atttcatgag gcaaaaaggc 1200
tttggcacag attttaagaa actagaatct ttctacattc aaaaggtttt ggatattatt 1260
gcaaccataa acaaggatc cattgtctgg caggaggttt ttgatgataa agcaaaagctt 1320
gcgcggggca caatagttga agtatggaaa gacagcgcat atcctgagga actcagtaga 1380
gtcacagcat ctggcttccc tgtaatcctt tctgtcctt ggtacttaga tttgattagc 1440
tatggacaag attggaggaa atactataaa gtggaacctc ttgattttgg cggtaactcag 1500
aaacagaaac aacttttcat tgggtggagaa gctgtgtctat ggggagaata tgtggatgca 1560
actaacctca ctccaagatt atggcctcgg gcaagtgtctg ttggtgagag actctggagt 1620
tccaaagatg tcagagatat ggatgacgcc tatgacagac tgacaaggca ccgctgcagg 1680
atggtcgaac gtggaatagc tgcacaacct ctttatgtct gatattgtaa ccatgagaac 1740
atgtaaaaaa tggaggggaa aaaggccaca gcaatctgta ctacaatcaa ctttattttg 1800
aaatcatgta aaataagata ttagactttt ttgaataaaa tattttttatt gattgaa 1857

```

<210> 17

<211> 536

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 17

```

Met Pro Gln Ser Pro Arg Ser Ala Pro Gly Leu Leu Leu Leu Gln Ala
1          5          10          15
Leu Val Ser Leu Val Ser Leu Ala Leu Val Ala Pro Ala Arg Leu Gln
20          25          30
Pro Ala Leu Trp Pro Phe Pro Arg Ser Val Gln Met Phe Pro Arg Leu
35          40          45
Leu Tyr Ile Ser Ala Glu Asp Phe Ser Ile Asp His Ser Pro Asn Ser
50          55          60
Thr Ala Gly Pro Ser Cys Ser Leu Leu Gln Glu Ala Phe Arg Arg Tyr
65          70          75          80
Tyr Asn Tyr Val Phe Gly Phe Tyr Lys Arg His His Gly Pro Ala Arg
85          90          95
Phe Arg Ala Glu Pro Gln Leu Gln Lys Leu Leu Val Ser Ile Thr Leu

```

100										105				110			
Glu	Ser	Glu	Cys	Glu	Ser	Phe	Pro	Ser	Leu	Ser	Ser	Asp	Glu	Thr	Tyr		
		115					120					125					
Ser	Leu	Leu	Val	Gln	Glu	Pro	Val	Ala	Val	Leu	Lys	Ala	Asn	Ser	Val		
		130				135					140						
Trp	Gly	Ala	Leu	Arg	Gly	Leu	Glu	Thr	Phe	Ser	Gln	Leu	Val	Tyr	Gln		
145					150					155					160		
Asp	Ser	Phe	Gly	Thr	Phe	Thr	Ile	Asn	Glu	Ser	Ser	Ile	Ala	Asp	Ser		
				165					170					175			
Pro	Arg	Phe	Pro	His	Arg	Gly	Ile	Leu	Ile	Asp	Thr	Ser	Arg	His	Phe		
			180					185					190				
Leu	Pro	Val	Lys	Thr	Ile	Leu	Lys	Thr	Leu	Asp	Ala	Met	Ala	Phe	Asn		
		195					200					205					
Lys	Phe	Asn	Val	Leu	His	Trp	His	Ile	Val	Asp	Asp	Gln	Ser	Phe	Pro		
	210					215				220							
Tyr	Gln	Ser	Thr	Thr	Phe	Pro	Glu	Leu	Ser	Asn	Lys	Gly	Ser	Tyr	Ser		
225					230					235				240			
Leu	Ser	His	Val	Tyr	Thr	Pro	Asn	Asp	Val	Arg	Met	Val	Leu	Glu	Tyr		
				245					250					255			
Ala	Arg	Leu	Arg	Gly	Ile	Arg	Val	Ile	Pro	Glu	Phe	Asp	Thr	Pro	Gly		
			260					265					270				
His	Thr	Gln	Ser	Trp	Gly	Lys	Gly	Gln	Lys	Asn	Leu	Leu	Thr	Pro	Cys		
		275					280					285					
Tyr	Asn	Gln	Lys	Thr	Lys	Thr	Gln	Val	Phe	Gly	Pro	Val	Asp	Pro	Thr		
	290					295				300							
Val	Asn	Thr	Thr	Tyr	Ala	Phe	Phe	Asn	Thr	Phe	Phe	Lys	Glu	Ile	Ser		
305					310					315					320		
Ser	Val	Phe	Pro	Asp	Gln	Phe	Ile	His	Leu	Gly	Gly	Asp	Glu	Val	Glu		
			325						330					335			
Phe	Gln	Cys	Trp	Ala	Ser	Asn	Pro	Asn	Ile	Gln	Gly	Phe	Met	Lys	Arg		
			340					345					350				
Lys	Gly	Phe	Gly	Ser	Asp	Phe	Arg	Arg	Leu	Glu	Ser	Phe	Tyr	Ile	Lys		
		355					360					365					
Lys	Ile	Leu	Glu	Ile	Ile	Ser	Ser	Leu	Lys	Lys	Asn	Ser	Ile	Val	Trp		
	370					375					380						
Gln	Glu	Val	Phe	Asp	Asp	Lys	Val	Glu	Leu	Gln	Pro	Gly	Thr	Val	Val		
385					390					395					400		
Glu	Val	Trp	Lys	Ser	Glu	His	Tyr	Ser	Tyr	Glu	Leu	Lys	Gln	Val	Thr		
				405					410					415			
Gly	Ser	Gly	Phe	Pro	Ala	Ile	Leu	Ser	Ala	Pro	Trp	Tyr	Leu	Asp	Leu		
			420					425					430				
Ile	Ser	Tyr	Gly	Gln	Asp	Trp	Lys	Asn	Tyr	Tyr	Lys	Val	Glu	Pro	Leu		
		435					440					445					

<210> 18

<211> 1750

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 18

ggagcagtc	tgccgcagtc	cccgcgtagc	gcccccgggc	tgctgctgct	gcaggcgctg	60
gtgtcgctag	tgctcgctggc	cctagtggcc	ccggcccgac	tgcaacctgc	gctatggccc	120
ttcccgcgt	cgtgagcat	gttcccgcgg	ctgttgtaga	tctccgcgga	ggacttcagc	180
atcgaccaca	gtcccaattc	cacagcgggc	ccttctctgct	cgctgctaca	ggaggcggtt	240
cggcgatatt	acaactatgt	ttttggtttc	tacaagagac	atcatggccc	tgctagattt	300
cgagctgagc	cacagttgca	gaagtcctg	gtctccatta	ccctcgagtc	agagtgcgag	360
tccttcccta	gtctgtcttc	agatgaaacc	tattctctgc	ttgtacaaga	accagtagcc	420
gtcctcaagg	ccaacagcgt	ttggggagcg	ttacgaggtt	tagagacgtt	tagccagtta	480
gtttaccaag	actctttcgg	gactttcacc	atcaatgaat	ccagtatagc	tgattctcca	540
agattccctc	atagaggat	tttaattgat	acatctagac	acttctctgc	tgtgaagaca	600
attttaaaaa	ctctggatgc	catggctttt	aataagttta	atgttcttca	ctggcacata	660
gtggacgacc	agtctttccc	ttatcagagt	accacttttc	ctgagctaag	caataaggga	720
agctactctt	tgtctcatgt	ctatacacca	aacgatgtcc	ggatgggtgt	ggagtacgcc	780
cggtctccgag	ggattcgagt	cataccagaa	tttgataccc	ctggccatac	acagtcttgg	840
ggcaaaggac	agaaaaacct	tctaactcca	tggtacaatc	aaaaaactaa	aactcaagtg	900
tttgggcctg	tagaccaaac	tgtaaacaca	acgtatgcat	tctttaacac	atttttcaaa	960
gaaatcagca	gtgtgtttcc	agatcagttc	atccacttgg	gaggagatga	agtagaattt	1020
caatgttggg	catcaaattc	aaacatccaa	ggtttcatga	agagaaaggg	ctttggcagc	1080
gatttttagaa	gactagaatc	cttttatatt	aaaaagattt	tggaaattat	ttcatcctta	1140
aagaagaact	ccattgtttg	gcaagaagtt	tttgatgata	aggtggagct	tcagccgggc	1200
acagtagtcg	aagtgtggaa	gagtgcagat	tattcatatg	agctaaagca	agtcacaggc	1260
tctggcttcc	ctgccatcct	ttctgctcct	tggtacttag	acctgatcag	ctatgggcaa	1320
gactggaaaa	actactacaa	agttgagccc	cttaattttg	aaggctctga	gaagcagaaa	1380
caacttgta	ttggtggaga	agcttgccctg	tggggagaat	ttgtggatgc	aactaacctt	1440
actccaagat	tatggcctcg	agcaagcgct	gttgggtgaga	gactctggag	ccctaaaact	1500
gtcactgacc	tagaaaatgc	ctacaaacga	ctggccgtgc	accgctgcag	aatggctcagc	1560
cgtggaatag	ctgcacaacc	tctctatact	ggatactgta	actatgagaa	taaaatatag	1620
aagtgcagca	cgtctacagc	attccagcta	tgatcatggt	gattctgaaa	tcagttaaat	1680
taagatttgt	taggctgttt	tttttttaaa	taaaccatct	ttttattgat	tgaatctttc	1740
taaaaaaaaa						1750

&lt;210&gt; 19

&lt;211&gt; 12263

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 19

aatgtagtct	tatgcaatac	tcttgtagtc	ttgcaacatg	gtaacgatga	gttagcaaca	60
tgcccttaca	ggagagaaaa	agcaccgtgc	atgccgattg	gtggaagtaa	ggtggtacga	120
tcgtgcctta	ttaggaaggc	aacagacggg	tctgacatgg	attggacgaa	ccactgaatt	180
gccgcattgc	agagatattg	tatttaagtg	cctagctcga	tacataaacg	ggtctctctg	240
gttagaccag	atctgagcct	gggagctctc	tggttaacta	gggaaccac	tgcttaagcc	300
tcaataaagc	ttgccttgag	tgcttcaagt	agtgtgtgcc	cgtctgttgt	gtgactctgg	360
taactagaga	tccctcagac	ccttttagtc	agtgtggaaa	atctctagca	gtggcgccc	420
aacagggact	tgaaagcgaa	agggaaacca	gaggagctct	ctcgacgcag	gactcggtt	480
gctgaagcgc	gcacggcaag	aggcgagggg	cggcgactgg	tgagtacgcc	aaaaattttg	540
actagcggag	gctagaagga	gagagatggg	tgcgagagcg	tcagtattaa	gcgggggaga	600

attagatcgc	gatgggaaaa	aattcgggta	aggccagggg	gaaagaaaaa	atataaatta	660
aaacatatag	tatgggcaag	cagggagcta	gaacgattcg	cagttaatcc	tggcctgtta	720
gaaacatcag	aaggctgtag	acaaatactg	ggacagctac	aaccatccct	tcagacagga	780
tcagaagaac	ttagatcatt	atataatata	gtagcaaccc	tctatttgtt	gcatcaaagg	840
atagagataa	aagacaccaa	ggaagcttta	gacaagatag	aggaagagca	aaacaaaagt	900
aagaccaccg	cacagcaagc	ggccgctgat	cttcagacct	ggaggaggag	atatgaggga	960
caattggaga	agtgaattat	ataaatataa	agtagtaaaa	attgaaccat	taggagtagc	1020
accaccaaag	gcaaagagaa	gagtgggtgca	gagagaaaaa	agagcagtgg	gaataggagc	1080
tttgttcctt	gggttccttg	gagcagcagg	aagcactatg	ggcgagcgt	caatgacgct	1140
gacggtacag	gccagacaat	tattgtctgg	tatagtgcag	cagcagaaca	atttgcgtgag	1200
ggctattgag	gcgcaacagc	atctgttgca	actcacagtc	tggggcatca	agcagctcca	1260
ggcaagaatc	ctggctgtgg	aaagatacct	aaaggatcaa	cagctcctgg	ggatttgggg	1320
ttgctctgga	aaactcattt	gcaccactgc	tgtgccttgg	aatgctagt	ggagtaataa	1380
atctctggaa	cagattttgga	atcacacgac	cttgatggag	tgggacagag	aaattaacaa	1440
ttacacaagc	ttaatacact	ccttaattga	agaatcgcaa	aaccagcaag	aaaagaatga	1500
acaagaatta	ttggaattag	ataaatgggc	aagtttgtgg	aattggttta	acataacaaa	1560
ttggctgtgg	tatataaaat	tattcataat	gatagtagga	ggcttggtag	gtttaagaat	1620
agtttttgc	gtactttcta	tagtgaatag	agttaggcag	ggatattcac	cattatcggt	1680
tcagaccac	ctcccaaccc	cgaggggacc	cgacaggccc	gaaggaatag	aagaagaagg	1740
tggagagaga	gacagagaca	gatccattcg	attagtgaac	ggatctcgac	ggatcagata	1800
agcttgatat	cgaattcggg	accctagtta	tttaatagtaa	tcaattacgg	ggcatttagt	1860
tcatagccca	tatatggagt	tccgcgttac	ataacttacg	gtaaatggcc	cgctggctg	1920
accgcccaac	gacccccgcc	cattgacgtc	aataatgacg	tatgttccca	tagtaacgcc	1980
aatagggact	ttccattgac	gtcaatgggt	ggactattta	cggtaaactg	cccacttggc	2040
agtacatcaa	gtgtatcata	tgccaagtac	gccccctatt	gacgtcaatg	acggtaaatg	2100
gcccgcctgg	cattatgccc	agtacatgac	cttatgggac	tttccctact	ggcagtacat	2160
ctaogtatta	gtcatcgcta	ttaccatggt	cgaggtgagc	cccacgttct	gcttcactct	2220
ccccactccc	ccccctccc	cacccccaat	tttgatttta	tttatttttt	aattattttg	2280
tgcagcgatg	ggggcggggg	gggggggggg	gcgcgcgcca	ggcgggggcg	ggcggggcga	2340
ggggcggggc	ggggcgaggc	ggagaggtgc	ggcggcagcc	aatcagagcg	gcgcgctccg	2400
aaagtttctt	tttatggcga	ggcgggcgcg	gcggcgggcc	tataaaaagc	gaagcgcgcg	2460
gcggggcgga	gtcgctgcga	cgctgccttc	gccccgtgcc	ccgctccgcc	gccgctcgc	2520
gcccggcgcc	ccggtctgta	ctgaccgcgt	tactcccaca	ggtgagcggg	cgggacggcc	2580
cttctcctcc	gggctgtaat	tagcgcttgg	tttaatgacg	gcttgtttct	tttctgtggc	2640
tgcgtgaaag	ccttgagggg	ctccgggagg	gccccttgtg	cgggggggag	cggtcgggg	2700
ggtgcgtgcg	tgtgtgtgtg	cgtggggagc	gcgcgctgcg	gcccgcgctg	cccgccggct	2760
gtgagcgctg	cgggcgcggc	gcggggcttt	gtgcgctccg	cagtgtgcgc	gaggggagcg	2820
cgccgggggg	cggtgccccg	cggtgcgggg	ggggctgcga	ggggaacaaa	ggctgcgtgc	2880
ggggtgtgtg	cgtggggggg	tgagcagggg	gtgtgggcgc	ggcggtcggg	ctgtaacccc	2940
cccctgcacc	ccctccccc	agttgctgag	cacggccccg	cttcgggtgc	ggggctccgt	3000
acggggcggtg	gcgcgggggt	cgccgtgccc	ggcggggggt	ggcggcagg	gggggtgccg	3060
ggcgggggcg	ggccgcctcg	ggccggggag	ggctcggggg	aggggcgcgg	cgcccccg	3120
agcgccggcg	gctgtcgagg	cgcggcgagc	cgagccatt	gccttttatg	gtaatcgtgc	3180
gagagggcg	agggaacttc	tttgtcccaa	atctgtgcgg	agccgaaatc	tgggagggcg	3240
cgccgcaccc	cctctagcgg	gcgcggggcg	aagcgggtgc	gcgcgggcag	gaaggaaatg	3300
ggcggggagg	gccttcgtgc	gtcgccgcgc	cgccgtcccc	ttctccctct	ccagcctcgg	3360
ggctgtccgc	ggggggacgg	ctgccttcgg	gggggacggg	gcagggcggg	gttcggcttc	3420
tggcgtgtga	ccggcggtc	tagagcctct	gctaaccatg	ttcatgcctt	cttctttttc	3480
ctacagctcc	tgggcaacgt	gctggttatt	gtgctgtctc	atcatttttg	caaagaattc	3540
ctgcagcccc	gggatccac	tagtccagt	tgggtgaatt	gatcccttca	cctaatacga	3600
ctcactatag	gctagcctcg	agctgatccg	ggccgggcgg	gaagtgcggg	cccagggtc	3660
cggtcggca	gaccgggcgg	aaagcagccg	agcgcccatg	gagctgtgcg	ggctggggct	3720
gccccggccg	cccatgctgc	tggcgtgct	gttggcgaca	ctgctggcgg	cgatgttggc	3780
gctgctgact	caggtggcgc	tgggtggtgca	ggtggcgagg	gcggctcggg	ccccgagcgt	3840
ctcggccaag	ccggggccgg	cgctgtggcc	cctgccgctc	tcggtgaaga	tgacccccgaa	3900
cctgctgcat	ctcgccccgg	agaacttcta	catcagccac	agccccaatt	ccacggcggg	3960
ccccctctgc	accctgctgg	aggaagcgtt	ccagcatat	catggctata	tttttggttt	4020
ctacaagtgg	catcatgaac	ctgctgaatt	ccaggctaaa	accaggttc	agcaacttct	4080
tgtctcaatc	acccttcagt	cagagtgtga	tgctttcccc	aacatatctt	cagatgagtc	4140

ttatacttta	cttgtgaaag	aaccagtggc	tgtccttaag	gccaacagag	tttggggagc	4200
attacgaggt	ttagagacct	ttagccagtt	agtttatcaa	gattccttatg	gaactttcac	4260
catcaatgaa	tccaccatta	ttgattctcc	aaggttttct	cacagaggaa	ttttgattga	4320
tacatccaga	cattatctgc	cagttaagat	tattcttaaa	actctggatg	ccatggcctt	4380
taataagttt	aatgttcttc	actggcacat	agttgatgac	cagtctttcc	catatcagag	4440
catcactttt	cctgagttaa	gcaataaagg	aagctattct	ttgtctcatg	tttatacacc	4500
aaatgatgtc	cgtatggtga	ttgaatatgc	cagattacga	ggaattcgag	tcctgccaga	4560
atlttgatacc	cctgggcata	cactatcttg	gggaaaagg	cagaaagacc	tcctgactcc	4620
atgttacagt	agacaaaaca	agttggactc	ttttggacct	ataaacctta	ctctgaatac	4680
aacatacagc	ttccttacta	catttttcaa	agaaattagt	gaggtgtttc	cagatcaatt	4740
cattcatttg	ggaggagatg	aagtgaatt	taaatgttg	gaatcaaata	caaaaattca	4800
agatttcatg	aggcaaaaag	gctttggcac	agattttaag	aaactagaat	ctttctacat	4860
tcaaaagggt	ttggatatta	ttgcaacat	aaacaaggga	tccattgtct	ggcaggagg	4920
ttttgatgat	aaagcaaagc	ttgcgccggg	cacaatagtt	gaagtatgga	aagacagcgc	4980
atatacctgag	gaactcagta	gagtcacagc	atctggcctc	cctgtaatac	tttctgctcc	5040
ttggtaactta	gatttgatta	gctatggaca	agattggagg	aaatactata	aagtggaaac	5100
tcttgatttt	ggcggtaactc	agaaacagaa	acaacttttc	attggtggag	aagcttgtct	5160
atggggagaa	tatgtggatg	caactaacct	cactccaaga	ttatggcctc	gggcaagtgc	5220
tgttggtgag	agactctgga	gttccaaaga	tgtcagagat	atggatgacg	cctatgacag	5280
actgacaagg	caccgctgca	ggatggctga	acgtggaata	gctgcacaac	ctctttatgc	5340
tggatattgt	aaccatgaga	acatgtaaaa	aatggagggg	aaaaaggcca	cagcaatctg	5400
tactacaatc	aactttattt	tgaatcatg	taaaataaga	tattagactt	ttttgaataa	5460
actcgagaat	tcacgcgtcg	agcatgcata	tagggcgggc	aattccgccc	ctctccctcc	5520
ccccccccta	acgttactgg	ccgaagccgc	ttggaataag	gccggtgtgc	gtttgtctat	5580
atgtgatttt	ccaccatatt	gccgtctttt	ggcaatgtga	gggcccgga	acctggccct	5640
gtcttcttga	cgagcattcc	taggggtctt	tcccctctcg	ccaaagggaat	gcaagggtctg	5700
ttgaatgtcg	tgaagggaagc	agttcctctg	gaagcttctt	gaagacaaac	aacgtctgta	5760
gcgacccttt	gcaggcagcg	gaacccccca	cctggcgaca	ggtgcctctg	cggccaaaag	5820
ccacgtgtat	aagatacacc	tgcaaaggcg	gcacaacccc	agtgccacgt	tgtgagttgg	5880
atagttgtgg	aaagagtcaa	atggctctcc	tcaagcgtat	tcaacaagg	gctgaaggat	5940
gcccagaagg	tacccattg	tatgggatct	gatctggggc	ctcgggtgcac	atgctttaca	6000
tgtgtttagt	cgaggttaaa	aaaacgtcta	ggccccccga	accacgggga	cgtggttttc	6060
ctttgaaaaa	cacgatgata	agcttgccac	aaccgggat	cctctatgac	aagctccagg	6120
ctttgggttt	cgtctgtgct	ggcggcagcg	ttcgaggac	gggcgacggc	cctctggccc	6180
tggcctcaga	acttccaaac	ctccgaccag	cgctacgtcc	tttaccgaa	caactttcaa	6240
ttccagtacg	atgtcagctc	ggcgcgcag	cccgctgct	cagtcctcga	cgaggccttc	6300
cagcgctatc	gtgacctgct	tttcggttcc	gggtcttggc	cccgctctta	cctcacagg	6360
aaacggcata	cactggagaa	gaatgtgttg	gttgtctctg	tagtcacacc	tggatgtaac	6420
cagcttccta	ctttggagtc	agtggagaat	tataccctga	ccataaatga	tgaccagtgt	6480
ttactcctct	ctgagactgt	ctggggagct	ctccgaggtc	tggagacttt	tagccagctt	6540
gtttggaaat	ctgctgagg	cacattcttt	atcaacaaga	ctgagattga	ggactttccc	6600
cgttttcctc	accggggctt	gctgttggt	acatctcgcc	attacctgcc	actctctagc	6660
atcctggaca	ctctggatgt	catggcgta	aataaattga	acgtgttcca	ctggcatctg	6720
gtagatgata	cttccttccc	atatgagagc	ttcacttttc	cagagctcat	gagaaagggg	6780
tcctacaacc	ctgtcaccca	catctacaca	gcacaggatg	tgaaggagg	cattgaatac	6840
gcacggctcc	ggggtatccg	tgtgcttgca	gagtttgaca	ctcctggcca	cactttgtcc	6900
tggggaccag	gtatccctgg	attactgact	ccttgctact	ctgggtctga	gccctctggc	6960
acctttggac	cagtgaatcc	cagtctcaat	aatacctatg	agttcatgag	cacattcttc	7020
ttagaagtca	gctctgtctt	cccagatttt	tatcttcata	ttggaggaga	tgaggttgat	7080
ttcacctgct	ggaagtccaa	cccagagatc	caggacttta	tgaggagaa	aggcttcggt	7140
gaggacttca	agcagctgga	gtccttctac	atccagagc	tgtctggacat	cgtctcttct	7200
tatggcaagg	gctatgtggt	gtggcaggag	gtgtttgata	ataaagtaaa	gattcagcca	7260
gacacaatca	tacaggtgtg	gcgagaggat	attccagtga	actatatgaa	ggagctggaa	7320
ctggctacca	aggccggctt	ccgggccctt	ctctctgccc	cctggtacct	gaaccgtata	7380
tcctatggcc	ctgactggaa	ggatttctac	gtagtggaa	ccctggcatt	tgaaggtaac	7440
cctgagcaga	aggctctggt	gattgggtgga	gaggcttgta	tgtggggaga	atatgtggac	7500
aacacaaacc	tggctcccag	gctctggccc	agagcagggg	ctgttgccga	aaggctgtgg	7560
agcaacaagt	tgacatctga	cctgacattt	gcctatgaac	gtttgtcaca	cttccgctgt	7620
gagttgctga	ggcgagggtg	ccaggcccaa	cccctcaatg	taggcttctg	tgagcaggag	7680

tttgaacaga	cctgaagagt	cgaccggggc	ggccgcttcc	ctttagttag	ggttaatgaa	7740
gggctcgagt	ctagagggcc	cgcggttcga	aggtaagcct	atccctaacc	ctctcctcgg	7800
tctcgattct	acgcgtaccg	gttagtaatg	agtttggaat	taattctgtg	gaatgtgtgt	7860
cagttagggt	gtggaaagtc	cccaggctcc	ccaggcaggc	agaagtatgc	aaagcatgca	7920
tctcaattag	tcagcaacca	ggtgtggaaa	gtccccaggc	tccccagcag	gcagaagtat	7980
gcaaagcatg	catctcaatt	agtcagcaac	catagtcccc	cccctaactc	cgcccatccc	8040
gcccctaact	ccgcccagtt	ccgcccattc	tccgccccat	ggctgactaa	ttttttttat	8100
ttatgcagag	gccgaggccg	cctctgcctc	tgagctattc	cagaagtagt	gaggaggcct	8160
ttttggaggc	ctaggctttt	gcaaaaagct	cccgggagct	tgtatatcca	ttttcggatc	8220
tgatcagcac	gtggttgaaa	ttaatcatcg	gcatagtata	tcggcatagt	ataatacgac	8280
aaggtgagga	actaaacat	ggccaagcct	ttgtctcaag	aagaatccac	cctcattgaa	8340
agagcaacgg	ctacaatcaa	cagcatcccc	atctctgaag	actacagcgt	cgccagcgca	8400
gctctctcta	gcgacggccg	catcttcaat	ggtgtcaatg	tatatcattt	tactggggga	8460
ccttgtgcag	aactcgtggt	gctgggcact	gctgtgctg	cggcagctgg	caacctgact	8520
tgtatcgtcg	cgatcggaaa	tgagaacagg	ggcacttga	gccctgcgg	acggtgccga	8580
caggtgcttc	tcgatctgca	tcctgggatc	aaagccatag	tgaaggacag	tgatggacag	8640
ccgacggcag	ttgggattcg	tgaattgctg	ccctctggtt	atgtgtggga	gggctaagca	8700
caattcgagc	tcgggtacctt	taagaccaat	gacttacaag	gcagctgtag	atcttagcca	8760
ctttttaaaa	gaaaaggggg	gactggaagg	gctaattcac	tcccaacgaa	gacaagatct	8820
gctttttgct	tgtactgggt	ctctctggtt	agaccagatc	tgagcctggg	agctctctgg	8880
ctaactaggg	aacccactgc	ttaagcctca	ataaagcttg	ccttgagtgc	ttcaagtagt	8940
gtgtgcccgt	ctgttgtgtg	actctggtaa	ctagagatcc	ctcagaccct	tttagtcagt	9000
gtggaaaatc	tctagcagta	gtagttcatg	tcactttatt	attcagtatt	tataacttgc	9060
aaagaaatga	atatcagaga	gtgagaggaa	cttgtttatt	gcagcttata	atggttacia	9120
ataaagcaat	agcatcacia	atttcacaaa	taaagcattt	ttttcactgc	attctagtgt	9180
tggtttgtcc	aaactcatca	atgtatctta	tcagtgtctg	ctctagctat	cccgccccta	9240
actccgcccc	tcccgcccct	aactccgccc	agttccgccc	attctccgcc	ccatggctga	9300
ctaatttttt	ttattttatgc	agaggccgag	gccgcctcgg	cctctgagct	attccagaag	9360
tagtgaggag	gctttttttg	aggcctaggg	acgtacccaa	tccgcccctat	agtgagtcgt	9420
attacgcgcg	ctcactggcc	gtcgttttac	aacgtctgga	ctgggaaaac	cctggcggtta	9480
cccaacttaa	tcgccttgca	gcacatcccc	ctttcgccag	ctggcgtaat	agcgaagagg	9540
cccgaccgga	tcgccccttc	caacagttgc	gcagcctgaa	tggcgaatgg	gacgcgccct	9600
gtagcggcgc	attaagcgcg	gcgggtgtgg	tggttacgcg	cagcgtgacc	gctacacttg	9660
ccagcgccct	agcgcccgct	cctttcgctt	tcttcccttc	ctttctcgcc	acgttcgcgg	9720
gctttccccc	tcaagctcta	aatcgggggc	tccctttagg	gttccgattt	agtgttttac	9780
ggcacctcga	ccccaaaaaa	cttgattagg	gtgatggttc	acgtagtggg	ccatcgccct	9840
gataagcggg	ttttcgccct	ttgacgttgg	agtcacagtt	ctttaatagt	ggactcttgt	9900
tccaaactgg	aacaacactc	aaccctatct	cggctctattc	ttttgattta	taagggtattt	9960
tgccgatttc	ggcctatttg	ttaaaaaatg	agctgattta	acaaaaattt	aacgcgaatt	10020
ttaacaaaat	attaacgctt	acaatttagg	tggcactttt	cggggaaatg	tgcgcggaac	10080
ccctattttg	ttatttttct	aaatacattc	aaatatgtat	ccgctcatga	gacaataacc	10140
ctgataaatg	cttcaataat	attgaaaaag	gaagagtatg	agtattcaac	atttccgtgt	10200
cgcccttatt	cccttttttg	cggcattttg	ccttctctgtt	tttgtctacc	cagaaacgct	10260
ggtgaaagta	aaagatgctg	aagatcagtt	gggtgcacga	gtgggttaca	tcgaactgga	10320
tctcaacagc	ggtaagatcc	ttgagagttt	tccgcccga	gaacgttttc	caatgatgag	10380
cactttttaa	gttctgctat	gtggcgcggt	attatcccg	attgacgcgg	ggcaagagca	10440
actcggctcg	cgcatacact	attctcagaa	tgacttggtt	gagtactcac	cagtcacaga	10500
aaagcatctt	acggatggca	tgacagtaag	agaattatgc	agtgtgcca	taaccatgag	10560
tgataaacact	gcggccaact	tacttctgac	aacgatcgga	ggaccgaagg	agctaaccgc	10620
ttttttgcac	aacatggggg	atcatgtaac	tgccttgat	cgttgggaac	cggagctgaa	10680
tgaagccata	ccaaacgacg	agcgtgacac	cacgatgcct	gtagcaatgg	caacaacggt	10740
gcgcaaacta	ttactggcg	aactacttac	tctagcttcc	cggcaacaat	taatagactg	10800
gatggaggcg	gataaagttg	caggaccact	tctgcgctcg	gcccttccgg	ctggctgggt	10860
tattgctgat	aaatctggag	ccggtgagcg	tgggtctcgc	ggtatcattg	cagcactggg	10920
gccagatggt	aagccctccc	gtatcgtagt	tatctacacg	acggggagtc	aggcaactat	10980
ggatgaacga	aatagacaga	tcgctgagat	agggtcctca	ctgattaagc	attggtaact	11040
gtcagaccaa	gtttactcat	atatacttta	gattgattta	aaacttcatt	tttaatttaa	11100
aaggatctag	gtgaagatcc	tttttgataa	tctcatgacc	aaaatccctt	aacgtgagtt	11160
ttcgttccac	tgagcgtcag	accccgtaga	aaagatcaaa	ggatcttctt	gagatccttt	11220

ttttctgcgc	gtaatctgct	gcttgcaaac	aaaaaaacca	ccgctaccag	cggtgggttg	11280
tttgccggat	caagagctac	caactctttt	tccgaaggta	actggcttca	gcagagcgca	11340
gataccaaat	actgttcttc	tagtgtagcc	gtagttaggc	caccacttca	agaactctgt	11400
agcaccgcct	acatacctcg	ctctgctaata	cctgttacca	gtggctgctg	ccagtggcga	11460
taagtcgtgt	cttaccgggt	tggactcaag	acgatagtta	ccggataagg	cgcagcggtc	11520
gggctgaacg	gggggttcgt	gcacacagcc	cagcttggag	cgaacgacct	acaccgaact	11580
gagataccta	cagcgtgagc	tatgagaaaag	cgccacgctt	cccgaaggga	gaaaggcgga	11640
cagggtatccg	gtaagcggca	gggtcggaac	aggagagcgc	acgagggagc	ttccaggggg	11700
aaacgcctgg	tatctttata	gtcctgtcgg	gtttcgccac	ctctgacttg	agcgtcgatt	11760
tttgtgatgc	tcgtcagggg	ggcggagcct	atggaaaaac	gccagcaacg	cggccttttt	11820
acggttcctg	gccttttgct	ggccttttgc	tcacatgttc	tttctctcgt	tatcccctga	11880
ttctgtggat	aaccgtatta	ccgcctttga	gtgagctgat	accgctcgcc	gcagccgaac	11940
gaccgagcgc	agcgagtcag	tgagcgagga	agcggaagag	cgccaataac	gcaaaccgcc	12000
tctccccgcg	cgttggccga	ttcattaatg	cagctggcac	gacaggtttc	ccgactggaa	12060
agcgggcagt	gagcgcaacg	caattaatgt	gagttagctc	actcattagg	caccccaggc	12120
tttacctttt	atgcttccgg	ctcgtatgtt	gtgtggaatt	gtgagcggat	aacaatttca	12180
cacaggaaac	agctatgacc	atgattacgc	caagcgcgca	attaaccctc	actaaaggga	12240
acaaaagctg	gagctgcaag	ctt				12263

&lt;210&gt; 20

&lt;211&gt; 11110

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 20

aatgtagtct	tatgcaatac	tcttgtagtc	ttgcaacatg	gtaacgatga	gttagcaaca	60
tgccttacaa	ggagagaaaa	agcaccgtgc	atgccgattg	gtggaagtaa	ggtggtacga	120
tcgtgcctta	ttaggaaggc	aacagacggg	tctgacattg	attggacgaa	ccactgaatt	180
gccgcattgc	agagatattg	tattttaagt	cctagctcga	tacataaacg	ggtctctctg	240
gttagaccag	atctgagcct	gggagctctc	tggctaacta	gggaaccac	tgcttaagcc	300
tcaataaagc	ttgccttgag	tgcttcaagt	agtgtgtgcc	cgtctgttgt	gtgactctgg	360
taactagaga	tccttcagac	ccttttagtc	agtgtggaaa	atctctagca	gtggcgcccg	420
aacagggact	tgaaagcgaa	agggaaacca	gaggagctct	ctcgacgcag	gactcggctt	480
gctgaagcgc	gcacggcaag	aggcgagggg	cggcgactgg	tgagtacgcc	aaaaattttg	540
actagcggag	gctagaagga	gagagatggg	tgcgagagcg	tcagtattaa	gcgggggaga	600
attagatcgc	gatgggaaaa	aattcggtta	aggccagggg	gaaagaaaaa	atataaatta	660
aaacatatag	tatgggcaag	caggagccta	gaacgattcg	cagttaatcc	tggcctgtta	720
gaaacatcag	aaggctgtag	acaaatactg	ggacagctac	aaccatccct	tcagacagga	780
tcagaagaac	ttagatcatt	atataataca	gtagcaaccc	tctatttgtg	gcatcaaagg	840
atagagataa	aagacaccaa	ggaagcttta	gacaagatag	aggaagagca	aaacaaaagt	900
aagaccaccg	cacagcaagc	ggccgctgat	cttcagacct	ggaggaggag	atatgaggga	960
caattggaga	agtgaattat	ataaatataa	agtagtaaaa	attgaaccat	taggagtagc	1020
acccaccaag	gcaaagagaa	gagtgggtgca	gagagaaaaa	agagcagtgg	gaataggagc	1080
tttgttcctt	gggttcttgg	gagcagcagg	caagcactatg	ggcgacgcgt	caatgacgct	1140
gacggtacag	gccagacaat	tattgtctgg	tatagtgcag	cagcagaaca	atttgctgag	1200
ggctattgag	gcgcaacagc	atctgttgca	actcacagtc	tggggcatca	agcagctcca	1260
ggcaagaatc	ctggctgtgg	aaagatacct	aaaggatcaa	cagctcctgg	ggatttgggg	1320
ttgctctgga	aaactcattt	gcaccactgc	tgtgccttgg	aatgctagtt	ggagtaataa	1380
atctctggaa	cagatttgga	atcacacgac	ctggatggag	tgggacagag	aaattaacaa	1440
ttacacaagc	ttaatacact	ccttaattga	agaatcgcaa	aaccagcaag	aaaagaatga	1500
acaagaatta	ttggaattag	ataaatgggc	aagtttgtgg	aatttggtta	acataacaaa	1560
ttggctgtgg	tatataaaat	tattcataat	gatatagtaga	ggcttggtag	gtttaagaat	1620



agtttttgc	gtactttcta	tagtgaatag	agttaggcag	ggatattcac	cattatcggt	1680
tcagaccac	ctcccaaccc	cgaggggacc	cgacaggccc	gaaggaatag	aagaagaagg	1740
tgagagaga	gacagagaca	gatccattcg	attagtgaac	ggatctcgac	ggtatcgata	1800
agcttgggag	ttccgcgtta	cataacttac	ggtaaattggc	ccgcctggct	gaccgcccac	1860
cgacccccgc	ccattgacgt	caataatgac	gtatgttccc	atagtaacgc	caatagggac	1920
tttccattga	cgatcaatggg	tggagtattt	acggtaaaact	gcccacttgg	cagtacatca	1980
agtgtatcat	atgccaagta	cgccccctat	tgacgtcaat	gacggtaaat	ggccccgctg	2040
gcattatgcc	cagtacatga	ccttatggga	ctttcctact	tggcagtaca	tctacgtatt	2100
agtcacgcgt	attaccatgg	tgatgcgggt	ttggcagtac	atcaatgggc	gtggatagcg	2160
gtttgactca	cggggatttc	caagtctcca	ccccattgac	gtcaatggga	gtttgttttg	2220
gcacaaaat	caacgggact	ttccaaaatg	tcgtaacaac	tccgccccat	tgacgcaa	2280
gggcggtagg	cgtgtacggg	gggaggtcta	tataagcaga	gctcgtttag	tgaaccgtca	2340
gatcgccctg	agacgccatc	cacgctgttt	tgacctccat	agaagacacc	gactctagag	2400
gatccactag	tccagtgtgg	tggaaattgat	cccttcacct	aatacgactc	actataggct	2460
agcctcgagc	tgatccgggc	cgggcgggaa	gtcgggtccc	gaggctccgg	ctcggcagac	2520
cgggcggaaa	gcagccgagc	ggccatggag	ctgtgcgggc	tggggctgcc	ccggccgccc	2580
atgctgctgg	cgctgctgtt	ggcgacactg	ctggcggcga	tgttggcgct	gctgactcag	2640
gtggcgctgg	tgggtcagggt	ggcgaggcg	gctcgggccc	cgagcgcttc	ggccaagccg	2700
gggcccggcg	tgtggccctc	gccgctctcg	gtgaagatga	ccccgaacct	gctgcatctc	2760
gccccggaga	acttctacat	cagccacagc	cccaattcca	cggggggccc	ctcctgcacc	2820
ctgctggagg	aagcgtttcg	acgatatcat	ggctatatatt	ttggtttcta	caagtggcat	2880
catgaacctg	ctgaattcca	ggctaaaacc	caggttccagc	aacttcttgt	ctcaatcacc	2940
cttcagtcag	agtgtgatgc	tttccccaac	atatcttcag	atgagcttta	tactttactt	3000
gtgaaagaac	cagtggctgt	ccttaaggcc	aacagagttt	ggggagcatt	acgagggtta	3060
gagaccttta	gccagttagt	ttatcaagat	tcttatggaa	ctttcaccat	caatgaatcc	3120
accattattg	attctccaag	gttttctcac	agaggaattt	tgattgatac	atccagacat	3180
tatctgccag	ttaagattat	tcttaaaact	ctggatgcca	tggcttttaa	taagttta	3240
gttcttctact	ggcacatagt	tgatgaccag	tctttcccat	atcagagcat	cacttttcc	3300
gagttaaagc	ataaaggaag	ctattctttg	ctctatgttt	atacaccaa	tgatgtccgt	3360
atggtgatag	aatatggcag	attcagatcc	tgccagaatt	tgactccatg	ttacagtaga	3420
gggcatacac	tatcttgggg	aaaaggtcag	aaagacctcc	tgaatacaac	atacagcttc	3480
caaaacaagt	tggactcttt	tggacctata	aacctactc	atcaattcat	tcatttggga	3540
cttactacat	ttttcaaaga	aattagttag	gtgtttccag	aaattcaaga	tttcatgagg	3600
ggagatgaag	tggaaatttaa	atgttgggaa	tcaaattcaa	tctacattca	aaaggttttg	3660
caaaaaggct	ttggcacaga	ttttaagaaa	ctagaatctt	aggagggttt	tgatgataaa	3720
gatattattg	caaccataaa	caagggatcc	attgtctggc	acagcgcata	tcttgaggaa	3780
gcaaaagctg	cgccgggcac	aatagttgaa	gtatggaaag	ctgctccttg	gtacttagat	3840
ctcagtagag	tcacagcatc	tggcttccct	gtaatccttt	tggaaacctct	tgattttggc	3900
ttgattagct	atggacaaga	ttggaggaaa	tactataaag	cttgtctatg	gggagaatat	3960
ggtactcaga	aacagaaaca	acttttcatt	ggtggagaag	caagtgtctg	tggtgagaga	4020
gtggatgcaa	ctaacctcac	tccaagatta	tggcctcggg	atgacagact	gacaaggcac	4080
ctctggagtt	ccaaaagatg	cagagatatg	gatgacgcct	tttatgctgg	atattgtaac	4140
cgctgcagga	tggtcgaacg	tggaaatagct	gcacaacctc	caatctgtac	tacaatcaac	4200
catgagaaca	tgtaaaaaat	ggaggggaaa	aaggccacag	tgaataaact	cgagaattca	4260
tttattttga	aatcatgtaa	aataagatat	tagacttttt	tccctcccc	ccccctaacg	4320
cgcgctcgagc	atgcatctag	ggcgccaat	tccgcccctc	tgtctatatg	tgattttcca	4380
ttactggccg	aagccgcttg	gaataaggcc	ggtgtgcgtt	tggccctgtc	ttcttgacga	4440
ccatattgcc	gtcttttggc	aatgtgaggg	cccggaaacc	aggtctgttg	aatgtcgtga	4500
gcattcctag	gggtctttcc	cctctcgcca	aaggaatgca	gtctgtagcg	accctttgca	4560
aggaagcagt	tcctctggaa	gcttcttgaa	gacaaacaac	ccaaaagcca	cgtgtataag	4620
gtacgcggaa	ccccccacct	ggcgacaggt	gcctctgcgg	gagttggata	gttgtggaaa	4680
atacacttcg	aaaggcggca	caacccaggt	gccacgttgt	gaaggatgcc	cagaagggtac	4740
gagtcaaatg	gctctcctca	agcgtattca	acaaggggct	ctttacatgt	gtttagtcga	4800
cccattgtat	gggatctgat	ctggggcctc	ggtgcacatg	ggttttcctt	tgaaaaacac	4860
ggttaaaaaa	acgtctaggc	ccccgaacc	acggggacgt	ctccaggctt	tggttttcgc	4920
gatgataagc	ttgcacaaac	ccgggatcct	ctatgacaag	ctggccctgg	cctcagaact	4980
tgctgctggc	ggcagcgttc	gcaggacggg	cgacggccct	ctttcaattc	cagtacgatg	5040
tccaaacctc	cgaccagcgc	tacgtccttt	acccgaacaa	ggccttcag	cgctatcggt	5100
tcagctcggc	cgcgcagccc	ggctgctcag	tcctcgacga			5160

acctgctttt	cggttccggg	tcttggtccc	gtccttacct	cacagggaaa	cggcatacac	5220
tggagaagaa	tgtgttggtt	gtctctgtag	tcacacctgg	atgtaaccag	cttcctactt	5280
tggagtcagt	ggagaattat	accctgacca	taaatgatga	ccagtggtta	ctcctctctg	5340
agactgtctg	gggagctctc	cgaggtctgg	agacttttag	ccagcttggt	tggaaatctg	5400
ctgagggcac	attctttatc	aacaagactg	agattgagga	ctttccccgc	tttcctcacc	5460
ggggcttgct	gttgataaca	tctcgccatt	acctgccact	ctctagcatc	ctggacactc	5520
tggatgtcat	ggcgtacaat	aaattgaacg	tgttccactg	gcatctggta	gatgatcctt	5580
ccttcccata	tgagagcttc	acttttccag	agctcatgag	aaaggggtcc	tacaaccctg	5640
tcacccacat	ctacacagca	caggatgtga	aggaggtcat	tgaatacgca	cggctccggg	5700
gtatccgtgt	gcttgacagag	tttgacactc	ctggccacac	tttgtcctgg	ggaccaggta	5760
tccttggtatt	actgactcct	tgctactctg	ggtctgagcc	ctctggcacc	tttggaccag	5820
tgaatcccag	tctcaataat	acctatgagt	tcatgagcac	attcttctta	gaagtcagct	5880
ctgtcttccc	agatttttat	cttcactctg	gaggagatga	ggttgatttc	acctgctgga	5940
agtccaaccc	agagatccag	gactttatga	ggagaaaagg	cttcggtgag	gacttcaagc	6000
agctggagtc	cttctacatc	cagacgctgc	tggacatcgt	ctcttcttat	ggcaagggtc	6060
atgtgggtgtg	gcaggaggtg	tttgataata	aagtaaagat	tcagccagac	acaatcatac	6120
aggtgtggcg	agaggatatt	ccagtgaact	atatgaagga	gctggaactg	gtcaccaagg	6180
ccggcttccg	ggcccttctc	tctgccccct	ggtacctgaa	ccgtatatcc	tatggccctg	6240
actggaagga	tttctacgta	gtggaacccc	tggcatttga	aggtacccct	gagcagaagg	6300
ctctggtgat	tgggtgagag	gcttgtatgt	ggggagaata	tgtggacaac	acaaacctgg	6360
tccccagggt	ctggcccaga	gcaggggctg	ttggcgaag	gctgtggagc	aacaagttga	6420
catctgacct	tatgaacgtt	tgtcacactt	ccgctgtgag	ttgctgaggc	gaacagacct	6480
gaggtgtcca	ggcccaaccc	ctcaatgtag	gcttctgtga	gcaggagttt	ctcgagtcta	6540
gaagagtcca	cccgggcggc	cgcttccctt	tagtgagggt	taatgaagg	ctcgagtcta	6600
gagggcccg	ggttcgaagg	taagcctatc	cctaaccctc	tcctcggtct	cgattctacg	6660
cgtaccgggt	agtaatgagt	ttggaattaa	ttctgtggaa	tgtgtgtcag	ttaggggtgtg	6720
gaaagtcccc	aggctcccca	ggcaggcaga	agtatgcaaa	gcatgcatct	caattagtca	6780
gcaaccagggt	gtggaagtc	cccaggctcc	ccagcaggca	gaagtatgca	aagcatgcat	6840
ctcaattagt	cagcaaccat	agtcgccccc	ctaactccgc	ccatcccgcc	cctaactccg	6900
cccagttccg	cccattctcc	gccccatggc	tgactaattt	tttttattta	tgcagaggcc	6960
gaggccgct	ctgcctctga	gctattccag	aagtagtgag	gaggcttttt	tggaggccta	7020
ggcttttgca	aaaagctccc	gggagcttgt	atatccattt	tcggatctga	tcagcacgtg	7080
ttgacaatta	atcatcgcca	tagtatatcg	gcatagtata	atacgacaag	gtgaggaaact	7140
aaaccatggc	caagcctttg	tctcaagaag	aatccaccct	cattgaaaga	gcaacggcta	7200
caatcaacag	catccccatc	tctgaagact	acagcgtcgc	cagcgcagct	ctctctagcg	7260
acggccgcag	cttcactggt	gtcaatgtat	atcattttac	tgggggacct	tgtgcagaac	7320
tcgtgggtgct	gggcaactgct	gtgctgctgc	cagctggcaa	cctgacttgt	atcgtcgcga	7380
tcggaaatga	gaacaggggc	atcttgagcc	cctgcggacg	gtgccgacag	gtgcttctcg	7440
atctgcatcc	tgggatcaaa	gccatagtga	aggacagtga	tggacagccg	acggcagttg	7500
ggattcgtga	attgctgccc	tctggttatg	tgtgggaggg	ctaagcacia	ttcgagctcg	7560
gtacctttaa	gaccaatgac	ttacaaggca	gctgtagatc	ttagccactt	tttaaaagaa	7620
aaggggggac	tgggaagggt	aattcactcc	caacgaagac	aagatctgct	ttttgcttgt	7680
actgggtctc	tctggttaga	ccagatctga	gcctgggagc	tctctggcta	actagggaac	7740
ccactgctta	agcctcaata	aagcttgcc	tgagtgtctc	aagtagtggt	tgcccgtctg	7800
ttgtgtgact	ctggtaacta	gagatccctc	agaccctttt	agtcagtgtg	gaaaatctct	7860
agcagtagta	gttcagtca	tcttattatt	cagtatttat	aacttgcaaa	gaaatgaata	7920
tcagagagtg	agaggaaact	gtttattgca	gcttataatg	gttacaaata	aagcaatagc	7980
atcacaaatt	tcacaaataa	agcatttttt	tactgcat	ctagtgtgtg	tttgtccaaa	8040
ctcatcaatg	tatcttatca	tgtctggctc	tagctatccc	gcccctaact	ccgcccattc	8100
cgcccctaag	tccgcccagt	tccgcccatt	ctccgccc	tggctgacta	atttttttta	8160
tttatgcaga	ggccgagggc	gcctcgccct	ctgagctatt	ccagaagtag	tgaggaggct	8220
tttttgagg	cctagggacg	tacccaattc	gccctatagt	gagtcgtatt	acgcgcgctc	8280
actggccgtc	gttttacaac	gtcgtgactg	ggaaaaccct	ggcgttacc	aacttaatcg	8340
ccttgacaga	catccccctt	tcgccagctg	gcgtaatagc	gaagaggccc	gcaccgatcg	8400
cccttcccaa	cagttgcgca	gcctgaatgg	cgaatgggac	gcgccctgta	gcggcgcat	8460
aagcgcggcg	ggtgtggtgg	ttacgcgcag	cgtgaccgct	acacttgcca	gcgccctagc	8520
gcccgtctct	ttcgtttctt	tcccttccct	tctcgccacg	ttcgccggct	ttccccgtca	8580
agctctaaat	cgggggctcc	ctttagggtt	ccgatttagt	gctttacggc	acctcgacct	8640
caaaaaactt	gattagggtg	atggttcacg	tagtgggcca	tcgccctgat	agacgggttt	8700

tcgccctttg	acgttggagt	ccacgttctt	taatagtga	ctcttggtcc	aaactggaac	8760
aacactcaac	cctatctcgg	tctattcttt	tgatttataa	gggattttgc	cgatttcggc	8820
ctattgggta	aaaaatgagc	tgatttaaca	aaaatttaac	gcgaatttta	acaaaatatt	8880
aacgcttaca	atthaggtgg	cacttttcgg	ggaaatgtgc	gcggaacccc	tatttggtta	8940
tttttctaaa	tacattcaaa	tatgtatccg	ctcatgagac	aataaccctg	ataaatgctt	9000
caataatatt	gaaaaaggaa	gagtatgagt	attcaacatt	tccgtgtcgc	ccttattccc	9060
ttttttgcgg	cattttgcct	tcctgttttt	gctcaccag	aaacgctggt	gaaagtaaaa	9120
gatgctgaag	atcagttggg	tgacagagt	ggttacatcg	aactggatct	caacagcggg	9180
aagatccttg	agagttttcg	ccccgaagaa	cgttttccaa	tgatgagcac	ttttaaagtt	9240
ctgctatgtg	gcgcgggtatt	atcccgtatt	gacgcggggc	aagagcaact	cggtcgccgc	9300
atacactatt	ctcagaatga	cttggttgag	tactcaccag	tcacagaaaa	gcattcttacg	9360
gatggcatga	cagtaagaga	attatgcagt	gctgccataa	ccatgagtga	taacactgcg	9420
gccaaacttac	ttctgacaac	gatcggagga	ccgaaggagc	taaccgcttt	tttgacaaac	9480
atggggggtc	atgtaactcg	ccttgatcgt	tgggaaccgg	agctgaatga	agccatacca	9540
aacgacgagc	gtgacaccac	gatgcctgta	gcaatggcaa	caacgttgcg	caaactatta	9600
actggcgaaac	tacttactct	agcttcccgg	caacaattaa	tagactggat	ggaggcggat	9660
aaagtgtcag	gaccacttct	gcgctcggcc	cttcgggctg	gctgggttat	tgctgataaa	9720
tctggagccg	gtgagcgtgg	gtctcgcggt	atcattgcag	cactggggcc	agatggtaag	9780
ccctcccgtg	tcgtagttat	ctacacgacg	gggagtcagg	caactatgga	tgaacgaaat	9840
agacagatcg	ctgagatagg	tgccctactg	attaagcatt	ggtaactgtc	agaccaagtt	9900
tactcatata	tacttttagat	tgatttaaaa	cttcattttt	aattttaaag	gatctagggtg	9960
aagatccttt	ttgataatct	catgaccaaa	atcccttaac	gtgagttttc	gttccactga	10020
gcgtcagacc	ccgtagaaaa	gatcaaagga	tcttcttgag	atcctttttt	tctgcgcgta	10080
atctgctgct	tgcaaaacaaa	aaaaccaccg	ctaccagcgg	tggtttgttt	gccggatcaa	10140
gagctaccaa	ctctttttcc	gaaggttaact	ggcttcagca	gagcgcagat	accaaatact	10200
gttcttctag	tgtagccgta	gttaggccac	cacttcaaga	actctgtagc	accgcctaca	10260
tacctcgctc	tgctaatact	gttaccagt	gctgctgcca	gtggcgataa	gtcgtgtctt	10320
accgggttgg	actcaagacg	atagttaccg	gataaggcgc	agcggtcggg	ctgaacgggg	10380
ggttcgtgca	cacagcccag	cttgagcga	acgacctaca	ccgaactgag	atacctacag	10440
cgtgagctat	gagaaagcgc	cacgcttccc	gaaggagaa	aggcggacag	gtatccggta	10500
agcggcaggg	tcggaacagg	agagcgcacg	aggagccttc	cagggggaaa	cgctcggtat	10560
ctttatagtc	ctgtcggggt	tcgccacctc	tgacttgagc	gtcgattttt	gtgatgctcg	10620
tcaggggggc	ggagcctatg	gaaaaacgcc	agcaacgcgg	ccttttttacg	gttcctggcc	10680
ttttgctggc	cttttgctca	catgttcttt	cctgcgttat	cccctgattc	tgtggataac	10740
cgtattaccg	cctttgagt	agctgatacc	gctcgcgcga	gccgaacgac	cgagcgcagc	10800
gagtcagtga	gcgaggaagc	ggaagagcgc	ccaatacgca	aaccgcctct	ccccgcgcgt	10860
tgcccgattc	attaatgcag	ctggcacgac	aggtttcctg	actggaaagc	gggcagtga	10920
cgcaacgcaa	ttaatgtgag	ttagctcact	cattaggcac	cccaggcttt	acactttatg	10980
cttcgcggctc	gtatgttgtg	tgggaattgtg	agcggataac	aatttcacac	aggaaacagc	11040
tatgaccatg	attacgccaa	gcgcgcgaatt	aaccctcact	aaagggaaca	aaagctggag	11100
ctgcaagctt						11110

&lt;210&gt; 21

&lt;211&gt; 1278

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 21

tcgaggtgag	ccccacgttc	tgtttcactc	tcccatctc	ccccccctcc	ccacccccaa	60
ttttgtattt	atttattttt	taattatttt	gtgcagcgat	gggggcgggg	gggggggggg	120
cgcgcgccag	gcggggcggg	gcggggcgag	gggcggggcg	gggcgaggcg	gagaggtgag	180
gcggcagcca	atcagagcgg	cgcgctccga	aagtttctct	ttatggcgag	gcggcgcgcg	240
cggcgccctt	ataaaaaagc	aagcgcgcgg	cgggcgggag	tcgctgcgtt	gccttcgccc	300

cgtgccccgc	tccgcgcgcg	ctcgcgcgcg	ccgccccggc	tctgactgac	cgcgttactc	360
ccacaggtga	gcgggcggga	cgcccttct	cctccgggct	gtaattagcg	cttggtttaa	420
tgacggctcg	tttcttttct	gtggctgcgt	gaaagcctta	aagggtccg	ggagggccct	480
ttgtgcgggg	gggagcggt	cggggggtgc	gtgcgtgtgt	gtgtgcgtgg	ggagcgccgc	540
gtgcggcccg	cgctgcccgg	cgctgtgag	cgctgcgggc	gcggcgcggg	gctttgtgcg	600
ctccgcgtgt	gcgcgagggg	agcgcggccg	ggggcggtgc	cccgcggtgc	gggggggctg	660
cgaggggaac	aaaggctgcg	tgcggggtgt	gtgcgtgggg	gggtgagcag	gggtgtggg	720
cgcggcggtc	gggtgtaac	ccccctgc	acccctcc	ccgagttgct	gagcacggcc	780
cggcttcggg	tgcggggctc	cgtgcggggc	gtggcgcggg	gctcgccgtg	ccgggcgggg	840
ggtggcgcca	ggtgggggtg	ccggcgggg	cgggcccgcc	tcgggcccgg	gagggctcgg	900
gggaggggag	cggcgccccc	ggagcgccgg	cggtgtcga	ggcgcgcgca	gccgcagcca	960
ttgcctttta	tggtaatcgt	gcgagagggc	gcagggaactt	cctttgtccc	aaatctggcg	1020
gagccgaaat	ctgggagggc	ccgccgcacc	ccctctagcg	ggcgcgggcg	aagcgggtgcg	1080
gcgcggcgag	gaaggaaatg	ggcggggagg	gccttcgtgc	gtcgccgcgc	cgcggtcccc	1140
ttctccatct	ccagcctcgg	ggctgccgca	gggggacggc	tgccctcggg	ggggacgggg	1200
cagggcgggg	ttcggcttct	ggcgtgtgac	cggcgggggtt	tatatcttcc	cttctctgtt	1260
cctccgcagc	cagccatg					1278

&lt;210&gt; 22

&lt;211&gt; 1278

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 22

tcgaggtgag	ccccacgttc	tgettactc	tccccatctc	ccccccctcc	ccacccccaa	60
ttttgtattt	atttattttt	taattatttt	gtgcagcgat	gggggcgggg	gggggggggg	120
cgcgcgccag	gcggggcggg	gcggggcgag	gggcggggcg	gggcgaggcg	gagaggtgcg	180
gcggcagcca	atcagagcgg	cgcgtccga	aagtttctt	ttatggcgag	gcggcgggcg	240
cggcgccct	ataaaaagcg	aagcgcgcgg	cgggcgggag	tcgctgcgtt	gccttcgccc	300
cgtgccccgc	tccgcgcgcg	ctcgcgcgcg	ccgccccggc	tctgactgac	cgcgttactc	360
ccacaggtga	gcgggcggga	cgcccttct	cctccgggct	gtaattagcg	cttggtttaa	420
tgacggctcg	tttcttttct	gtggctgcgt	gaaagcctta	aagggtccg	ggagggccct	480
ttgtgcgggg	gggagcggt	cggggggtgc	gtgcgtgtgt	gtgtgcgtgg	ggagcgccgc	540
gtgcggcccg	cgctgcccgg	cggtgtgag	cgctgcgggc	gcggcgcggg	gctttgtgcg	600
ctccgcgtgt	gcgcgagggg	agcgcggccg	ggggcggtgc	cccgcggtgc	gggggggctg	660
cgaggggaac	aaaggctgcg	tgcggggtgt	gtgcgtgggg	gggtgagcag	gggtgtggg	720
cgcggcggtc	gggtgtaac	ccccctgc	acccctcc	ccgagttgct	gcgcacggcc	780
cggcttcggg	tgcggggctc	cgtgcggggc	gtggcgcggg	gctcgccgtg	ccgggcgggg	840
ggtggcgcca	ggtgggggtg	ccgggcgggg	cgggcccgcc	tcgggcccgg	gagggctcgg	900
gggaggggag	cggcgccccc	ggagcgccgg	cggtgtcga	ggcgcgcgca	gccgcagcca	960
ttgcctttta	tggtaatcgt	gcgagagggc	gcagggaactt	cctttgtccc	aaatctggcg	1020
gagccgaaat	ctgggagggc	ccgccgcacc	ccctctagcg	ggcgcgggcg	aagcgggtgcg	1080
gcgcggcgag	gaaggaaatg	ggcggggagg	gccttcgtgc	gtcgccgcgc	cgcggtcccc	1140
ttctccatct	ccagcctcgg	ggctgccgca	gggggacggc	tgccctcggg	ggggacgggg	1200
cagggcgggg	ttcggcttct	ggcgtgtgac	cggcgggggtt	tatatcttcc	cttctctgtt	1260
cctccgcagc	cagccatg					1278

&lt;210&gt; 23

&lt;211&gt; 1729

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 23  
gaattcggta ccctagttat taatagtaat caattacggg gtcattagtt catagcccat 60  
atatggagtt ccgcgttaca taacttacgg taaatggccc gcctggctga ccgccaacg 120  
acccccgccc attgacgtca ataatgacgt atgttcccat agtaacgcca atagggactt 180  
tccattgacg tcaatgggtg gactatttac ggtaaaactgc ccacttggca gtacatcaag 240  
tgtatcatat gccaagtacg cccctattg acgtcaatga cggtaaatgg ccgcctggc 300  
attatgcccga gtacatgacc ttatgggact ttctacttg gcagtacatc tacgtattag 360  
tcatcgctat taccatggtc gaggtgagcc ccacgttctg cttcactctc cccatctccc 420  
ccccctcccc accccaatt ttgtatttat ttatttttta attattttgt gcagcgatgg 480  
gggcgggggg gggggggggg cgcgcgccag gcgggcgggg gcgggcgag gggcgggcg 540  
gggcgaggcg gagaggtgcg gcgagcgcca atcagagcgg cgcgctccga aagtttcctt 600  
ttatggcgag gcgagggcg cgcgcgccct aaaaaagcg aagcgcgcg cgggcgagg 660  
tcgctgcgac gctgccttcg ccccgcgccc cgctccgccc ccgcccggc 720  
cggtctgac tgacgcggtt actcccacag gtgagcgggc gggacggccc ttctcctccg 780  
ggctgtaatt agcgcttggg ttaatgacgg cttgtttctt ttctgtggct gcgtgaaagc 840  
cttgaggggc tccgggaggg ccctttgtgc gggggggagc ggctcgggg gtgcgtgcgt 900  
gtgtgtgtgc gtggggagcg ccgcgtgcgg ccgcgctgc ccggcggtg tgagcgctgc 960  
gggcgcgggc cggggctttg tgcgctccgc agtgtgcgag aggggagcgc ggccgggggc 1020  
ggtgccccgc ggtgcggggg gggctgcgag gggaaacaaag gctgcgtgcg ggggtgtgtc 1080  
gtgggggggt gagcaggggg tgtgggcgcg gcggtcgggc tgtaaccccc ccctgcaccc 1140  
ccctccccga gttgctgagc acggcccggc ttcggtgcg gggctccgta cggggcggtg 1200  
cgcggggctc gccgtgcgg gcggggggtg gcggcaggtg ggggtgccg gcggggcggg 1260  
gccgcctcgg gccggggagg gctcggggga ggggcgcggc ggcccccgga gcgccggcg 1320  
ctgtcgaggg gcggcgagcc gcagccattg ccttttatgg taatcgtagc agagggcgca 1380  
gggacttcct ttgtcccaaa tctgtgcgga gccgaaatct gggaggcgcc gccgcacccc 1440  
ctctagcggg cgcgggggcga agcggtagcg cgccggcagg aaggaaatgg gcggggaggg 1500  
ccttcgtgcg tcgccgcgcc gccgtcccct tctcctctc cagcctcggg gctgtccgag 1560  
gggggacggc tgccctcggg ggggacgggg cagggcgggg ttcggttctt ggcgtgtgac 1620  
cggcgctct agagcctctg ctaaccatgt tcatgccttc ttctttttcc tacagctcct 1680  
gggcaacgtg ctggttattg tgctgtctca tcattttggc aaagaattc 1729

<210> 24

<211> 366

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 24

tagttattaa tagtaatcaa ttacggggtc attagttcat agcccatata tggagttccg 60  
cgttacataa cttacggtaa atggcccgcc tggtgacgg cccaacgacc ccgcccatt 120  
gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc attgacgtca 180  
atgggtggac tatttacggt aaactgccc cttggcagta catcaagtgt atcatatgcc 240  
aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt atgccagta 300  
catgacctta tgggactttc ctacttggca gtacatctac gtattagtca tcgctattac 360  
catggt

<210> 25

<211> 1295

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 25  
 ccaattttgt atttatttat tttttaatta ttttgtgcag cgatgggggc gggggggggg 60  
 ggggggcgcg cgccaggcgg ggcggggcgg ggcgaggggc ggggcggggc gaggcggaga 120  
 ggtgcggcgg cagccaatca gagcggcggc ctccgaaagt ttccttttat ggcgagggcg 180  
 cggcggcggc ggccctataa aaagcgaagc gcgcggcggg cgggagtcgc tgcgacgctg 240  
 ccttcgcccc gtgccccgct ccgcccgcgc ctccgcccgc ccgccccggc tctgactgac 300  
 cgcgtttact ccacaggtga gcgggcggga cggcccttct cctccgggct gtaattagcg 360  
 cttggtttaa tgacggcttg tttcttttct gtggctgcgt gaaagccttg aggggctccg 420  
 ggagggccct ttgtgcgggg gggagcggct cgggggggtg gtgctgtgtg gtgtgcgtgg 480  
 ggagcgcgcg gtgcggcccc cgctgcccgg cggctgtgag cgctgcgggc gcggcgcggg 540  
 gctttgtgcg ctccgcagtg tgcgcgaggg gagcgcggcc gggggcgggt ccccgcggtg 600  
 cggggggggc tgcgagggga acaaagcgtg cgtgcggggg gtgtgcgtgg ggggggtgagc 660  
 agggggtgtg ggcgcggcgg tcgggctgta accccccctt gcacccccct ccccgagttg 720  
 ctgagcacgg cccggcttcg ggtgcggggc tccgtacggg gcgtggcgcg gggctcgcgcg 780  
 tgccggggcg ggggtggcgg caggtggggg tgcggggcgg ggcggggccg cctcggggcg 840  
 gggagggctc gggggagggg cgcggcggcc cccggagcgc cggcggctgt cgaggcgcgg 900  
 cgagccgcag ccattgcctt ttatgtaaat cgtgcgagag ggcgagggga cttcctttgt 960  
 cccaaatctg tgcggagccg aaatctggga ggcgcggccg cccccctct agcgggcgcg 1020  
 gggcgaagcg gtgcggcgcc ggcaggaagg aaatgggcgg ggagggcctt cgtgcgtcgc 1080  
 cgcgcgcgcg tcccccttc cctctccagc ctccgggctg tccgcggggg gacggctgcc 1140  
 ttcggggggg acggggcagg gcgggggtcg gcttctggcg tgtgaccggc ggctctagag 1200  
 cctctgctaa ccatgttcac gccttcttct ttttctaca gtcctggggc aacgtgctgg 1260  
 ttattgtgct gtctcatcat tttggcaaa aattc 1295

<210> 26

<211> 1278

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:/Note =  
 Synthetic Construct

<400> 26  
 tcgaggtgag cccacagttc tgcttacttc tccccatctc cccccctcc ccacccccaa 60  
 ttttgtattt atttattttt taattatttt gtgcagcgat gggggcgggg gggggggggg 120  
 gcgcgcgcag gcggggcggg gcggggcgag gggcggggcg gggcgaggcg gagaggtgcg 180  
 gcggcagcca atcagagcgg cgcgctccga aagtttctt ttatggcgag gcggcggcgg 240  
 cggcggccct ataaaaagcg aagcgcgcgg cgggcgggag tcgctgcgtt gccttcgccc 300  
 cgtgccccgc tccgcgcgcg ctccgcgcgc ccgccccggc tctgactgac cgcgttactc 360  
 ccacaggtga gcggggcggga cggcccttct cctccgggct gtaattagcg cttggtttaa 420  
 tgacggctcg tttcttttct gtggctgcgt gaaagcctta aagggtccg ggagggccct 480  
 ttgtgcgggg gggagcggct cgggggggtg gtgctgtgtg gtgtgcgtgg ggagcgcgcg 540  
 gtgcggcccc cgctgcccgg cggctgtgag cgctgcgggc gcggcgcggg gctttgtgcg 600  
 ctccgcgtgt gcgcgagggg agcgcggccg ggggcgggtg cccgcgggtg gggggggctg 660  
 cgaggggaac aaaggctgcg tgcgggggtg gtgctgtggg gggtagcag ggggtgtggg 720  
 cgcggcggtc gggctgtaac cccccctgc accccccctc ccgagttgct gcgcacggcc 780  
 cggcttcggg tgccggggctc cgtgcggggc gtggcgcggg gctcgcctg ccgggcgggg 840  
 ggtggcggca ggtgggggtg ccgggcgggg cggggccgcg tcgggcgggg gagggctcgg 900  
 gggagggggc cggcggcccc ggagcgcggc cggctgtcga ggcgcgcgga gccgcagcca 960  
 ttgcctttta tggtaatcgt gcgagaggc gcaggactt cctttgtccc aaatctggcg 1020  
 gagccgaat ctgggagggc ccgcgcacc cctctagcg ggcgcgggcg aagcgggtgcg 1080  
 gcgcggcgag gaaggaaatg ggcggggagg gccttcgtg gtccgcgcgc gccgctcccc 1140  
 ttctccatct ccagcctcgg ggctgccgca gggggacggc tgccttcggg ggggacgggg 1200  
 cagggcgggg ttcggcttct ggcgtgttac cggcgggggt tatacttcc cttctctgtt 1260  
 cctccgcagc cagccatg 1278

<210> 27

<211> 229

<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 27  
gtattagtca tcgctattac catggtgatg cggttttggc agtacatcaa tgggcgtgga 60  
tagcggtttg actcacgggg atttccaagt ctccacccca ttgacgtcaa tgggagtttg 120  
ttttggcacc aaaatcaacg ggactttcca aaatgtcgta acaactccgc cccattgacg 180  
caaatgggcg gtaggcgtgt acggtgggag gtctatataa gcagagctc 229

<210> 28  
<211> 281  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 28  
tggcattatg cccagtacat gaccttatgg gactttccta ctgggcagta catctacgta 60  
ttagtcatcg ctattaccat ggtgatgcgg ttttggcagt acatcaatgg gcgtggatag 120  
cggtttgact cacggggatt tccaagtctc caccocattg acgtcaatgg gagtttggtt 180  
tggcaccaaa atcaacggga ctttccaaaa tgtcgtaaca actccgcccc attgacgcaa 240  
atgggcggta ggcgtgtacg gtgggaggtc tatataagca g 281

<210> 29  
<211> 282  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 29  
attatgccca gtacatgacc ttatgggact ttcctacttg gcagtacatc tacgtattag 60  
tcatcgctat taccatgggtg atgcggtttt ggcagtacat caatgggagt ggatagcggg 120  
ttgactcacg gggatttcca agtctccacc ccattgacgt caatgggagt ttgttttggc 180  
accaaaatca acgggacttt ccaaaatgtc gtaacaactc cgccccattg acgcaaattg 240  
gcggtaggcg tgtacggtgg gaggtctata taagcagagc tc 282

<210> 30  
<211> 512  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 30  
ttgcgttaca taacttacgg taaatggccc gcctgggtga ccgcccacg acccccggcc 60  
attgacgtca ataatgacgt atgttcccat agtaacgcca atagggactt tccattgacg 120  
tcaatgggtg gactatttac ggtaaactgc ccacttggca gtacatcaag tgtatcatat 180

gccaaagtacg	ccccctattg	acgtcaatga	cggtaaatgg	cccgcctggc	attatgccca	240
gtacatgacc	ttatgggact	ttcctacttg	gcagtacatc	tacgtattag	tcacgcgtat	300
taccatgggtg	atgcgggtttt	ggcagtagac	caatgggcgt	ggatagcggg	ttgactcacg	360
gggatttcca	agtctccacc	ccattgacgt	caatgggagt	ttgttttggc	accaaaatca	420
acgggacttt	ccaaaatgtc	gtaacaactc	cgccccattg	acgcaaattg	gcggtagggc	480
tgtacgggtg	gaggtctata	taagcagagc	tc			512

&lt;210&gt; 31

&lt;211&gt; 308

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 31

tcggcgaagc	ctcgcgcggc	cggccaggac	gaggagcgcc	actaggttga	acatccgcac	60
gagccgcggg	gccagggtctc	ggacggggctc	tcgagactcg	atctcgtgca	tgtcggcggt	120
cgcgggtgag	gttatagacc	atctgctagg	cgggtccggg	gagacaggca	cattactggc	180
ctcgcgcgcc	agcctaggcg	tgtctagagc	tcgaccgcgc	gtccggagcg	ccattcgacc	240
ggcgggtagc	gagaagaacg	ccggagaccg	caggttataa	caacgtcatg	cataaattaa	300
gaatgggc						308

&lt;210&gt; 32

&lt;211&gt; 1848

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 32

ctgcagtga	taataaaatg	tgtgtttgtc	cgaaatacgc	gtttgagatt	tctgtcccga	60
ctaaattcat	gtcgcgcgat	agtgggtgtt	atcgccgata	gagatggcga	tattggaaaa	120
atcgatat	gaaaatatgg	catattgaaa	atgtcgccga	tgtgagtttc	tgtgtaactg	180
atatcgccat	ttttccaaaa	gttgattttt	gggcatacgc	gatatactggc	gatacgctta	240
tatcgtttac	gggggatggc	gatagacgcc	tttgggtgact	tgggcgattc	tgtgtgtcgc	300
aaatatcgca	gtttcgatat	aggtgacaga	cgatatgagg	ctatatcgcc	gatagaggcg	360
acatcaagct	ggcacatggc	caatgcata	cgatctatac	attgaatcaa	tattggccat	420
tagccatatt	attcattggg	tatatagcat	aaatcaatat	tggctattgg	ccattgcata	480
cgttgtatcc	atatcataat	atgtacattt	atattggctc	atgtccaaca	ttaccgccat	540
gttgacattg	attattgact	agttattaat	agtaatcaat	tacgggggtca	ttagtgcata	600
gcccataatat	ggagttccgc	gttacataac	ttacggtaaa	tggcccgcct	ggctgaccgc	660
ccaacgaccc	ccgcccattg	acgtcaataa	tgacgtatgt	tcccatagta	acgccaatag	720
ggactttcca	ttgacgtcaa	tgggtggagt	atttacggta	aactgcccac	ttggcagtag	780
atcaagtgtg	tcatatgcca	agtagcggcc	ctattgacgt	caatgacggg	aaatggcccg	840
cctggcatta	tgcccagtag	atgaccttat	gggactttcc	tacttggcag	tacatctacg	900
tattagtcat	cgctattacc	atgggtgatgc	ggttttggca	gtacatcaat	gggcgtggat	960
agcggtttga	ctcacgggga	tttccaagtc	tccaccccat	tgacgtcaat	gggagtttgt	1020
tttggcacca	aaatcaacgg	gactttccaa	aatgtcgtaa	caactccgcc	ccattgacgc	1080
aaatgggcgg	taggcgtgta	cgggtggagg	tctatataag	cagagctcgt	ttagtgaacc	1140
gtcagatcgc	ctggagacgc	catccacgct	gttttgacct	ccatagaaga	caccgggacc	1200
gatccagcct	ccgcggccgg	gaacgggtga	ttggaacgcg	gattccccgt	gccaagagtg	1260
acgtaagtac	cgcctataga	gtctatagc	ccaccccctt	ggcttcttag	gcatgctata	1320
ctgttttttg	cttgggtgtc	atacaccctc	gcttcctcat	gttatagggtg	atgggtatagc	1380
ttagcctata	ggtgtgggtt	attgaccatt	attgaccact	cccctattgg	tgacgatact	1440



ttccattact	aatccataac	atggctcttt	gcacaactct	ctttattggc	tatatgccaa	1500
tacactgtcc	ttcagagact	gacacggact	ctgtattttt	acaggatggg	gtctcattta	1560
ttattttacaa	attcacatat	acaacaccac	cgtccccagt	gcccgagtt	tttattaaac	1620
ataacgtggg	atctccagcg	aatctcgggt	acgtgttccg	gacatggggc	tcttctccgg	1680
tagcggcgga	gcttctacat	ccagccctgc	tcccatctc	ccactcatgg	tcctcggcag	1740
ctccttgctc	ctaacagtgg	aggccagact	taggcacagc	acgatgccca	ccaccaccag	1800
tgtgccca	aggccgtggc	ggtagggtat	gtgtctgaaa	atgagctc		1848

&lt;210&gt; 33

&lt;211&gt; 1176

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 33

cccgggcccc	gcaccccaag	gcgggccaacg	ccaaaactct	ccctcctcct	cttcctcaat	60
ctcgtctctcg	ctcttttttt	ttttcgcaaa	aggaggggag	agggggtaaa	aaaatgctgc	120
actgtgcggc	gaagccgggtg	agtgcggcgc	gcgggggccaa	tcagcgtgcg	ccgttccgaa	180
agttgccttt	tatggctcga	gcgcccgcg	cggcgcccta	taaaacccag	cggcgcgacg	240
cgccaccacc	gccgagaccg	cgtccgcccc	gcgagcacag	agcctcgctt	ttgccgatcc	300
gccgcccgtc	cacaccgccc	gccaggtaag	cccggccagc	cgaccggggc	atgcggccgc	360
ggcccccttcg	cccgtgcaga	gccgccgtct	gggccgcagc	ggggggcgca	tgggggggga	420
accggaccgc	cgtggggggc	gcgggagaag	cccctggggc	tccggagatg	ggggacacc	480
cacgccagtt	cggaggcgcg	aggccgcgct	cgggaggcgc	gctccggggg	tgccgctctc	540
ggggcggggg	caaccggcgg	ggtctttgtc	tgagccgggc	tcttgccaat	ggggatcgca	600
gggtggggcg	ggcgtagccc	ccgccaggcc	cgggtggggc	tggggcgcca	ttgccggtgc	660
gcgctggtcc	tttgggcgct	aactgcgtgc	gcgctgggaa	ttggcgctaa	ttgcgcgtgc	720
gcgctgggac	tcaaggcgct	aattgcgcgt	gcgttctggg	gcccgggggtg	ccgcggcctg	780
ggctggggcg	aaggcgggct	cggccggaag	gggtggggtc	gccgcggctc	ccgggcgctt	840
gcgcgcactt	cctgcccag	ccgctggccg	cccaggggtg	tggccgctgc	gtgcgcgcgc	900
gccgacccgg	cgctgtttga	accgggcgga	ggcggggcgtg	gcgcccgggt	gggaggggggt	960
tggggcctgg	cttctgccc	cgcgcgcgg	ggacgcctcc	gaccagtgtt	tgctttttat	1020
ggtaataacg	cggccggccc	ggcttccttt	gtccccaatc	tgggcgcgcg	ccggcgcccc	1080
ctggcgccct	aaggactcgg	cgcgcgcgaa	gtggccaggg	cgggggcgac	ttcggctcac	1140
agcgcgcccc	gctattctcg	cagctcacca	tggatg			1176

&lt;210&gt; 34

&lt;211&gt; 49

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 34

cttctggcgt	gtgaccggcg	gggtttatat	cttccttcc	caagcttgg	49
------------	------------	------------	-----------	-----------	----

&lt;210&gt; 35

&lt;211&gt; 66

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

<400> 35  
 cttctggcgt gtgaccggcg gggtttatat cttcccttct ctgttcctcc gcagcccaa 60  
 gcttgg 66

<210> 36  
 <211> 68  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 Synthetic Construct

<400> 36  
 cttctggcgt gtgaccggcg gggtttatat cttcccttct ctgttcctcc gcagccagcc 60  
 aagcttgg 68

<210> 37  
 <211> 69  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 Synthetic Construct

<400> 37  
 cttctggcgt gtgaccggcg gggtttatat cttcccttct ctgttcctcc gcagccagcc 60  
 atgcatgat 69

<210> 38  
 <211> 1278  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:/Note =  
 Synthetic Construct

<400> 38  
 tcgaggtgag cccacagttc tgcttactc tccccatctc cccccctcc ccaccccaa 60  
 ttttgtattt atttattttt taattatttt gtgcagcgat gggggcgggg gggggggggg 120  
 cgcgcgccag gcggggcggg gcggggcgag gggcgggcg gggcgaggcg gagaggtgag 180  
 gcggcagcca atcagagcgg cgcgtccga aagtttctt ttatggcgag gcggcgcgcg 240  
 cggcggccct ataaaaagcg aagcgcgcgg cggcggggag tcgctgcgtt gccttcgccc 300  
 cgtgccccgc tccgcgccgc ctgcgccgc ccgccccggc tctgactgac cgcgttactc 360  
 ccacaggtga gcgggcggga cggcccttct cctccgggct gtaattagcg cttggtttaa 420  
 tgacggctcg tttcttttct gtggctgcgt gaaagcctta aagggtccg ggagggccct 480  
 ttgtgcgggg gggagcggtc cgggggggtg gtgcgtgtgt gtgtgcgtgg ggagcggcg 540  
 gtgcggcccc cgctgccgg cggtgtgag cgtgcggggc gcggcgcggg gctttgtgag 600  
 ctccgcgtgt gcgcgagggg agcgcggccg gggcggtgc ccgcgggtgc gggggggctg 660  
 cgaggggaac aaaggtgag tgccgggtgt gtgcgtgggg gggtagagag ggggtgtggg 720  
 cgcggcggtc gggctgtaac cccccctgc accccctcc ccgagttgct gcgcacggcc 780  
 cggcttcggg tgccgggctc cgtgcggggc gtggcgcggg gctcgccgtg ccgggcgggg 840  
 ggtggcggga ggtgggggtg ccgggcgggg cggggccgcc tcgggcccgg gagggctcgg 900  
 gggagggggc cggcgccccc ggagcgccgg cggtgtcga ggcgcggcga gccgcagcca 960  
 ttgcctttta tggtaatcgt gcagagggg gcagggactt ctttgtccc aaatctggcg 1020  
 gagccgaat ctgggagggc ccgcccacc ccctctagcg ggcgcggggc aagcgtgagc 1080

gcgccggcag	gaaggaaatg	ggcggggag	gccttcgtgc	gtcgccgcgc	cgccgtcccc	1140
ttctccatct	ccagcctcgg	ggctgccgca	gggggacggc	tgcccttcggg	ggggacgggg	1200
cagggcgggg	ttcggcttct	ggcgttgatc	cggcggggtt	tatatcttcc	cttctctgtt	1260
cctccgcagc	cagccatg					1278

&lt;210&gt; 39

&lt;211&gt; 1176

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 39

cccgggccca	gcacccaag	gcggccaaag	ccaaaactct	ccctcctcct	cttcctcaat	60
ctcgtctctg	ctcttttttt	ttttcgcaaa	aggaggggag	aggggggtaa	aaaatgctgc	120
actgtgcggc	gaagccgggtg	agtgcgcggc	gcggggccaa	tcagcgtgcg	ccgttccgaa	180
agttgccttt	tatggctcga	gcggccgcgg	cggcgcccta	taaaaccag	cggcgcgacg	240
cgcaccacc	gccgagaccg	cgtccgcccc	gcgagcacag	agcctcgcct	ttgccgatecc	300
gccgcccgtc	cacacccgcc	gccaggtaa	cccggccagc	cgaccggggc	atgcggccgc	360
ggcccccttc	cccgtgcaga	gccgcctct	gggcccgcagc	ggggggcgca	tgggggggga	420
accggaccgc	cgtggggggc	gcgggagaag	cccctggggc	tccggagatg	ggggacaccc	480
cacgccagtt	cggagggcgc	aggccgcgct	cgggaggcgc	gctccggggg	tgccgctctc	540
ggggcggggg	caaccggcgc	ggtctttgtc	tgagccgggc	tcttgccaat	ggggatcgca	600
gggtgggcgc	ggcgtagccc	ccgccaggcc	cgggtggggc	tggggcgcca	ttgccggtgc	660
gcgctggtcc	tttggcgct	aactgcgtgc	gcgctgggaa	ttggcgctaa	ttgcgcgtgc	720
gcgctgggac	tcaaggcgt	aattgcgcgt	gcgttctggg	gcccgggggtg	ccgcggcctg	780
ggctggggcg	aagcggggct	cggccggaag	gggtgggggtc	gccgcggctc	ccgggcgctt	840
gcgcgcactt	cctgcccgag	ccgctggccg	cccagagggtg	tggccgctgc	gtgcgcgcgc	900
gccgaccggc	cgtgttttga	accgggcgga	ggcgggggctg	gcgcccgggt	gggaggggggt	960
tggggccttg	cttcttgccg	cgcgcgcggc	ggacgcctcc	gaccagtgtt	tgctttttat	1020
ggtaataacg	cggccggccc	ggcttctttt	gtccccaatc	tgggcgcgcg	ccggcgcccc	1080
ctggcgccct	aaggactcgg	cgcgcgcgaa	gtggccaggg	cgggggcgac	ttcggtctac	1140
agcgcgcccc	gctattctcg	cagctcacca	tggatg			1176

&lt;210&gt; 40

&lt;211&gt; 1345

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
Synthetic Construct

&lt;400&gt; 40

tcgaggtgag	ccccacgttc	tgcttcactc	tccccatctc	ccccccctcc	ccacccccaa	60
ttttgtattt	atttatTTTT	taattatTTT	gtgcagcgat	gggggcgggg	gggggggggg	120
cgcgcgccag	gcggggcggg	gcggggcgag	gggcggggcg	gggcgaggcg	gagaggtgcg	180
gcggcagcca	atcagagcgg	cgcgtccga	aagtttctct	ttatggcgag	gcggcggcgg	240
cggcgccctt	ataaaaagcg	aagcgcgcgg	cgggcgggag	tcgctgcgtt	gccttcgccc	300
cgtgccccgc	tccgcgccgc	ctcgcgccgc	ccgccccggc	tctgactgac	cgcgttactc	360
ccacaggtga	gcgggcggga	cggcccttct	cctccgggct	gtaattagcg	cttggtttaa	420
tgacggctcg	tttcttttct	gtggctgcgt	gaaagcctta	aagggtcccg	ggagggccct	480
ttgtgcgggg	gggagcggct	cggggggtgc	gtgctgtgtg	gtgtgcgtgg	ggagcgccgc	540
gtgcggcccc	cgtgccccgg	cggctgtgag	cggcgcgggc	gcggcgcggg	gctttgtgcg	600
ctccgcgtgt	gcgcgagggg	agcgcggccg	ggggcggtgc	cccgcgggtg	gggggggctg	660
cgaggggaac	aaaggctgcg	tgccgggtgt	gtgctgtggg	gggtgagcag	gggtgtgggg	720
cgcggcggtc	gggctgtaac	ccccccctgc	acccccctcc	ccgagttgct	gagcagggcc	780

cggttcggg	tgcggggctc	cgtgcggggc	gtggcgcggg	gctcgccgtg	ccgggcgggg	840
ggtggcgga	ggtgggggtg	ccgggcgggg	cggggcggcc	tcgggcgggg	gagggctcgg	900
gggagggg	cggcggcccc	ggagcgccgg	cggtgtcga	ggcgcgcgga	gccgcagcca	960
ttgcctttta	tgtaaatcgt	gagagagggc	gcagggaactt	cctttgtccc	aaatctggcg	1020
gagccgaaat	ctgggaggcg	ccgccgcacc	ccctctagcg	ggcgcgggcg	aagcgggtcg	1080
gcgcgggcag	gaaggaaatg	ggcggggagg	gccttcgtgc	gtcgccgcgc	cgccgtcccc	1140
ttctccatct	ccagcctcgg	ggctgccgca	gggggacggc	tgcttcggg	ggggacgggg	1200
cagggcgggg	ttcggttct	ggcgtgtgac	cggcggctct	agagcctctg	ctaaccatgt	1260
tcattgccttc	ttctttttcc	tacagctcct	gggcaacgtg	ctggttggtg	tgctgtctca	1320
tcattttggc	aaagaattca	agctt				1345

&lt;210&gt; 41

&lt;211&gt; 684

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:/Note =  
 Synthetic Construct

&lt;400&gt; 41

tcaatattgg	ccattagcca	tattattcat	tggttatata	gcataaatca	atattggcta	60
ttggccattg	catacgttgt	atctatatca	taatatgtac	atztatattg	gctcatgtcc	120
aatatgaccg	ccatgttggc	attgattatt	gactagttat	taatagtaat	caattacggg	180
gtcattagtt	catagcccat	atatggagtt	ccgcgttaca	taacttacgg	taaatggccc	240
gcctggctga	ccgcccaacg	acccccgcc	attgacgtca	ataatgacgt	atgttcccat	300
agtaacgcca	atagggaactt	tccattgacg	tcaatgggtg	gagtatttac	ggtaaaactgc	360
ccacttggca	gtacatcaag	tgtatcatat	gccaagtccg	ccccctattg	acgtcaatga	420
cggtaaatgg	ccgcctggc	attatgccca	gtacatgacc	ttacgggact	ttcctacttg	480
gcagtacatc	tacgtattag	tcacgctat	taccatggtg	atgcggtttt	ggcagtacac	540
caatggg	cggtgacggg	ttgactcacg	gggatttcca	agtctccacc	ccattgacgt	600
caatgggag	ttgttttggc	acaaaaatca	acgggacttt	ccaaaatgtc	gtaataaacc	660
cgccccgttg	acgcaaatgg	gcgg				684